

Quick Permutation Test (QuiPT)

Piotr Sobczyk¹, Michał Burdukiewicz²

¹Wrocław University of Technology, Institute of Mathematics and Computer Science, Poland

²University of Wrocław, Department of Genomics, Poland

Outline

- 1 n-grams
 - n-gram definition
 - Positioned n-grams
- 2 Permutation test
 - Advantages
 - Drawbacks
 - p-value resolution
- 3 QuiPT
 - Contingency tables
 - Multinomial distribution of target-feature relationship
 - Advantages over permutation test
- 4 Simulation scheme
 - Power of the test
 - False significant features
- 5 Conclusion
 - Advantages over permutation test

Outline

- 1 n-grams
 - n-gram definition
 - Positioned n-grams
- 2 Permutation test
 - Advantages
 - Drawbacks
 - p-value resolution
- 3 QuiPT
 - Contingency tables
 - Multinomial distribution of target-feature relationship
 - Advantages over permutation test
- 4 Simulation scheme
 - Power of the test
 - False significant features
- 5 Conclusion
 - Advantages over permutation test

n-grams (k-tuples) are sets of n characters derived from the input sequence(s). They may form continuous sub-sequences or be discontinuous.

	X1	X2	X3	X4	X5	X6
1	4	1	4	1	3	3
2	1	4	3	1	2	2
3	1	4	1	1	1	1

Sample sequences.

1	2	3	4
2	0	2	2
2	2	1	1
5	0	0	1

Unigrams.


X1_1_0	X2_1_0	X3_1_0	X4_1_0	X5_1_0	X6_1_0	X1_2_0
0	1	0	1	0	0	0
1	0	0	1	0	0	0
1	0	1	1	1	1	0

Part of possible unigrams with position information.

Positioned n-gram data is binary.

Number of possible positioned n-grams:

$$n_{max} = L \times m^n$$



Outline

- 1 n-grams
 - n-gram definition
 - Positioned n-grams
- 2 **Permutation test**
 - Advantages
 - Drawbacks
 - p-value resolution
- 3 QuiPT
 - Contingency tables
 - Multinomial distribution of target-feature relationship
 - Advantages over permutation test
- 4 Simulation scheme
 - Power of the test
 - False significant features
- 5 Conclusion
 - Advantages over permutation test

- 1 Calculate test statistic for the given positioned n-gram and etiquettes (T_R).

- 1 Calculate test statistic for the given positioned n-gram and etiquettes (T_R).
- 2 Permute counts of n-grams and calculate permuted test statistic (T_P).

- 1 Calculate test statistic for the given positioned n-gram and etiquettes (T_R).
- 2 Permute counts of n-grams and calculate permuted test statistic (T_P).
- 3 Repeat step 2. N times.

- 1 Calculate test statistic for the given positioned n-gram and etiquettes (T_R).
- 2 Permute counts of n-grams and calculate permuted test statistic (T_P).
- 3 Repeat step 2. N times.
- 4 Calculate p-value using:

$$\text{p-value} = \frac{N_{T_P > T_R}}{N}$$

$N_{T_P > T_R}$ is number of times when T_P was bigger than T_R

- Model independent.

- Model independent.
- Statistic independent.

- Computationally expensive (number of cases, number of features).

- Computationally expensive (number of cases, number of features).
- Single feature analysis (no feature interaction).

The number of permutations is inversely proportional to the resolution of p-value.

Example: with 10×10^6 permutation the smallest possible p-values are: 0, 1×10^{-6} , 2×10^{-6} and so on.

Outline

- 1 n-grams
 - n-gram definition
 - Positioned n-grams
- 2 Permutation test
 - Advantages
 - Drawbacks
 - p-value resolution
- 3 **QuiPT**
 - Contingency tables
 - Multinomial distribution of target-feature relationship
 - Advantages over permutation test
- 4 Simulation scheme
 - Power of the test
 - False significant features
- 5 Conclusion
 - Advantages over permutation test

The binary positioned n-gram data tabulated by binary label can be easily described in 2d contingency table.

sequence ID	feature	target
1	1	0
2	1	0
3	0	0
4	1	1
5	0	1
...

Positioned n-grams with a label.

	target	feature
0	$n_{1,1}$	$n_{1,0}$
1	$n_{0,1}$	$n_{0,0}$

Contingency table.

Test statistics used by QuiPT (information gain, Kullback-Leibler divergence) measure inbalance of contingency tables.

If probability that target equals 1 is p and probability that feature equals 1 is q and feature and target are independent then each of them has the following probabilities

$$P(\text{Target}, \text{Feature}) = (1, 1)) = p \cdot q$$

$$P(\text{Target}, \text{Feature}) = (1, 0)) = p \cdot (1 - q)$$

$$P(\text{Target}, \text{Feature}) = (0, 1)) = (1 - p) \cdot q$$

$$P(\text{Target}, \text{Feature}) = (0, 0)) = (1 - p) \cdot (1 - q)$$

$$F(n_{1,1}, n_{1,0}, n_{0,1}, n_{0,0}) = \binom{n}{n_{1,1}} (p \cdot q)^{n_{1,1}} \binom{n - n_{1,1}}{n_{1,0}} (p \cdot (1 - q))^{n_{1,0}} \\ \binom{n - n_{1,1} - n_{1,0}}{n_{0,1}} ((1 - p) \cdot q)^{n_{0,1}} \\ \binom{n - n_{1,1} - n_{1,0} - n_{0,1}}{n_{0,0}} ((1 - p) \cdot (1 - q))^{n_{0,0}}$$

In addition to this: $n_{1\cdot} = n_{1,1} + n_{1,0}$ and $n_{\cdot 1} = n_{1,1} + n_{0,1}$ are known and fixed.


- $n_{1,1}$ is from range $[0, \min(n_{\cdot,1}, n_{1,\cdot})]$.

- $n_{1,1}$ is from range $[0, \min(n_{\cdot,1}, n_{1,\cdot})]$.
- The probability of certain contingency table is given as the conditional distribution, as impose restrictions on two parameters $n_{\cdot,1}$ and $n_{1,\cdot}$.

- $n_{1,1}$ is from range $[0, \min(n_{\cdot,1}, n_{1,\cdot})]$.
- The probability of certain contingency table is given as the conditional distribution, as impose restrictions on two parameters $n_{\cdot,1}$ and $n_{1,\cdot}$.
- The test statistic is computed for each possible value of $n_{1,1}$.

- $n_{1,1}$ is from range $[0, \min(n_{.,1}, n_{1,.})]$.
- The probability of certain contingency table is given as the conditional distribution, as impose restrictions on two parameters $n_{.,1}$ and $n_{1,.}$
- The test statistic is computed for each possible value of $n_{1,1}$.
- The distribution of test statistics under hypothesis that target and feature are independant is computed using values from 3.

Information Gain Probability



	Target	Feature	Freq
1	0	0	50
2	1	0	50
3	0	1	50
4	1	1	50

- QuiPT is faster.

- QuiPT is faster.
- Using the exact distribution of possible values of the criterion QuiPT yields precise small p-values without increasing the computation time.

Outline

- 1 n-grams
 - n-gram definition
 - Positioned n-grams
- 2 Permutation test
 - Advantages
 - Drawbacks
 - p-value resolution
- 3 QuiPT
 - Contingency tables
 - Multinomial distribution of target-feature relationship
 - Advantages over permutation test
- 4 Simulation scheme
 - Power of the test
 - False significant features
- 5 Conclusion
 - Advantages over permutation test

- 1 Random 4000 sequences (20 nucleotides each). The half of the sequences has label 0.


- 1 Random 4000 sequences (20 nucleotides each). The half of the sequences has label 0.
- 2 Choose a single position between 3 and 18 (to avoid border cases).


- 1 Random 4000 sequences (20 nucleotides each). The half of the sequences has label 0.
- 2 Choose a single position between 3 and 18 (to avoid border cases).
- 3 Resample nucleotides at chosen position. The dominant nucleotide has probability of occurrence $p_d = 0.25$. Other nucleotides have probability of occurrence $p_o = (1 - p_d)/3$.

- ❶ Random 4000 sequences (20 nucleotides each). The half of the sequences has label 0.
- ❷ Choose a single position between 3 and 18 (to avoid border cases).
- ❸ Resample nucleotides at chosen position. The dominant nucleotide has probability of occurrence $p_d = 0.25$. Other nucleotides have probability of occurrence $p_o = (1 - p_d)/3$.
- ❹ Perform QuiPT (Information Gain) and choose significant features (with p-value < 0.001).

- 1 Random 4000 sequences (20 nucleotides each). The half of the sequences has label 0.
- 2 Choose a single position between 3 and 18 (to avoid border cases).
- 3 Resample nucleotides at chosen position. The dominant nucleotide has probability of occurrence $p_d = 0.25$. Other nucleotides have probability of occurrence $p_o = (1 - p_d)/3$.
- 4 Perform QuiPT (Information Gain) and choose significant features (with p-value < 0.001).
- 5 Iterate steps 1-4 over other values of p_d - 0.38, 0.51, 0.65, 0.78, 0.91.

- 1 Random 4000 sequences (20 nucleotides each). The half of the sequences has label 0.
- 2 Choose a single position between 3 and 18 (to avoid border cases).
- 3 Resample nucleotides at chosen position. The dominant nucleotide has probability of occurrence $p_d = 0.25$. Other nucleotides have probability of occurrence $p_o = (1 - p_d)/3$.
- 4 Perform QuiPT (Information Gain) and choose significant features (with p-value < 0.001).
- 5 Iterate steps 1-4 over other values of p_d - 0.38, 0.51, 0.65, 0.78, 0.91.
- 6 Repeat steps 1-5 200 times.





Outline

- 1 n-grams
 - n-gram definition
 - Positioned n-grams
- 2 Permutation test
 - Advantages
 - Drawbacks
 - p-value resolution
- 3 QuiPT
 - Contingency tables
 - Multinomial distribution of target-feature relationship
 - Advantages over permutation test
- 4 Simulation scheme
 - Power of the test
 - False significant features
- 5 Conclusion
 - Advantages over permutation test

Quick permutation test is a powerful and quick equivalent of permutation test in binary feature-binary target testing scenario.