

# Quick Permutation Test (QuiPT)

Piotr Sobczyk<sup>1</sup>, Michał Burdukiewicz<sup>2</sup>

<sup>1</sup>Wrocław University of Technology, Institute of Mathematics and Computer Science, Poland

<sup>2</sup>University of Wrocław, Department of Genomics, Poland

# Outline

- 1 n-grams
  - n-gram definition
  - Positioned n-grams
- 2 Permutation test
  - Testing framework
  - Advantages
  - Drawbacks
  - p-value resolution
- 3 QuiPT
  - Contingency tables
  - Multinomial distribution of target-feature relationship
  - Advantages over permutation test
- 4 Simulation scheme
  - Power of the test
  - False significant features
- 5 Conclusion

# Outline

- 1 n-grams
  - n-gram definition
  - Positioned n-grams
- 2 Permutation test
  - Testing framework
  - Advantages
  - Drawbacks
  - p-value resolution
- 3 QuiPT
  - Contingency tables
  - Multinomial distribution of target-feature relationship
  - Advantages over permutation test
- 4 Simulation scheme
  - Power of the test
  - False significant features
- 5 Conclusion

n-grams (k-tuples) are sets of  $n$  characters derived from the input sequence(s). They may form continuous sub-sequences or be discontinuous.

Sample sequences.

1	2	3	4
3	2	0	1
1	1	3	1
1	1	2	2

Unigrams.

X1_1_0	X2_1_0	X3_1_0	X4_1_0	X5_1_0	X6_1_0	X1_2_0
1	0	1	0	0	1	0
0	1	0	0	0	0	0
0	1	0	0	0	0	0


A fraction of possible unigrams with position information.

Positioned n-gram data is binary.

Number of possible positioned n-grams:

$$n_{max} = L \times m^n$$





# Outline

- 1 n-grams
  - n-gram definition
  - Positioned n-grams
- 2 **Permutation test**
  - Testing framework
  - Advantages
  - Drawbacks
  - p-value resolution
- 3 QuiPT
  - Contingency tables
  - Multinomial distribution of target-feature relationship
  - Advantages over permutation test
- 4 Simulation scheme
  - Power of the test
  - False significant features
- 5 Conclusion

- 1 Calculate test statistic for the given positioned n-gram and etiquettes ( $T_R$ ).

- 1 Calculate test statistic for the given positioned n-gram and etiquettes ( $T_R$ ).
- 2 Permute counts of n-grams and calculate permuted test statistic ( $T_P$ ).

- 1 Calculate test statistic for the given positioned n-gram and etiquettes ( $T_R$ ).
- 2 Permute counts of n-grams and calculate permuted test statistic ( $T_P$ ).
- 3 Repeat step 2. N times.

- 1 Calculate test statistic for the given positioned n-gram and etiquettes ( $T_R$ ).
- 2 Permute counts of n-grams and calculate permuted test statistic ( $T_P$ ).
- 3 Repeat step 2.  $N$  times.
- 4 Calculate p-value using:

$$\text{p-value} = \frac{N_{T_P > T_R}}{N}$$

$N_{T_P > T_R}$  is number of times when  $T_P$  was bigger than  $T_R$

- Model independent.

- Model independent.
- Statistic independent.



- Computationally expensive (number of cases, number of features).

- Computationally expensive (number of cases, number of features).
- Single feature analysis (no feature interaction).

- Computationally expensive (number of cases, number of features).
- Single feature analysis (no feature interaction).
- Unfeasible precise estimation of low p-values.

The number of permutations is inversely proportional to the interval between p-values.

Example: with  $10 \times 10^6$  permutation the smallest possible p-values are: 0,  $1 \times 10^{-6}$ ,  $2 \times 10^{-6}$  and so on.

# Outline

- 1 n-grams
  - n-gram definition
  - Positioned n-grams
- 2 Permutation test
  - Testing framework
  - Advantages
  - Drawbacks
  - p-value resolution
- 3 QuiPT
  - Contingency tables
  - Multinomial distribution of target-feature relationship
  - Advantages over permutation test
- 4 Simulation scheme
  - Power of the test
  - False significant features
- 5 Conclusion

The binary positioned n-gram data tabulated by binary label can be easily described in 2d contingency table.

sequence ID	feature	target
1	1	0
2	1	0
3	0	0
4	1	1
5	0	1
...	...	...

Positioned n-grams with a label.

	target	feature
0	$n_{1,1}$	$n_{1,0}$
1	$n_{0,1}$	$n_{0,0}$

Contingency table.



Test statistics used by QuiPT (information gain, Kullback-Leibler divergence) measure inbalance of contingency tables.

If probability that target equals 1 is  $p$  and probability that feature equals 1 is  $q$  and feature and target are independent then each of them has the following probabilities

$$P(\text{Target}, \text{Feature}) = (1, 1)) = p \cdot q$$

$$P(\text{Target}, \text{Feature}) = (1, 0)) = p \cdot (1 - q)$$

$$P(\text{Target}, \text{Feature}) = (0, 1)) = (1 - p) \cdot q$$

$$P(\text{Target}, \text{Feature}) = (0, 0)) = (1 - p) \cdot (1 - q)$$

$$F(n_{1,1}, n_{1,0}, n_{0,1}, n_{0,0}) = \binom{n}{n_{1,1}} (p \cdot q)^{n_{1,1}} \binom{n - n_{1,1}}{n_{1,0}} (p \cdot (1 - q))^{n_{1,0}} \\ \binom{n - n_{1,1} - n_{1,0}}{n_{0,1}} ((1 - p) \cdot q)^{n_{0,1}} \\ \binom{n - n_{1,1} - n_{1,0} - n_{0,1}}{n_{0,0}} ((1 - p) \cdot (1 - q))^{n_{0,0}}$$

In addition to this:  $n_{1\cdot} = n_{1,1} + n_{1,0}$  and  $n_{\cdot 1} = n_{1,1} + n_{0,1}$  are known and fixed.


- $n_{1,1}$  is from range  $[0, \min(n_{\cdot,1}, n_{1,\cdot})]$ .

- $n_{1,1}$  is from range  $[0, \min(n_{\cdot,1}, n_{1,\cdot})]$ .
- The probability of certain contingency table is given as the conditional distribution, as impose restrictions on two parameters  $n_{\cdot,1}$  and  $n_{1,\cdot}$ .

- $n_{1,1}$  is from range  $[0, \min(n_{.,1}, n_{1,.})]$ .
- The probability of certain contingency table is given as the conditional distribution, as impose restrictions on two parameters  $n_{.,1}$  and  $n_{1,.}$
- The test statistic is computed for each possible value of  $n_{1,1}$ .

- $n_{1,1}$  is from range  $[0, \min(n_{.,1}, n_{1,.})]$ .
- The probability of certain contingency table is given as the conditional distribution, as impose restrictions on two parameters  $n_{.,1}$  and  $n_{1,.}$
- The test statistic is computed for each possible value of  $n_{1,1}$ .
- The distribution of test statistics under hypothesis that target and feature are independant is computed using values from 3.

Information Gain    Probability



	Target	Feature	Freq
1	0	0	50
2	1	0	50
3	0	1	50
4	1	1	50



- QuiPT is faster.

- QuiPT is faster.
- Using the exact distribution of possible values of the criterion QuiPT yields precise small p-values without increasing the computation time.

# Outline

- 1 n-grams
  - n-gram definition
  - Positioned n-grams
- 2 Permutation test
  - Testing framework
  - Advantages
  - Drawbacks
  - p-value resolution
- 3 QuiPT
  - Contingency tables
  - Multinomial distribution of target-feature relationship
  - Advantages over permutation test
- 4 Simulation scheme
  - Power of the test
  - False significant features
- 5 Conclusion

- 1 Random 4000 sequences (20 nucleotides each). The half of the sequences has label 0.

- 1 Random 4000 sequences (20 nucleotides each). The half of the sequences has label 0.
- 2 Choose a single position between 3 and 18 (to avoid border cases).


- 1 Random 4000 sequences (20 nucleotides each). The half of the sequences has label 0.
- 2 Choose a single position between 3 and 18 (to avoid border cases).
- 3 Resample nucleotides at chosen position. The dominant nucleotide has probability of occurrence  $p_d = 0.25$ . Other nucleotides have probability of occurrence  $p_o = (1 - p_d)/3$ .


- ① Random 4000 sequences (20 nucleotides each). The half of the sequences has label 0.
- ② Choose a single position between 3 and 18 (to avoid border cases).
- ③ Resample nucleotides at chosen position. The dominant nucleotide has probability of occurrence  $p_d = 0.25$ . Other nucleotides have probability of occurrence  $p_o = (1 - p_d)/3$ .
- ④ Perform QuiPT (Information Gain) and choose significant features (with p-value  $< 0.001$ ).

- 1 Random 4000 sequences (20 nucleotides each). The half of the sequences has label 0.
- 2 Choose a single position between 3 and 18 (to avoid border cases).
- 3 Resample nucleotides at chosen position. The dominant nucleotide has probability of occurrence  $p_d = 0.25$ . Other nucleotides have probability of occurrence  $p_o = (1 - p_d)/3$ .
- 4 Perform QuiPT (Information Gain) and choose significant features (with p-value  $< 0.001$ ).
- 5 Iterate steps 1-4 over other values of  $p_d$  - 0.38, 0.51, 0.65, 0.78, 0.91.



- 1 Random 4000 sequences (20 nucleotides each). The half of the sequences has label 0.
- 2 Choose a single position between 3 and 18 (to avoid border cases).
- 3 Resample nucleotides at chosen position. The dominant nucleotide has probability of occurrence  $p_d = 0.25$ . Other nucleotides have probability of occurrence  $p_o = (1 - p_d)/3$ .
- 4 Perform QuiPT (Information Gain) and choose significant features (with p-value  $< 0.001$ ).
- 5 Iterate steps 1-4 over other values of  $p_d$  - 0.38, 0.51, 0.65, 0.78, 0.91.
- 6 Repeat steps 1-5 200 times.





# Outline

- 1 n-grams
  - n-gram definition
  - Positioned n-grams
- 2 Permutation test
  - Testing framework
  - Advantages
  - Drawbacks
  - p-value resolution
- 3 QuiPT
  - Contingency tables
  - Multinomial distribution of target-feature relationship
  - Advantages over permutation test
- 4 Simulation scheme
  - Power of the test
  - False significant features
- 5 Conclusion

Quick permutation test is a powerful and quick equivalent of permutation test in binary feature-binary target testing scenario.