


A mirror symmetry conjecture: *The fundamental group of the Stringy Kähler Moduli Space acts on $D^b(X)$ via spherical twists*



Michela Barbieri

The B-side: Toric Geometric Invariant Theory

We start with algebraic torus $T \cong (\mathbb{C}^*)^r$ acting on a vector space \mathbb{C}^n . These actions the actions are simple to write out:

$$(\lambda_1, \dots, \lambda_r) \cdot (z_1, \dots, z_n) = (\lambda_1^{q_{11}} \lambda_2^{q_{12}} \dots \lambda_r^{q_{1r}} z_1, \dots, \lambda_1^{q_{n1}} \lambda_2^{q_{n2}} \dots \lambda_r^{q_{nr}} z_n)$$

We get an $n \times r$ integer matrix $Q = (q_{ij})$, called the weight matrix.
We construct an algebraic variety that parametrises the actions quotients. *Geometric Invariant Theory* (GIT) is the theory that tells *the unstable locus* to throw away before we quotient so that we get a good/geometric quotient [1]. GIT quotients are often denoted $X//G$.

Example 1 Consider \mathbb{C}^* acting on \mathbb{C}^2 linearly, i.e. $\lambda \cdot (x, y) = (\lambda x, \lambda y)$. If we take the quotient space, we see that the orbit of the origin cannot be separated from any other orbit. So we have to remove the origin and as expected, we get GIT quotient

$$\mathbb{C}^2//\mathbb{C}^* = \mathbb{C}^2 \setminus \{0\} / \mathbb{C}^* = \mathbb{P}^1$$




Figure 1: Orbits of the \mathbb{C}^* action on \mathbb{C}^2 .

In general GIT quotients are not unique and depend on a choice of *stability condition* $\phi \in \mathbb{Z}^r$, where for a given stability condition we denote the GIT quotient $X//_{\phi}G$. Different choices of ϕ have us remove different unstable loci and give us non-isomorphic (but birational) quotients.

Example 2 Consider \mathbb{C}^* acting on \mathbb{C}^3 via $\lambda \cdot (x, y, z) = (\lambda x, \lambda y, \lambda^{-1} z)$. The stability conditions space is \mathbb{Z} . For $\phi > 0$, we have unstable locus $Z_+ = \{x = y = 0\}$. We have GIT quotient

$$\mathbb{C}^3//_{\phi}\mathbb{C}^* = \mathbb{C}^3 \setminus \{x = y = 0\} / \mathbb{C}^* = \theta(-1) \mathbb{P}_{x,y}^1.$$

For $\phi < 0$, we have unstable locus $Z_- = \{z = 0\}$.

$$\mathbb{C}^3//_{\phi}\mathbb{C}^* = \mathbb{C}^3 \setminus \{z = 0\} / \mathbb{C}^* = \mathbb{A}_{x,y}^1.$$

Use of Colour

With the `color` package you can use as much colour as you like: but the `Red` and `DarkBlue` colours defined here are more useful than the obvious `red` and `blue` versions, which tend to seem too bright when printed.

Use spacing as well as colour to highlight a key question or issue:

Who will rid me of this turbulent priest?

Praesent elementum malesuada mauris. Duis aliquet dolor ut nunc. Pellentesque euismod augue a odio.

Mirror Symmetry Conjecture


Of course, the principal reason for choosing `LATEX` is its ability with equations:

$$U_{\text{doublet}} = \frac{mg}{2\pi\mu a} \left(\int_0^\infty \left\{ 1 - \frac{2 \sinh^2 s - 2s^2}{\sinh 2s + 2s} \right\} ds \right)^{-1} \approx \frac{1.55mg}{6\pi\mu a},$$

but it's equally easy to include tables:

L	U_1		U_2		U_3		U_3 error
	MR	SD	MR	SD	MR	SD	
2.01	0.65528	0.64739	0.63461	0.62691	0.00498	0.00451	9%
2.10	0.73857	0.73126	0.59718	0.58784	0.03517	0.02570	27%
2.50	0.87765	0.87482	0.49545	0.48829	0.07393	0.05853	21%
3.00	0.93905	0.93806	0.41694	0.41356	0.07824	0.06970	11%
4.00	0.97964	0.97945	0.31859	0.31774	0.06925	0.06639	4%
6.00	0.99581	0.99579	0.21586	0.21575	0.05078	0.05019	1%

and, of course, images:



Sectioning

First Subsection

Aenean feugiat, mauris vitae accumsan venenatis, tortor nunc facilisis velit, vel aliquam felis dui non ante. Aenean commodo, sem vel malesuada placerat, erat lacus lacinia magna, quis euismod quam nulla vitae turpis. Vivamus sapien. Vestibulum ante ipsum primis in faucibus orci luctus et ultrices posuere cubilia Curae; Nunc a dolor. Sed diam leo, fringilla non, fermentum non, ultricies tempor, dolor. Mauris vehicula, urna ac bibendum scelerisque, enim nisl vehicula libero, id blandit enim lacus et neque. Nunc posuere elit at erat. Integer commodo, eros dapibus blandit facilisis, dolor ipsum dictum dui, sit amet iaculis enim elit sed magna. Nunc libero nunc, malesuada sit amet, tincidunt at, pharetra ut, arcu.

Second Subsection

Integer ante. Mauris tincidunt adipiscing mi. Donec sollicitudin, lacus quis pellentesque placerat, quam elit pharetra lacus, sit amet placerat nisi quam in lorem. Etiam id nulla a est vestibulum tempus. Nam et nisl non arcu venenatis semper. Duis sed libero. Cras lectus. In semper urna in leo. Sed mattis lacinia arcu. Phasellus ut metus. Phasellus mi dolor, condimentum ut, ornare nec, tempus ac, mi. Integer ante sem, vestibulum in, gravida vel, condimentum non, ipsum.

References

[1] D. Mumford, J. Fogarty, and F. Kirwan. *Geometric invariant theory*, volume 34. Springer Science & Business Media, 1994.