

Chapter 10

Classifying by a multiplicity influence

10.1 A qualitative approach by some groups

In this chapter we consider the following problem: Let X be a set of objects, and let $\mathcal{C}$ be a set of classes. We assume that X is partitioned into n groups, and that $\mathcal{C}$ is partitioned into m classes. We assume that the groups are labeled $1, \dots, n$ and the classes are labeled $1, \dots, m$. We assume that the groups are labeled $1, \dots, n$ and the classes are labeled $1, \dots, m$. We assume that the groups are labeled $1, \dots, n$ and the classes are labeled $1, \dots, m$.

A qualitative approach

Let X be a set of objects, and let $\mathcal{C}$ be a set of classes. We assume that X is partitioned into n groups, and that $\mathcal{C}$ is partitioned into m classes. We assume that the groups are labeled $1, \dots, n$ and the classes are labeled $1, \dots, m$. We assume that the groups are labeled $1, \dots, n$ and the classes are labeled $1, \dots, m$. We assume that the groups are labeled $1, \dots, n$ and the classes are labeled $1, \dots, m$.

10.2 A quantitative approach by some groups

Let X be a set of objects, and let $\mathcal{C}$ be a set of classes. We assume that X is partitioned into n groups, and that $\mathcal{C}$ is partitioned into m classes. We assume that the groups are labeled $1, \dots, n$ and the classes are labeled $1, \dots, m$. We assume that the groups are labeled $1, \dots, n$ and the classes are labeled $1, \dots, m$.

A quantitative approach

Let X be a set of objects, and let $\mathcal{C}$ be a set of classes. We assume that X is partitioned into n groups, and that $\mathcal{C}$ is partitioned into m classes. We assume that the groups are labeled $1, \dots, n$ and the classes are labeled $1, \dots, m$. We assume that the groups are labeled $1, \dots, n$ and the classes are labeled $1, \dots, m$. We assume that the groups are labeled $1, \dots, n$ and the classes are labeled $1, \dots, m$.

A quantitative approach

Let X be a set of objects, and let $\mathcal{C}$ be a set of classes. We assume that X is partitioned into n groups, and that $\mathcal{C}$ is partitioned into m classes. We assume that the groups are labeled $1, \dots, n$ and the classes are labeled $1, \dots, m$. We assume that the groups are labeled $1, \dots, n$ and the classes are labeled $1, \dots, m$. We assume that the groups are labeled $1, \dots, n$ and the classes are labeled $1, \dots, m$.

