

why? because $(U^*AU)^* = U^*A^*(U^*)^* = U^*AU$ (since $A^* = A$, if A is hermetian).

but...if T is upper-triangular, AND hermetian, it must be LOWER triangular, as well. which means when A is hermetian, T is DIAGONAL. furthermore, since the diagonal elements of T equal their own complex-conjugates, T is a REAL matrix.

observation #2: any real symmetric matrix is hermetian. thus any real symmetric matrix is unitarily similar to a real diagonal matrix, D .

observation #3: the eigenvalues of A are the eigenvalues of D , and vice-versa.

proof: $\det(xI - A) = \det(U^{-1}(xI - A)U) = \det(U^{-1}(xI)U - U^{-1}AU) = \det(xI - D)$

observation #4: since the eigenvalues of A (a real symmetric matrix) are real, the eigenvectors are likewise real. thus we may take U to be a real unitary matrix, that is, an orthogonal one.

now suppose that a real, symmetric matrix A has an eigenvalue of (algebraic) multiplicity m . since A is orthogonally similar to a diagonal matrix D with the same eigenvalues (via an orthogonal matrix Q), D has m diagonal entries that are the same.

note that $Q^{-1}AQ = D$, means $AQ = QD$. since Q is invertible, its columns are linearly independent (it's square of full rank).

suppose $\{u_{j_1}, \dots, u_{j_m}\}$ are the columns of Q corresponding to the diagonal entry λ of D repeated m times (the eigenvalue of algebraic multiplicity m). then:

$$A(u_{j_k}) = (u_{j_k})^T D = \lambda u_{j_k} \quad k = 1, \dots, m$$

giving m linearly independent eigenvectors of the eigenspace E_λ .


Reply With Quote

Opalg

MHB Oldtimer

MHB Moderator

MHB Math Helper



Status: Offline

Join Date: Feb 2012

Location: Leeds, UK

Posts: 1,480

Thanks: 481 times

Thanked: 4510 times

Awards:

DECEMBER 23RD, 2012, 12:21

#3

Originally Posted by matqkks

I have been trying to prove the following result:
If A is real symmetric matrix with an eigenvalue λ of multiplicity m then λ has m linearly independent e.vectors.
Is there a simple proof of this result?

This is a slight variation of Deveno's argument. I will assume you already know that the eigenvalues of a real symmetric matrix are all real.

Let A be an $n \times n$ real symmetric matrix, and assume as an inductive hypothesis that all $(n-1) \times (n-1)$ real symmetric matrices are diagonalisable. Let λ be an eigenvalue of A , with a normalised eigenvector x_1 . Using the Gram-Schmidt process, form an orthonormal basis $\{x_1, x_2, \dots, x_n\}$ with that eigenvector as its first element.

Let P be the $n \times n$ matrix whose columns are x_1, x_2, \dots, x_n , and denote by $T: \mathbb{R}^n \rightarrow \mathbb{R}^n$ the linear transformation whose matrix with respect to the standard basis is A . Then P is an orthogonal matrix ($P^T = P^{-1}$), and the matrix of T with respect to the basis $\{x_1, x_2, \dots, x_n\}$ is $P^T A P$. The (i, j) -element of that matrix is $(P^T A P)_{ij} = \langle A x_j, x_i \rangle$. In particular, the elements in the first column are

$$(P^T A P)_{i1} = \langle A x_1, x_i \rangle = \langle \lambda x_1, x_i \rangle = \begin{cases} \lambda & (i = 1) \\ 0 & (i > 1) \end{cases}$$

(because the vectors x_i are orthonormal). Thus the first column of $P^T A P$ has λ as its top element, and 0 for each of the other elements. Since $P^T A P$ is symmetric, the top row also consists of a λ followed by all zeros. Hence the matrix $P^T A P$ looks like this:

$$\begin{bmatrix} \lambda & 0 & \dots & 0 \\ 0 & & & \\ \vdots & & B & \\ 0 & & & \end{bmatrix},$$

where B is an $(n-1) \times (n-1)$ real symmetric matrix. By the inductive hypothesis, B is diagonalisable, so there is an orthogonal $(n-1) \times (n-1)$ matrix Q such that $Q^T B Q$ is

diagonal. Let

$$R = \begin{bmatrix} 1 & 0 & \dots & 0 \\ 0 & & & \\ \vdots & & Q & \\ 0 & & & \end{bmatrix}.$$

Then $R^T P^T A P R$ is diagonal, as required.

Was sich überhaupt sagen lässt, lässt sich klar sagen; und wovon man nicht reden kann, darüber muss man schweigen.

(Anything that can be said at all, can be said clearly; and whereof one cannot speak, thereon one must be silent.)

– Ludwig Wittgenstein's good advice for forum contributors, in *Tractatus Logico-Philosophicus*.

[Reply With Quote](#)

[▲ Go to First Post](#)

« Singular Value decomposition | An inverse of the adjoint »

Similar Threads

[SOLVED] Incorporate axis symmetric case

By dwsmith in forum Mathematics Software and Calculator Discussion

Replies: 0

Last Post: December 6th, 2012, 14:56

Inverse of a Symmetric Matrix

By OhMyMarkov in forum Linear and Abstract Algebra

Replies: 2

Last Post: October 6th, 2012, 11:28

[SOLVED] Eigenvalues

By Sudharaka in forum Pre-Algebra and Algebra

Replies: 2

Last Post: May 27th, 2012, 09:02

Matrix Theory...showing that matrix is Unitary

By cylv89 in forum Linear and Abstract Algebra

Replies: 3

Last Post: March 27th, 2012, 17:00

A and B are two symmetric matrices

By Yankel in forum Linear and Abstract Algebra

Replies: 4

Last Post: January 27th, 2012, 09:17

-- Math Help Boards

Have a small screen? Select Math Help Boards for Small Screens!

Powered by vBulletin Copyright © 2000 - 2012, Jelsoft Enterprises Ltd.

Search Engine Optimisation provided by DragonByte SEO v1.0.15 (Pro) - vBulletin Mods & Addons Copyright © 2014 DragonByte Technologies Ltd.

Feedback Buttons provided by Advanced Post Thanks / Like (Pro) - vBulletin Mods & Addons Copyright © 2014 DragonByte Technologies Ltd.

© 2012-2014 Math Help Boards

[FORUMS](#)

[RULES](#)

[POTW](#)

[CONTACT](#)

[TOP](#)

