

# Photons? Waves? Rays?

Wave superposition

Ray optics

Huygens

Snell's law

TIR

Refraction

Apparent depth

Magnification

Lenses

Curved refractive  
surfaces

Lensmaker's equation

- You've had PHY 251, so you know all about photons from Planck and Einstein. Light comes in discrete packets of energy  $E = hc/\lambda$  and momentum  $p = Ec$ .
- But then we stepped back a few decades, and learned from Maxwell that light can be described in terms of electromagnetic waves.
- But if we step back even further, physicists described light propagation in terms of ray optics. What's with *that*?
- How do waves travel in straight lines? The answer: they only kinda sorta do.
- But before we address this question, let's consider some issues of adding up waves.

# Adding waves

Wave superposition

Ray optics

Huygens

Snell's law

TIR

Refraction

Apparent depth

Magnification

Lenses

Curved refractive  
surfaces

Lensmaker's equation

- Back in Lecture 2, we learned a few things about adding waves up. A movie to illustrate how phase affects wave addition is [here](#).
- If we add  $N$  waves  $|A|e^{i\omega+\theta_0}$  with identical magnitudes  $|A|$  and frequencies  $\omega$  but random phase offsets  $\theta_0$  (*incoherent* superposition), we found a net or resultant magnitude  $|R|$  of

$$|R|_{\text{incoherent}} = \sqrt{N \cdot A_i^2} = \sqrt{N}|A|. \quad (1)$$


- If all the waves have identical frequencies and identical phase offsets (*coherent* superposition), we found

$$|R|_{\text{coherent}} = \sqrt{R^2} = \sqrt{N^2 A^2} = N|A|. \quad (2)$$

- What about intermediate cases, where the phases are not completely random but are not absolutely identical either?

- Let's now consider the addition of waves where the phase is neither uniform, nor completely random.
- Instead, let's consider phases that vary uniformly about zero over a uniform range.
- How is the sum of many waves affected by the  $\pm$  range of starting phases? See at right.
- We see that if the phase is within about  $\pm\lambda/4$ , the net effect is only moderately reduced. This is known as the *Rayleigh quarter wave criterion*.

## Adding waves II



John William Strutt, 3<sup>rd</sup> Baron Rayleigh (1842–1919)

# Rays and Fermat's principle

Wave superposition

Ray optics

Huygens

Snell's law

TIR

Refraction

Apparent depth


Magnification

Lenses

Curved refractive  
surfaces

Lensmaker's equation


- Consider light traveling from point A to point B. If we consider point A to be emitting spherical waves, why do we draw a straight line ray path to point B?



- Let's think about two simple cases: a wave traveling a distance  $z$  along a straight line (path A), and a wave traveling through a point  $y$  off of the straight line (path B).
- In each case the wave propagates according to  $e^{-i\vec{k}\cdot\vec{x}}$ , where  $\vec{k}$  diverges to all directions in the case of a spherical wave, while  $\vec{x}$  describes the vector path.

## Rays and Fermat II

- Again, we're considering the straight line path A versus the off-axis path B in  $e^{-in\vec{k}\cdot\vec{x}}$ .




- Let's handle the path by saying that we have spherical waves (all directions in  $\vec{k}$ ) and a distance  $\ell$  along a particular path. That is,  $n\vec{k} \cdot \vec{x} \Rightarrow nk\ell$  where  $\ell$  measures the distance along a path and  $n$  is the refractive index, telling us the speed of wave front propagation.
- We refer to  $n\ell$  as the *optical path length*, and  $nk\ell$  as the accumulated optical phase.
- For the straight line path A, the distance traveled is  $\ell_A = z$ .
- For the off-axis path B, the distance traveled is

$$\ell_B = 2\sqrt{\left(\frac{z}{2}\right)^2 + y^2} = z\sqrt{1 + \left(\frac{2y}{z}\right)^2} \simeq z\left(1 + 2\left(\frac{y}{z}\right)^2\right)$$

where we have made use of the binomial expansion.

## Rays and Fermat III

- Again, we're considering the straight line path A versus the off-axis path B in  $e^{-i\vec{k}\cdot\vec{x}}$ .



- Path A has an optical path length of  $n\ell_A = nz$ .
- Path B has an optical path length of  $n\ell_B \simeq nz(1 + 2y^2/z^2)$
- These paths, plus those “bounded inside,” reinforce each other if their net optical path lengths are within  $\pm\lambda/4$ :

$$\begin{aligned}\ell_B - \ell_A &\simeq z \left( 1 + 2\left(\frac{y}{z}\right)^2 \right) - z \leq \frac{\lambda}{4} \\ \frac{2y^2}{z} &\leq \frac{\lambda}{4} \\ |y| &\leq \sqrt{\frac{\lambda z}{8}}\end{aligned}\quad (3)$$

## Rays and Fermat IV

Wave superposition

Ray optics

Huygens

Snell's law

TIR

Refraction

Apparent depth



Magnification

Lenses

Curved refractive  
surfaces

Lensmaker's equation

- From Eq. 3 we found that paths with  $|y| \leq \sqrt{\lambda z/8}$  satisfy the *Rayleigh quarter wave criterion*: they add up nearly coherently.
- Consider  $\lambda = 532$  nm (green laser pointer), and  $z = 5$  meters: this says  $|y| \leq \sqrt{(532 \times 10^{-9})(5)/8}$  or  $|y| \leq 0.58$  mm.
- So waves that travel within a millimeter of the straight line path satisfy the quarter-wave criterion, and add up nearly coherently.
- With  $z = 1$  mm, we have  $|y| \leq 8$   $\mu\text{m}$ .



# Rays and Fermat V

Wave superposition

Ray optics

Huygens

Snell's law

TIR

Refraction

Apparent depth

Magnification

Lenses

Curved refractive  
surfaces

Lensmaker's equation

- By the way, the Rayleigh quarter wave criterion is related to the method of stationary phase in complex integrals.
- Also, we'll see later on that we can do the “adding up waves” calculation using Fourier transforms, and find that the pattern we calculated is equivalent to the diffraction pattern of a rectangular aperture:  $[\sin(x)/x]^2$ .
- And finally, what we have shown is Fermat's principle, which is that light travels the path of least time (because that's the minimum optical path length  $n\ell$  or minimum phase  $nk\ell$  path, which does not get canceled out by other paths).



# The Huygens construction

Wave superposition

Ray optics

**Huygens**

Snell's law

TIR

Refraction


Apparent depth

Magnification

Lenses

Curved refractive  
surfaces

Lensmaker's equation



Christiaan Huy-  
gens (1629-1695)

# The Huygens construction

Wave superposition

Ray optics

**Huygens**

Snell's law

TIR

Refraction


Apparent depth

Magnification


Lenses

Curved refractive  
surfaces

Lensmaker's equation



Christiaan Huygens (1629-1695)



# The Huygens construction

Wave superposition

Ray optics

**Huygens**

Snell's law

TIR

Refraction


Apparent depth

Magnification


Lenses

Curved refractive  
surfaces

Lensmaker's equation



Christiaan Huy-  
gens (1629-1695)



# Huygens II

Wave superposition

Ray optics

**Huygens**

Snell's law

TIR

Refraction

Apparent depth

Magnification

Lenses

Curved refractive  
surfaces

Lensmaker's equation

- The Huygens construction shows us that we can make a plane wave from lots of little spherical wave point sources.
- A plane wave travels in a particular direction, like a ray does. . .
- So we have yet another way of justifying light traveling as rays!

# Snell's law

Wave superposition

Ray optics

Huygens

Snell's law

TIR

Refraction

Apparent depth


Magnification

Lenses

Curved refractive  
surfaces

Lensmaker's equation

- Apply Fermat's principle to a ray reaching an interface between two different refractive indices  $n_1$  and  $n_2$ .
- This was done by several people, including Ibn Sahl in 984, Thomas Harriot in 1602, the Dutch mathematician Willebrord Snellius in 1621, René Descartes in 1637. . . Most of them before Fermat's principle!
- We know it today as Snell's law:  $n_1 \sin \theta_1 = n_2 \sin \theta_2$ .



We use a darker shading for  $n_2$  to indicate  $n_2 > n_1$ .

# Deriving Snell's law

Wave superposition

Ray optics

Huygens

Snell's law

TIR

Refraction

Apparent depth

Magnification

Lenses


Curved refractive surfaces

Lensmaker's equation

We have shown that light travels according to the principle of stationary phase, or least time. Let's consider how light goes from point  $P_1$  to  $P_2$  as it travels across a refractive boundary. The time is distance over velocity, or

$$t = \frac{l_1}{v_1} + \frac{l_2}{v_2} = \frac{l_1 n_1}{c} + \frac{l_2 n_2}{c} \quad (4)$$

where we have used the phase velocity  $v_p = c/n$ . Light will travel along a straight line within one medium, but what about at an interface? What value of  $b$  minimizes the travel time/produces a path of stationary phase?



## Snell's law II

Wave superposition

Ray optics

Huygens

Snell's law

TIR

Refraction

Apparent depth

Magnification

Lenses

Curved refractive surfaces

Lensmaker's equation


To find the path of least time, let's write the lengths  $l_1$  and  $l_2$  in terms of  $b$ :

$$l_1 = \sqrt{y_1^2 + b^2} \quad l_2 = \sqrt{y_2^2 + (a - b)^2}$$

The travel time of Eq. 4 then becomes

$$t = \frac{n_1[y_1^2 + b^2]^{1/2}}{c} + \frac{n_2[y_2^2 + (a - b)^2]^{1/2}}{c} \quad (5)$$

To minimize the time, let's set the derivative as we vary  $b$  equal to zero:  $dt/db = 0$ .



## Snell's law III

Wave superposition

Ray optics

Huygens

Snell's law

TIR

Refraction

Apparent depth

Magnification

Lenses

Curved refractive  
surfaces

Lensmaker's equation

Again, we want to minimize the time of Eq. 5 of

$$t = \frac{n_1[y_1^2 + b^2]^{1/2}}{c} + \frac{n_2[y_2^2 + (a - b)^2]^{1/2}}{c}$$

by setting  $dt/db = 0$ :

$$\begin{aligned}\frac{dt}{db} = 0 &= \frac{1}{2} \frac{n_1[y_1^2 + b^2]^{-1/2}}{c} 2b + \frac{1}{2} \frac{n_2[y_2^2 + (a - b)^2]^{-1/2}}{c} 2(a - b)(-1) \\ 0 &= n_1 \frac{b}{\sqrt{y_1^2 + b^2}} - n_2 \frac{a - b}{\sqrt{y_2^2 + (a - b)^2}}\end{aligned}\quad (6)$$



## Snell's law IV

Wave superposition

Ray optics

Huygens

Snell's law

TIR

Refraction

Apparent depth

Magnification

Lenses

Curved refractive  
surfaces

Lensmaker's equation


Again our condition for minimizing the time was given in Eq. 6:

$$0 = n_1 \frac{b}{\sqrt{y_1^2 + b^2}} - n_2 \frac{a - b}{\sqrt{y_2^2 + (a - b)^2}}$$

If we consult again our original diagram, we see that we can also express this as

$$\begin{aligned} 0 &= n_1 \sin \theta_1 - n_2 \sin \theta_2 \\ \text{giving } n_1 \sin \theta_1 &= n_2 \sin \theta_2 \end{aligned} \quad (7)$$

so we have proved Snell's law.



# TIR: total internal reflection

Wave superposition

Ray optics

Huygens

Snell's law

TIR

Refraction

Apparent depth



Magnification

Lenses

Curved refractive  
surfaces

Lensmaker's equation

- Again, with  $n_2 > n_1$ , we can get total internal reflection.
- As  $\theta_2$  increases,  $\theta_1$  goes to  $90^\circ$  and then you get total internal reflection!
- Known as the critical angle  $\theta_c$ :  
 $n_1 \sin 90^\circ = n_2 \sin \theta_c$ , or  
 $\sin \theta_c = n_1/n_2$ .



# TIR for swimming pools

Wave superposition

Ray optics

Huygens

Snell's law

TIR

Refraction

Apparent depth


Magnification

Lenses

Curved refractive surfaces

Lensmaker's equation

- Remember Snell's law:  
$$n_1 \sin \theta_1 = n_2 \sin \theta_2.$$
- Sit at the bottom of a swimming pool (water:  $n_2 = 1.33$ ) and look up at the sky, where  $n_1 = 1$ . That is,  $n_2 > n_1$ .
- As  $\theta_2$  increases,  $\theta_1$  goes to  $90^\circ$  and then you get total internal reflection!



Pool photo: <http://lifshitz.ucdavis.edu/~dmartin/phy7/7C/Refraction/Refraction.html>

## TIR for X rays

Wave superposition

Ray optics

Huygens

Snell's law

TIR

Refraction

Apparent depth



Magnification

Lenses

Curved refractive surfaces

Lensmaker's equation

- Reminder: we get Total Internal Reflection (TIR) within  $n_2$  at a critical angle  $\sin \theta_c = n_1/n_2$  when  $n_2 > n_1$ .
- But for X rays  $n = 1 - \delta - i\beta$ , or just  $n = 1 - \delta$  if we ignore absorption.
- Therefore we have  $n_2 > n_1$  if  $n_2 = 1$  for vacuum/air, and  $n_1 = 1 - \delta$  for a medium.
- That is, external reflection for X rays is really *total internal reflection*!
- The critical angle is  $\sin \theta_c = 1 - \delta$  with  $\theta_c$  from the surface normal, or  $\cos \theta'_c = 1 - \delta$  from the surface.



## X-ray critical angle $\theta'_c$

Wave superposition

Ray optics

Huygens

Snell's law

TIR

Refraction

Apparent depth

Magnification

Lenses


Curved refractive  
surfaces

Lensmaker's equation

- Again, for X rays we have  $\cos \theta'_c = 1 - \delta$  with  $\theta'_c$  measured from the surface.
- Since  $\delta$  is small, let's use an approximation:

$$1 - \frac{(\theta'_c)^2}{2} \simeq \cos \theta'_c = 1 - \delta \quad \Rightarrow \quad \theta'_c \simeq \sqrt{2\delta} = \lambda \sqrt{2\alpha f_1}$$

- Since  $\delta$  is small, X rays only reflect at small grazing angles!



# Refraction at a curved interface

Wave superposition

Ray optics

Huygens

Snell's law

TIR

Refraction

Apparent depth

Magnification

Lenses


Curved refractive surfaces

Lensmaker's equation

Consider refraction at an interface with a radius of curvature  $-R$ . We'll follow a ray from the point  $O$  as it travels at some angle  $\alpha$  and hits the interface at a point  $P$ . Now Snell's law tells us

$$\begin{aligned}n_1 \sin \theta_1 &= n_2 \sin \theta_2 \\n_1 \theta_1 &\simeq n_2 \theta_2 \\n_1(\alpha - \varphi) &\simeq n_2(\alpha' - \varphi) \quad (8)\end{aligned}$$

where we have used the small angle approximation. We get the relationship  $\theta_1 = \alpha - \varphi$  from considering the triangles  $PCQ$ ,  $CPO$ , and  $POQ$ . We get the relationship  $\theta_2 = \alpha' - \varphi$  from considering the triangles  $PCQ$ ,  $CPI$ , and  $PIQ$ .



## Refraction II

Wave superposition

Ray optics

Huygens

Snell's law

TIR

Refraction

Apparent depth

Magnification

Lenses

Curved refractive surfaces

Lensmaker's equation


Again, we had from Eq. 8 the relationship

$$n_1(\alpha - \varphi) = n_2(\alpha' - \varphi)$$

in the small angle approximation. Now let's express these angles in terms of distances on the diagram, where we ignore the distance  $QV$ :

$$\begin{aligned} n_1\left(\frac{h}{s} - \frac{h}{-R}\right) &= n_2\left(\frac{h}{-s'} - \frac{h}{-R}\right) \\ \frac{n_1}{s} + \frac{n_2}{s'} &= \frac{n_2 - n_1}{R} \end{aligned} \quad (9)$$

which is Fowles Eq. 10.4. Notice that we defined the distances  $s'$  and  $R$  to be negative in anticipation of what will follow. This is a general relationship for refraction at a curved interface.




## Apparent depth

Let's consider some consequences of the expression of Eq. 9 of

$$\frac{n_1}{s} + \frac{n_2}{s'} = \frac{n_2 - n_1}{R}.$$

First of all, consider standing on the side of  $n_2$  and looking at an object at point  $O$  within  $n_1$ , and letting  $R \rightarrow \infty$  for a flat interface. As you look at the object and as your brain assumes that light travels in straight lines, you will think that the object is really located at point  $I$ . That is, the apparent depth  $-s'$  of the object will be at

$$\begin{aligned}\frac{n_1}{s} &= -\frac{n_2}{s'} \\ s' &= -\frac{n_2}{n_1}s\end{aligned}\quad (10)$$



so if you're trying to spear a fish you will miss!




## Magnification

Now let's consider multiple rays from one point to another, hitting a convex refractive surface with positive  $R$ . One ray hits the spherical surface at  $\theta_1 = 0$  so that  $\theta_2 = 0$ . Another ray goes through the centerline of the lens so that the angle  $\theta_1$  from the source to the refractive interface is equal (in the small angle approximation) to  $h/s$ . The two rays cross at a point  $s'$  on the other side of the surface (hence our choice of sign convention earlier for  $s'$ ). The second ray reaches that point at an angle  $\theta_2 = h'/s'$ . From Snell's law we have

$$\begin{aligned} n_1 \frac{h}{s} &= n_2 \frac{h'}{s'} \\ \text{giving } m \equiv \frac{h'}{h} &= -\frac{n_1 s'}{n_2 s} \quad (11) \end{aligned}$$

where the negative sign accounts for the fact that the object is inverted.



# Curved refractive surfaces

Wave superposition

Ray optics

Huygens

Snell's law

TIR

Refraction

Apparent depth

Magnification


Lenses

Curved refractive surfaces

Lensmaker's equation


Consider two refractive surfaces. For the first surface we have

$$\frac{n_1}{s_1} + \frac{n_2}{s'_1} = \frac{n_2 - n_1}{R_1} \quad (12)$$




Consider a second surface immediately following. For it, we have the opposite ordering for  $n_1$  and  $n_2$ , and we have the opposite radius of curvature, giving

$$\frac{n_2}{s_2} + \frac{n_1}{s'_2} = \frac{n_1 - n_2}{-R_2} \quad (13)$$



Here's the key: if the lens is truly thin, then

$$s'_1 = -s_2. \quad (14)$$



## Two surfaces II

We can rewrite our expression of Eq. 12 for the first refractive interface as

$$\frac{1}{s'_1} = \frac{n_2 - n_1}{n_2} \frac{1}{R_1} - \frac{n_1}{n_2} \frac{1}{s_1} \quad (15)$$

and our expression of Eq. 13 for the second refractive interface as

$$\frac{1}{s_2} = \frac{n_1 - n_2}{n_2} \frac{1}{R_2} - \frac{n_1}{n_2} \frac{1}{s'_2} \quad (16)$$

Now let's use the result of Eq. 14 of  $s'_1 = -s_2$  to combine these two:

$$\begin{aligned} \frac{n_1 - n_2}{n_2} \frac{1}{R_1} + \frac{n_1}{n_2} \frac{1}{s_1} &= \frac{n_1 - n_2}{n_2} \frac{1}{R_2} - \frac{n_1}{n_2} \frac{1}{s'_2} \\ \frac{n_1}{n_2} \left( \frac{1}{s_1} + \frac{1}{s'_2} \right) &= \frac{n_1 - n_2}{n_2} \left( -\frac{1}{R_1} + \frac{1}{R_2} \right) \\ \frac{1}{s_1} + \frac{1}{s'_2} &= \frac{n_2 - n_1}{n_1} \left( \frac{1}{R_1} - \frac{1}{R_2} \right) \equiv \frac{1}{f} \quad (\text{Fowles 10.7}) \end{aligned}$$

## Two surfaces III

Wave superposition

Ray optics

Huygens

Snell's law

TIR

Refraction

Apparent depth

Magnification

Lenses

Curved refractive  
surfaces

Lensmaker's equation

For the overall optical system, we'll write  $s$  for  $s_1$  and  $s'$  for  $s'_2$ .  
Therefore we have derived the lensmaker's equation in Eq. 17:

$$\frac{1}{s} + \frac{1}{s'} = \frac{1}{f} \quad \text{with} \quad \frac{1}{f} \equiv \frac{n_2 - n_1}{n_1} \left( \frac{1}{R_1} - \frac{1}{R_2} \right)$$

This allows us to calculate the focal length of a lens with any set of spherical surfaces. This equation assumes  $+R$  for centers of curvature located “downstream” of the lens, so for a double-convex lens we have  $R_1 = +|R_1|$  and  $R_2 = -|R_2|$ .