

Machine learning under physical constraints

Unsupervised learning in DA

Sixin Zhang
(sixin.zhang@toulouse-inp.fr)

Outline

Unsupervised learning of Linear Dynamical systems

MLE approach and EM algorithm


Outline

Unsupervised learning of Linear Dynamical systems

MLE approach and EM algorithm

A special ODS for financial time series


- Build a statistical model of financial time series to analyze stock volume changes under news impact



- Model dynamics using **Linear dynamical systems** (Kalman Filter).


A special ODS for financial time series

- ▶ Model daily volume dynamics x_t over t .
- ▶ Assume $x_0 \sim \mathcal{N}(\mu_0, \Sigma_0)$, $A \in \mathbb{R}$, $B \in \mathbb{R}$, $C > 0$
- ▶ Dynamics: $x_{t+1}|x_t \sim \mathcal{N}(Ax_t + B, C)$



A special ODS for financial time series

- Model within-day volume $y_t \in \mathbb{R}^d$ for day t ,



- Assume $E \in \mathbb{R}^d$, F p.d matrix, $\mathbb{1}$ vector of ones
- Observation: $y_t | x_t \sim \mathcal{N}(\mathbb{1}x_t + E, F)$

Problem formulation

- ▶ Can we estimate the densities $p(x_t|x_{t-1})$ and $p(y_t|x_t)$ for $t = 1, \dots, T$ from $y = (y_1, \dots, y_T)$?
- ▶ Problem: estimate $\theta = (\mu_0, \Sigma_0, A, B, C, E, F)$ from y .
- ▶ **Challenge:** assume x_t is not observed (because certain y_t may be impacted by news, thus not observed)
- ▶ Idea: Maximum likelihood estimation with EM algorithm
- ▶ Reference: Pattern Recognition and Machine Learning by Christopher M. Bishop

Outline

Unsupervised learning of Linear Dynamical systems

MLE approach and EM algorithm

MLE approach and EM algorithm

- Identifiability condition (to restrict E):

$$E \perp \mathbb{1}$$

- Estimate θ from $y = (y_1, \dots, y_T)$,

$$\max_{\theta} \log p(y|\theta)$$

- Estimating of the latent variables $x = (x_0, \dots, x_T)$ makes the maximisation of θ easier,

$$\text{E step : } \max_{\phi} \int \log \frac{p(x|y, \theta)}{q(x|y, \phi)} q(x|y, \phi) dx$$

$$\text{M step : } \max_{\theta} \int \log p(x, y|\theta) q(x|y, \phi) dx$$

MLE approach and EM algorithm

- ▶ KF smoother makes the **E step** possible: compute

$$p(x_t | y_1, \dots, y_T, \theta)$$

$$p(x_t, x_{t+1} | y_1, \dots, y_T, \theta)$$

- ▶ **Factorization** of $\log p$ makes the **M step** possible: in KF,

$$\log p(x, y | \theta) = \log p(x_0 | \theta) + \sum_{t=0}^{T-1} \log p(x_{t+1} | x_t, \theta) + \sum_{t=1}^T \log p(y_t | x_t, \theta).$$

Road-map: 3 sub-problems in M step

- ▶ Decompose the M-step using factorization,

$$\theta^{k+1} \in \arg \max_{\theta} \int \log p(x, y | \theta) q(x | y, \phi^{(k)}) dx$$

- ▶ **Problem 1:** estimate (μ_0, Σ_0)

$$\max_{\theta} \int \log p(x_0 | \theta) q(x_0 | y, \phi^{(k)}) dx_0$$

This requires to solve the **E step** for $q(x_0 | y, \phi^{(k)})$ at θ^k .

- ▶ Problem 2: estimate (A, B, C)
- ▶ Problem 3: estimate (E, F)

Problem 1: analytical formula of M step

$$\ell_1 = \int \log p(x_0|\theta) q(x_0|y, \phi^{(k)}) dx_0$$

- ▶ Under $q(x_0|y, \phi^{(k)})$, x_0 follows $\mathcal{N}(\mu_{0|T}, \Sigma_{0|T})$
- ▶ Develop the log-likelihood,

$$-\ell_1 \propto \mathbb{E}_{x_0}((x_0 - \mu_0)^\top \Sigma_0^{-1} (x_0 - \mu_0)) + \log |\Sigma_0|$$

- ▶ To maximize ℓ_1 , compute the critical points

$$\frac{\partial \ell_1}{\partial \mu_0} = 0, \frac{\partial \ell_1}{\partial \Sigma_0} = 0 \quad \Rightarrow \quad \mu_0^{(k)} = \mu_{0|T}, \Sigma_0^{(k)} = \Sigma_{0|T}$$

E step: How to compute $\mathcal{N}(\mu_{0|T}, \Sigma_{0|T})$?

- ▶ KF propagation and analysis: compute $p(x_T|y)$ recursively.
- ▶ **KF smoother**: a key to compute

$$x_{t+1}|y_1, \dots, y_T \Rightarrow x_t|y_1, \dots, y_T$$

- ▶ Analysis step: $x_t|y_1, \dots, y_t \sim \mathcal{N}(\mu_{t|t}, \Sigma_{t|t})$
- ▶ Propagation step:

$$x_{t+1}|y_1, \dots, y_t \sim \mathcal{N}(\mu_{t+1|t}, \Sigma_{t+1|t})$$

E step: KF smoother

- Smoother: given

$$x_{t+1}|y_1, \dots, y_T \sim \mathcal{N}(\mu_{t+1|T}, \Sigma_{t+1|T})$$

compute

$$x_t|y_1, \dots, y_T \sim \mathcal{N}(\mu_{t|T}, \Sigma_{t|T})$$

- We verify that

$$\mu_{t|T} = \mu_{t|t} + J_t(\mu_{t+1|T} - \mu_{t+1|t})$$

$$\Sigma_{t|T} = \Sigma_{t|t} + J_t(\Sigma_{t+1|T} - \Sigma_{t+1|t})J_t^\top$$

where $J_t = \Sigma_{t|t}^\top A^\top (C + A\Sigma_{t|t}A^\top)^{-1}$

E step: KF smoother (technical detail)

- Verify that $p(x_t|x_{t+1}, y_1, \dots, y_T, \theta)$ follows

$$\mathcal{N}(\mu_{t|t+1|T}, \Sigma_{t|t+1|T})$$

where

$$\mu_{t|t+1|T} = \mu_{t|t} + J_t[x_{t+1} - (A\mu_{t|t} + B)]$$

$$\Sigma_{t|t+1|T} = \Sigma_{t|t} - J_t A \Sigma_{t|t}$$

- This implies the joint distribution (x_{t+1}, x_t) given y is

$$\mathcal{N}\left(\begin{pmatrix} \mu_{t+1|T} \\ \mu_{t|T} \end{pmatrix}, \begin{pmatrix} \Sigma_{t+1|T}, & (J_t \Sigma_{t+1|T})^\top \\ J_t \Sigma_{t+1|T}, & \Sigma_{t|T} \end{pmatrix}\right)$$

M step: analytical formula of Problem 2

- ▶ Consider $\ell_2 = \sum_t \ell_{2,t}$, where

$$\begin{aligned}\ell_{2,t} &= \int \log p(x_{t+1}|x_t, \theta) q(x_{t+1}, x_t|y, \phi^{(k)}) dx_{t+1} dx_t \\ &\propto -\mathbb{E}_{(x_{t+1}, x_t)}[(x_{t+1} - Ax_t - B)^\top C^{-1}(x_{t+1} - Ax_t - B)] - \log |C|\end{aligned}$$

- ▶ Compute critical points of ℓ_2 using matrix calculus,

$$\frac{\partial \ell_2}{\partial A} = 0, \quad \frac{\partial \ell_2}{\partial B} = 0, \quad \frac{\partial \ell_2}{\partial C} = 0$$

- ▶ Exercise: find out the optimal solution.

M step: analytical formula of Problem 3

- ▶ Consider $\ell_3 = \sum_t \ell_{3,t}$, where

$$\begin{aligned}\ell_{3,t} &= \int \log p(y_t|x_t, \theta) q(x_t|y, \phi^{(k)}) dx_t \\ &\propto -\mathbb{E}_{x_t}[(y_t - \mathbb{1}x_t - E)^\top F^{-1}(y_t - \mathbb{1}x_t - E)] - \log |F|\end{aligned}$$

- ▶ Due to the constraint on E , we compute critical points of ℓ_3 using the Lagrangian method.
- ▶ Exercise: find out the optimal solution (check https://cs.nyu.edu/~zsx/rapport_mva_sixin.pdf).