

Lecture Notes Thermal Physics

Mick Veldhuis

September 4, 2019

Contents

| | | |
|---|---|---|
| 1 | Introduction: a short overview of important notions | 1 |
|---|---|---|

1 Introduction: a short overview of important notions

The mole

A **mole** of a certain atom is equivalent to an **Avogadro number** N_A of atoms. Where $N_A = 6.022 \cdot 10^{23}$.

The ideal gas law

Suppose we have N particles in a gas, then we can relate the pressure, volume, temperature, and amount of particles as follows:

$$pV = Nk_B T \quad (1)$$

where k_B is Boltzmann's constant. In order to apply this formula one assumes that (i) there are no intermolecular forces and (ii) that the particles are point-like and have zero size.

Combinatorics

Suppose we have n atoms, and r of those atoms are for instance in an excited state. If we want to calculate the number of possible configurations Ω we apply the following formula,

$$\Omega = \frac{n!}{(n-r)!r!} \equiv {}^nC_r \quad (2)$$

Since factorials grow incredibly quick, we shall often use $\ln \Omega$ instead of Ω .

$$\ln \Omega = \ln(n!) - \ln((n-r)!) - \ln(r!) \quad (3)$$

We also have **Sterling's formula**:

$$\ln(n!) \approx n \ln n - n \quad (4)$$

2 Heat