

# 1. Prove the results about the smoothness of the members of the Matérn family.

We use the theorem that if

$$\frac{d^{2\nu}}{d\tau^{2\nu}}\rho(\tau)$$

exists and is finite at  $\tau = 0$ , then the random field having  $\rho(\tau)$  as its correlation function is  $\nu$  times differentiable at 0.

Without loss of generality, let  $\phi = 1$ . The Matérn correlation function is given by

$$\rho(\tau) = \frac{\tau^\nu}{2^{\nu-1}\Gamma(\nu)}K_\nu(\tau), \quad \tau \geq 0, \nu > 0.$$

For small  $\tau$  and  $\nu > 0$ ,  $K_\nu(\tau) \approx \Gamma(\nu)2^{\nu-1}\tau^{-\nu}$ . Also,  $\frac{d}{d\tau}\tau^\nu K_\nu(\tau) = -\tau^\nu K_{\nu-1}(\tau)$  and  $K_\nu(\tau) = K_{-\nu}(\tau)$ . We will be taking  $\tau \rightarrow 0$ , so we use the approximation for  $K_\nu(\tau)$ . This leads to the derivative,

$$\begin{aligned} \frac{d}{d\tau}\rho(\tau) &= -\frac{\tau^\nu}{2^{\nu-1}\Gamma(\nu)}K_{\nu-1}(\tau) \\ &= \begin{cases} -\frac{\tau^\nu}{2^{\nu-1}\Gamma(\nu)}K_{\nu-1}(\tau), & \nu - 1 \geq 0 \\ -\frac{\tau^\nu}{2^{\nu-1}\Gamma(\nu)}K_{1-\nu}(\tau), & \nu - 1 < 0 \end{cases} \\ &\approx \begin{cases} -\frac{\tau^\nu}{2^{\nu-1}\Gamma(\nu)}\Gamma(\nu-1)2^{\nu-2}\tau^{-\nu+1}, & \nu - 1 \geq 0 \\ -\frac{\tau^\nu}{2^{\nu-1}\Gamma(\nu)}\Gamma(1-\nu)2^{-\nu}\tau^{\nu-1}, & \nu - 1 < 0 \end{cases} \\ &\approx \begin{cases} -\tau G_1(\nu), & \nu - 1 \geq 0 \\ -\tau^{2\nu-1}G_2(\nu), & \nu - 1 < 0 \end{cases}. \end{aligned}$$

Therefore,

$$\rho'(0) \begin{cases} = 0, & \nu \geq 1 \\ \in (-\infty, 0), & 1/2 \leq \nu < 1 \\ = -\infty, & 0 < \nu < 1/2 \end{cases}.$$

The second derivative is given by

$$\rho''(\tau) = \begin{cases} \frac{-\tau^{\nu-1}K_{\nu-1}(\tau) + \tau^\nu K_{\nu-2}(\tau)}{2^{\nu-1}\Gamma(\nu)}, & \nu \geq 2 \\ \frac{-\tau^{\nu-1}K_{\nu-1}(\tau) + \tau^\nu K_{2-\nu}(\tau)}{2^{\nu-1}\Gamma(\nu)}, & 1 \leq \nu < 2 \\ \frac{-\tau^{\nu-1}K_{1-\nu}(\tau) + \tau^\nu K_{2-\nu}(\tau)}{2^{\nu-1}\Gamma(\nu)}, & 0 < \nu < 1 \end{cases},$$

and is evaluated at  $\tau = 0$  to

$$\rho''(0) \begin{cases} = 0, & \nu \geq 2 \\ \in (-\infty, 0), & 1 \leq \nu < 2 \\ = -\infty, & 0 < \nu < 1 \end{cases}.$$

We have that the second derivative is finite when  $\nu \geq 1$ , leading to a random field that is one time mean square differentiable. To show this generalizes to  $\nu \geq d$ , we need to keep taking derivatives of  $\rho(\tau)$ . I suspect that on each even derivative, the orders of certain Bessel functions are negated when those orders are less than  $\nu$ . When this happens, the approximation will lead to a term having  $\tau$  raised to a negative exponent causing the derivative to evaluate to  $-\infty$ .

## 2. Use the spectral representation to show that the product of two valid correlation functions is a valid correlation function.

A valid correlation function is the characteristic function of some random variable,

$$\rho(\boldsymbol{\tau}) = E \left[ e^{i\boldsymbol{\tau}^\top \mathbf{X}} \right].$$

Suppose we have two valid correlation functions  $\rho_1(\boldsymbol{\tau})$  and  $\rho_2(\boldsymbol{\tau})$  associated with independent random variables  $\mathbf{X}_1$  and  $\mathbf{X}_2$ , respectively. Then the product is written

$$\begin{aligned} \rho(\boldsymbol{\tau}) &= \rho_1(\boldsymbol{\tau})\rho_2(\boldsymbol{\tau}) = E \left[ e^{i\boldsymbol{\tau}^\top \mathbf{X}_1} \right] E \left[ e^{i\boldsymbol{\tau}^\top \mathbf{X}_2} \right] \\ &= E \left[ e^{i\boldsymbol{\tau}^\top \mathbf{X}_1} e^{i\boldsymbol{\tau}^\top \mathbf{X}_2} \right] \\ &= E \left[ e^{i\boldsymbol{\tau}^\top (\mathbf{X}_1 + \mathbf{X}_2)} \right], \end{aligned}$$

so  $\rho$  is the characteristic function of  $\mathbf{X}_1 + \mathbf{X}_2$  and thus the product of two valid correlation functions is a valid correlation function. Note, our assumption of independence for  $\mathbf{X}_1$  and  $\mathbf{X}_2$  presents no issues.  $\mathbf{X}_1$  and  $\mathbf{X}_2$  may be dependent, but then we could simply define new independent random variables  $\mathbf{Y}_1$  and  $\mathbf{Y}_2$  with the same marginal distributions as  $\mathbf{X}_1$  and  $\mathbf{X}_2$ , resulting in the same correlation functions in either case.

## 3. The spectral density of a correlation in the Matérn family has tails whose thickness depends on the smoothness parameter. Conjecture: the smoothness of the corresponding random field depends on the number of moments of the spectral density. What can you say about this conjecture?

For correlation function

$$\rho(\tau) \propto (a\tau)^\nu K_\nu(a\tau), \quad \tau \geq 0, \nu > 0, a = 1/\phi > 0,$$

we have the corresponding spectral density

$$f(x) \propto \frac{1}{(1 + (x/a)^2)^{\nu+n/2}},$$

where  $n$  is the dimension  $\tau$  (and  $x$ ). This density has a form comparable to the  $t$ -distribution. Using integration by parts, we calculate the  $k$ th moment as

$$\begin{aligned} E(X^k) &\propto \int x^k (1 + (x/a)^2)^{-(\nu+n/2)} dx \\ &= -\frac{a^2}{2\nu + n - 2} \frac{x^{k-1}}{(1 + (x/a)^2)^{(2\nu+n-2)/2}} \Big|_{-\infty}^{\infty} + \int \frac{(k-1)a^2}{2\nu + n - 2} \frac{x^{k-2}}{(1 + (x/a)^2)^{(2\nu+n-2)/2}} dx. \end{aligned}$$

The first term (and hence the second term also) will be finite when  $2\nu + n - 2 \geq k - 1$ , or  $\nu \geq (k - n + 1)/2$ . In one dimension,  $n = 1$ , we see that when  $\nu \geq k/2$  the  $k$ th moment exists. This may be related to the theorem used in the first problem, that we need to have  $2d$ -differentiable correlation function to have a  $d$ -differentiable random field. Here, we need the  $2d$ th moment to exist, i.e.  $\nu \geq d$ , to have smoothness.