

The Width of the Conformal Fan: Dependence and the Variance of Realized Coverage

Peter Cotton

`peter.cotton@microprediction.com`

June 2026

Abstract

Fix a split-conformal calibration set; the coverage realized against an independent draw from the score marginal is $c = U_{(k)}$, distributed $\text{Beta}(k, n - k + 1)$ for independent calibration scores, with variance about $\alpha(1 - \alpha)/n$ — the fan. We show the fan’s width is set by the dependence among the calibration scores, through the covariance kernel of the sub-threshold count. Three results follow: a leading-order fan coefficient in a Bahadur regime; an exact aggregate bound, summed over all levels, showing negative association never widens the fan; and, for equicorrelated normals, the exact single-level law $\text{Var}(Z_{(k)}) = v_k + \rho(1 - v_k)$. The general single-level contraction turns out to fail at $n = 2$: we exhibit an explicit negatively associated pair, uniform marginals, whose extreme order statistic is strictly over-dispersed. For $n \geq 3$ it is open and unbroken across an extensive search, and holds on the Gaussian-copula, FGM, and without-replacement subclasses.

1 Setup

Split conformal returns a set with marginal coverage $1 - \alpha$. Fix the calibration set; the coverage you then realize on fresh test points is a random variable, and it is not $1 - \alpha$ on the nose. Write the scores on the probability-integral scale, $U_i = F(S_i)$. We assume throughout that the score marginal F is continuous, so each $U_i \sim \text{Unif}(0, 1)$; if F has atoms, read U_i as the randomised PIT $F(S_i^-) + V_i(F(S_i) - F(S_i^-))$ with $V_i \sim \text{Unif}(0, 1)$ independent, which restores exact uniformity. With $k = \lceil (n+1)(1-\alpha) \rceil$ the conformal threshold is the order statistic $U_{(k)}$, and the calibration-conditional coverage against an independent test draw from the marginal is

$$c = U_{(k)}. \tag{1}$$

For *independent* calibration scores this is the classical result (Vovk, 2012; Marques F., 2025): $c \sim \text{Beta}(k, n - k + 1)$, with

$$\mathbb{E}[c] = \frac{k}{n+1}, \quad \text{Var}(c) = \frac{k(n-k+1)}{(n+1)^2(n+2)} \approx \frac{\alpha(1-\alpha)}{n}. \tag{2}$$

This across-sample spread of realized coverage is its *dispersion*, the formal object of this note; the family of such laws swept out as the calibration dependence varies is the *fan*, for short. The per-dataset coverage scatters around $1 - \alpha$ with variance of order $1/n$ (standard deviation of order $n^{-1/2}$). The question of this note is what the dependence among the calibration scores does to it.

Two cautions fix the object of study. First, $c = U_{(k)}$ is coverage against a test draw *independent* of the calibration set; it is not the conformal conditional coverage $\mathbb{P}(U_{n+1} \leq U_{(k)} \mid U_1, \dots, U_n)$ for

a test point exchangeable with, and dependent on, the calibration scores, which need not equal $U_{(k)}$. The marginal conformal guarantee comes from the rank symmetry of U_{n+1} among the $n+1$ scores and is not what controls c . Second, the mean $\mathbb{E}[c] = k/(n+1)$ is an *independence* statement, not a consequence of uniform marginals or exchangeability: if $U_1 = \dots = U_n = V \sim \text{Unif}(0, 1)$ then $U_{(k)} = V$ and $\mathbb{E}[c] = \frac{1}{2}$ for every k . So under dependence both the mean and the variance of c respond; this note studies the variance, and flags the mean where it matters. The companion notes treat the orthogonal, within-dataset axis (Cotton, a,b); here every statement is about the across-dataset fluctuation of c . The results separate by type: Theorems 2 and 4 and Propositions 1–3 are finite-sample exact, Theorem 1 is asymptotic and conditional on a Bahadur representation, and the general per-level negative-association contraction (Conjecture 1) is open.

A third point fixes what kind of dependence is in scope, since it has to square with an independent test. What the analysis requires is only that the test draw be independent of the calibration scores; the calibration dependence is then necessarily *design-induced* — a property of how the calibration set is assembled, not of a data stream the test belongs to. This is the standard split-conformal test assumption (a fresh, independent future point); the only non-standard ingredient is letting the *calibration* set carry an engineered dependence. That ingredient is not exotic — it is what you get whenever the calibration set is *constructed or curated* rather than taken as an i.i.d. draw, and the fan is then a statement about a design choice.

Three concrete settings. *Space-filling calibration*. When the calibration inputs can be chosen — active or pool-based calibration, or any pipeline that selects which points to label — one spreads them evenly, on a stratified, Latin-hypercube, or low-discrepancy (quasi-Monte-Carlo) grid, to avoid clumping. Even coverage of the input space makes the resulting scores negatively associated with uniform marginals, and the fan says this engineering can only tighten the run-to-run variability of realized coverage, never loosen it. *Balanced subsampling from a pool*. Labeled data is often subsampled to a fixed calibration budget; drawing without replacement, or balancing on a stratifying covariate, or antithetically pairing, all induce negative association among the retained scores while the deployed test point remains exogenous. *Deduplication and diversity filtering*. Removing near-duplicates or enforcing diversity in a calibration set — routine when data are scraped or redundant — is again a negative-dependence-inducing design, and again a fan-narrowing one. In each case the practical payoff is the same: a calibration design that is more evenly spread buys a more reliable realized coverage from one deployment to the next, at no cost to the marginal guarantee, which the rank symmetry of the independent test point still delivers.

The contrast is the case where the test shares its source with the calibration — the next step of a time series, or the unsampled remainder of a without-replacement draw used as the test — which couples the test to the calibration, so that $c \neq U_{(k)}$ and a different analysis applies; that case is out of scope here. The maximally negatively associated floor is read the same way: a deterministic grid laid down by design, not a draw whose leftover is the test.

The whole problem passes through the count process $N(u) = \sum_{i=1}^n \mathbf{1}\{U_i \leq u\}$, because $U_{(k)} = \int_0^1 \mathbf{1}\{N(u) < k\} du$, so

$$\text{Var}(c) = \int_0^1 \int_0^1 \text{Cov}(\mathbf{1}\{N(u) < k\}, \mathbf{1}\{N(v) < k\}) du dv, \quad (3)$$

a functional of the law of the count process alone — equivalently, by Sklar’s theorem, of the score *copula*, since the uniform marginal has factored out (we return to this in Remark 7). Two facts about N drive everything.

Lemma 1 (Count variance). *If the scores are negatively associated (Joag-Dev and Proschan, 1983), then for every u , $\text{Var}(N(u)) \leq n u(1-u)$, with equality iff the sub-threshold indicators are pairwise*

uncorrelated.

Proof. The indicators $\mathbf{1}\{U_i \leq u\}$ are functions of single distinct coordinates, all nondecreasing in the same direction, so negative association is preserved and they are pairwise non-positively correlated. So $\text{Var}(N(u)) = \sum_i \text{Var} + \sum_{i \neq j} \text{Cov} \leq \sum_i u(1-u) = n u(1-u)$. $\square$

2 The fan coefficient

Let $p = 1 - \alpha$ and $b_i = \mathbf{1}\{U_i \leq p\}$, so $N(p) = \sum_i b_i$. (We reserve ξ for the i.i.d. normals of Proposition 3.) The representation is on the uniform scale, where the quantile density is $1/f(p) = 1$, so the usual $1/f$ factor is absent.

Theorem 1 (Fan coefficient). *Suppose F is continuous (or a randomised PIT is used), $k/n \rightarrow p \in (0, 1)$, and the scores satisfy a Bahadur representation $U_{(k)} - p = p - N(p)/n + r_n$ with the remainder negligible in L^2 relative to $n^{-1} \text{Var}(N(p))^{1/2}$. Then*

$$\text{Var}(c) = \frac{1}{n^2} \text{Var}(N(p)) + o\left(\frac{\text{Var}(N(p))}{n^2}\right). \quad (4)$$

If moreover $\text{Var}(N(p)) \asymp n$, writing it as $np(1-p) + n(n-1)\bar{c}$ with $\bar{c} = \frac{1}{n(n-1)} \sum_{i \neq j} \text{Cov}(b_i, b_j)$,

$$\text{Var}(c) = \frac{p(1-p)}{n} R(1 + o(1)), \quad R = 1 + \frac{n-1}{p(1-p)} \bar{c}. \quad (5)$$

So the leading fan coefficient is the count variance at the operating quantile. Under independence $\bar{c} = 0$, $R = 1$, recovering (2); negative association makes $\bar{c} < 0$ and $R < 1$, narrowing the fan. Two caveats the statement respects. The multiplicative form needs $\text{Var}(N(p)) \asymp n$: under strong negative dependence the count variance can be $o(n)$ (zero at the floor), and then only the additive form is safe. And the Bahadur regime itself fails under a fixed positive latent (de Finetti) dependence, where $N(p)/n$ has a non-degenerate random limit $Q([0, p])$ and $U_{(k)}$ converges to a random conditional quantile rather than linearizing at p ; the next section handles that regime exactly and separately.

3 Positive dependence: an exact decomposition

When the scores are extendable, drawn from an infinitely exchangeable sequence, de Finetti (de Finetti, 1937; Kerns and Székely, 2006) makes them conditionally independent given a latent directing measure Q . The law of total variance then gives an exact decomposition.

Theorem 2 (Positive-dependence decomposition). *For an extendable exchangeable calibration sequence with directing measure Q ,*

$$\text{Var}(c) = \underbrace{\mathbb{E}_Q[\text{Var}(U_{(k)} \mid Q)]}_{\text{average within-}Q \text{ order-stat variance}} + \underbrace{\text{Var}_Q(m_k(Q))}_{\text{between-}Q \text{ term}}, \quad m_k(Q) = \mathbb{E}[U_{(k)} \mid Q]. \quad (6)$$

The between- Q term is non-negative and vanishes under independence.

The decomposition is exact and needs no asymptotics, but note what conditioning gives: $U_i \mid Q$ are i.i.d. Q , so $U_{(k)} \mid Q$ is the k -th order statistic of Q on the marginal scale, *not* a uniform Beta variable, and its mean $m_k(Q)$ moves with Q . The within- Q term is thus not the independent fan (2), and the identity by itself does *not* prove $\text{Var}(c)$ exceeds the independent value: that comparison

would need a separate theorem relating the average within- Q order-stat variance to the uniform fan. What the identity does isolate is the between- Q term, one signature of positive dependence (it also shifts the within- Q law from uniform to Q): a per-dataset latent shifts the whole calibration sample, and the realized coverage rides with it. Empirically the net effect is a wider fan, e.g. a one-factor Gaussian copula at correlation 0.2 gives $\text{Var}(c) \approx 3.3$ times the independent value (and the comonotone limit $U_1 = \dots = U_n$ gives $\frac{1}{12}$, far above it), but that widening is observed, not proved by Theorem 2 alone.

4 Negative dependence narrows the fan

The negative regime is the one a genuine prior cannot reach. By Kerns and Székely (2006) a finite exchangeable sequence with $\rho < 0$ has a *signed* directing measure, so the clean conditioning of Theorem 2 is unavailable: there is no Q to condition on, and the between-dataset term would be a negative quasi-variance. That is precisely the regime where the fan contracts.

Theorem 3 (Endpoints). *Independent scores give $c \sim \text{Beta}(k, n - k + 1)$, the benchmark fan. At the exchangeable pairwise-correlation floor, take the discrete rank idealization in which the calibration scores are a uniformly random permutation of the grid $\{1/(n + 1), \dots, n/(n + 1)\}$ (sampling without replacement, pairwise correlation $-1/(n - 1)$, the smallest attainable for an exchangeable vector): the order statistic is deterministic, $U_{(k)} = k/(n + 1)$ a.s., so $\text{Var}(c) = 0$.*

Independence is the benchmark, not the maximum: positive dependence can widen the fan well past it (the comonotone limit reaches $\frac{1}{12}$). The claim of this note is the other direction, that *negative* association does not widen it, with independence the (conjectural) upper endpoint of the negative-dependence range. Two model cautions on the floor. The grid permutation has *discrete*-uniform marginals, not the continuous $\text{Unif}(0, 1)$ of $U_i = F(S_i)$, so it sits in the model only if one allows discrete or rank scores; the continuous-marginal analogue is a Latin hypercube (one point per stratum, randomly permuted), whose fan is small, of order $1/n^2$, but not exactly zero. And the endpoints agree with Theorem 1 at $R = 1$ and $R = 0$. The two are joined by an exact path.

Proposition 1 (An exact variance interpolation). *Let the calibration law be, with probability $1 - t$, n independent uniforms, and with probability t , a contest (a uniformly random permutation of the grid $\{1/(n + 1), \dots, n/(n + 1)\}$). Both components yield realized coverage with the same mean $k/(n + 1)$, the contest with zero variance, so by the law of total variance the between-component term vanishes and*

$$\text{Var}(c) = (1 - t) \frac{k(n - k + 1)}{(n + 1)^2(n + 2)}. \quad (7)$$

Proof. Condition on the component. $\mathbb{E}[c \mid \text{iid}] = \mathbb{E}[c \mid \text{contest}] = k/(n + 1)$, so $\text{Var}(\mathbb{E}[c \mid \cdot]) = 0$, and $\mathbb{E}[\text{Var}(c \mid \cdot)] = (1 - t) \frac{k(n - k + 1)}{(n + 1)^2(n + 2)}$. $\square$

So the fan collapses linearly to zero as $t \rightarrow 1$. Two caveats. The mixture's pairwise correlation is $\text{Corr}(U_i, U_j) = -t/(n + 1 - 2t)$, not $-t/(n - 1)$: the grid component contributes $\text{Cov} = -1/(12(n + 1))$ but mixing changes the marginal variance to $(n + 1 - 2t)/(12(n + 1))$; the correlation reaches the floor $-1/(n - 1)$ only at $t = 1$. And the mixture's one-dimensional marginal is neither continuous uniform nor discrete-grid uniform, so this is an exact *variance* interpolation, not a fixed-marginal copula path.

So we can prove the contraction exactly at both endpoints and along this whole interpolating family, and to leading order everywhere (Theorem 1). For an *arbitrary* negatively associated law the contraction also holds once it is summed over all levels.

Theorem 4 (Aggregate fan bound). *For negatively associated scores with uniform marginals,*

$$\sum_{k=1}^n \text{Var}(U_{(k)}) \leq \sum_{k=1}^n \frac{k(n-k+1)}{(n+1)^2(n+2)}, \quad (8)$$

the independent value, with equality holding under independence. Summed over all target levels, the fan is never widened by negative association.

Proof. The order statistics are a permutation of the sample, so $\sum_k U_{(k)}^2 = \sum_i U_i^2$ and, under uniform marginals, $\sum_k \mathbb{E}[U_{(k)}^2] = n \mathbb{E}[U_1^2] = n/3$ for every such law; hence $\sum_k \text{Var}(U_{(k)}) = \frac{n}{3} - \sum_k (\mathbb{E} U_{(k)})^2$. For the top j statistics, $\sum_{k>n-j} U_{(k)} = \int_0^1 \min(j, n-N(u)) du$. The function $\phi(m) = \max(-j, m-n)$ is convex and *nondecreasing*, which is exactly the class for which Shao (2000) compares a negatively associated sum to its independent counterpart, $\mathbb{E}\phi(N(u)) \leq \mathbb{E}\phi(\text{Bin}(n, u))$. (Because the means agree, $\mathbb{E}N(u) = nu = \mathbb{E}\text{Bin}$, one may add any affine term, so the comparison in fact extends to all convex ϕ , the full convex order; the monotone-convex form already suffices here.) Since $\min(j, n-m) = -\phi(m)$, this gives $\mathbb{E} \min(j, n-N(u)) \geq \mathbb{E} \min(j, n-\text{Bin}(n, u))$; integrating,

$$\sum_{k>n-j} \mathbb{E} U_{(k)} \geq \sum_{k>n-j} \frac{k}{n+1} \quad \text{for every } j = 1, \dots, n, \quad (9)$$

the right side being the independent profile's top- j sum. Both profiles are increasing and have the same total, $\sum_k \mathbb{E} U_{(k)} = \sum_i \mathbb{E} U_i = n/2 = \sum_k \frac{k}{n+1}$ (the case $j = n$ of (9), with equality). For two increasing vectors with equal sum, “every top- j partial sum of one dominates that of the other” is exactly the definition of majorization; so $(\mathbb{E} U_{(1)}, \dots, \mathbb{E} U_{(n)})$ majorizes $(\frac{1}{n+1}, \dots, \frac{n}{n+1})$. The map $x \mapsto \sum_k x_k^2$ is Schur-convex, hence larger on the majorizing vector, $\sum_k (\mathbb{E} U_{(k)})^2 \geq \sum_k (\frac{k}{n+1})^2$, and the displayed identity for $\sum_k \text{Var}(U_{(k)})$ finishes it. $\square$

The proof uses only identical uniform marginals and the convex-order domination $N(u) \leq_{\text{cx}} \text{Bin}(n, u)$, not exchangeability as such. Equality holds under independence; that equality *forces* independence is not established here and would need a separate tightness lemma.

5 The single level: a counterexample at $n = 2$, and a conjecture for $n \geq 3$

Everything above is proved for general negative association. The single conformal level, the operationally relevant one, is not, and the obstruction is exactly why the aggregate succeeds: the step $\mathbf{1}\{N(u) < k\}$ behind a single $\text{Var}(U_{(k)})$ is neither concave nor convex, so the convex order does not sign it, whereas the smooth aggregate functional $\min(j, n-N)$ is concave and it does. That gap is not merely technical. At $n = 2$ the single-level bound is *false*.

Proposition 2 (Counterexample at $n = 2$). *For each $a \in (0, \frac{\sqrt{11}-3}{2})$ there is a negatively associated pair with uniform marginals whose extreme order statistic is strictly over-dispersed: $\text{Var}(U_{(2)}) = \frac{1}{18} + \frac{a^2(1-6a-2a^2)}{18} > \frac{1}{18} = \text{Var}^{\text{iid}}(U_{(2)})$.*

Proof. Split $[0, 1]$ into a bottom bin $B = [0, 1-a]$ and a top bin $T = [1-a, 1]$. Draw the cell of (U_1, U_2) with $\mathbb{P}(B, T) = \mathbb{P}(T, B) = a$, $\mathbb{P}(B, B) = 1-2a$, $\mathbb{P}(T, T) = 0$, and, given the cell, make the coordinates independent uniforms on their bins: the two points are *never both large*. Each marginal is uniform (mass a spread uniformly on T , mass $1-a$ on B). For a pair,

negative association is equivalent to negative quadrant dependence $C(u, v) \leq uv$ (Joag-Dev and Proschan, 1983; Lehmann, 1966); here $uv - C(u, v) = a^2 (g_B(u) - g_T(u))(g_B(v) - g_T(v)) \geq 0$ with $g_B(x) = \min(x/(1-a), 1)$, $g_T(x) = \max(0, (x - (1-a))/a)$ and $g_B \geq g_T$ pointwise, so the law is genuinely NA. Its diagonal is $\delta(u) = \mathbb{P}(U_{(2)} \leq u) = (1-2a)u^2/(1-a)^2$ on $[0, 1-a]$ and $2u-1$ on $[1-a, 1]$; with $\mathbb{E}[U_{(2)}^m] = 1 - m \int_0^1 u^{m-1} \delta(u) du$ this gives the stated variance, positive on the displayed range of a . $\square$

The excess is small but exact: at $a = \frac{1}{10}$, $\text{Var}(U_{(2)}) = \frac{1673}{30000}$, a ratio 1.0038 to the independent fan; optimising the diagonal (a Bertino copula) reaches about 1.034. By the negation symmetry $U \mapsto 1 - U$, which preserves NA and uniform marginals, the mirror law (never both *small*) over-disperses the minimum by the same amount, so the bound fails at both extreme levels. This is exactly the diagonal the linear program of (13) below produces: the note's hope there — that no genuine NA law realises a centroid-violating diagonal — is false at $n = 2$. The mechanism is that uniform marginals and exchangeability pin only the *sum* of the order-statistic means (to $n/2$), never an individual $\mathbb{E}[U_{(k)}]$, so a mean-decreasing spread is available: the counterexample lowers $\mathbb{E}[U_{(2)}]$ while widening it, and the aggregate bound survives because the other level over-contracts to compensate ($\text{Var}(U_{(1)}) + \text{Var}(U_{(2)}) \approx 0.104 < \frac{2}{18}$).

Remark 1 (Why $n = 2$ is special). *At $n = 2$ the order-statistic variance is a functional of the copula diagonal $\delta(u) = C(u, u)$ alone, and negative association collapses to the single diagonal constraint $\delta(u) \leq u^2$ — for a pair, NA, NQD, and negative correlation of monotone functions all coincide. That one-dimensional constraint leaves room to push the deficit $u^2 - \delta$ toward $u = 1$, which is precisely the over-dispersion. For $n \geq 3$ the single-level variance depends on the off-diagonal negative-orthant probabilities across thresholds (the content (13) isolates), and NA is genuinely higher-order there — the same reason pairwise negative dependence is strictly weaker than NA for $n \geq 3$, and the reason the exchangeable correlation floor $-1/(n-1)$ tightens away from the countermonotone -1 available only at $n = 2$. The counterexample does not survive that tightening: padding it with an independent uniform (NA by closure, Joag-Dev and Proschan, 1983) already drops the ratio below one, 0.9995 at $n = 3$, and further below for $n = 4, 5, 6$; the independent coordinate dilutes the effect. An extensive search over exchangeable, moving-average, block, without-replacement, and non-exchangeable negatively-correlated Gaussian-copula laws at $3 \leq n \leq 10$ and every k finds no violation (`check_fan_conjecture.py`).*

Conjecture 1 (Per-level contraction, $n \geq 3$). *For $n \geq 3$, if the calibration scores are negatively associated then $\text{Var}(c)$ is at most the independent value (2) for each k individually.*

Two subclasses settle it outright. The Farlie–Gumbel–Morgenstern copula at negative parameter has $\text{Var}(U_{(k)})$ a concave quadratic in the parameter with positive slope at independence, hence below the independent value throughout the NA range; sampling without replacement reduces, via the same diagonal functional, to a finite centroid inequality that the hypergeometric single-crossing settles. And in the canonical continuous family the per-level statement is exact, and more, it is monotone in the correlation.

Proposition 3 (Exact per-level law for equicorrelated normals). *Let $(Z_1, \dots, Z_n)$ be equicorrelated standard normal with correlation $\rho \in [-\frac{1}{n-1}, 1]$, and let $v_k = \text{Var}(\xi_{(k)})$ be the variance of the k -th order statistic of n i.i.d. standard normals. Then, for every k ,*

$$\text{Var}(Z_{(k)}) = v_k + \rho(1 - v_k). \quad (10)$$

As $v_k < 1$ for $n \geq 2$, this is strictly increasing in ρ : negative correlation strictly shrinks the variance of every order statistic, positive correlation inflates it.

Proof. Write $Z_i = \bar{Z} + (Z_i - \bar{Z})$. For equicorrelated normals $\bar{Z} \sim N(0, s)$, $s = (1 + (n-1)\rho)/n$, is independent of the centred vector, and $\text{Cov}(Z_i - \bar{Z}, Z_j - \bar{Z}) = (1 - \rho)(\delta_{ij} - 1/n)$, exactly $1 - \rho$ times the covariance of $(\xi_i - \bar{\xi})$ for i.i.d. standard normals. Hence $Z_{(k)} \stackrel{d}{=} \sqrt{s}\eta + \sqrt{1 - \rho}(\xi - \bar{\xi})_{(k)}$ with $\eta \sim N(0, 1)$ independent of the second term, so $\text{Var}(Z_{(k)}) = s + (1 - \rho)\text{Var}((\xi - \bar{\xi})_{(k)})$. Both terms are affine in ρ ; matching v_k at $\rho = 0$ forces $\text{Var}((\xi - \bar{\xi})_{(k)}) = v_k - 1/n$, and substituting gives $v_k + \rho(1 - v_k)$. $\square$

Remark 2. *This is the per-level contraction, exact and monotone, on the Gaussian scale, unifying the two sides: the single line $v_k + \rho(1 - v_k)$ runs from inflation at $\rho > 0$ through the independent value at $\rho = 0$ to the floor at $\rho = -1/(n-1)$. Under the Gaussian copula $U_i = \Phi(Z_i)$, which restores uniform marginals, $\text{Var}(U_{(k)})$ is numerically monotone in ρ and below the independent value throughout $\rho < 0$ (`check_fan.py`). The transfer to the uniform scale is a dispersive-order statement, and it is half rigorous. Decompose along the common mean, $Z_{(k)} = \sqrt{1 - \rho}D + \bar{Z}$ with $D = (\xi - \bar{\xi})_{(k)}$ and $\bar{Z} \sim N(0, s)$ independent. D is log-concave, being the order statistic of a vector with log-concave (Gaussian) joint density (Prékopa, 1973); and convolution with a log-concave density increases the dispersive order (Lewis and Thompson, 1981), so adding the Gaussian shift $\bar{Z}$ disperses, monotonically in s . The residual is that this outweighs the simultaneous $\sqrt{1 - \rho}$ shrinkage of D as ρ grows, a one-parameter density-quantile monotonicity under the variance budget $\frac{n-1}{n}(1 - \rho) + s = 1$, which no off-the-shelf convolution theorem delivers but which we now settle.*

Tweedie's formula sharpens that residual to a single posterior-variance bound. Reading $Y := Z_{(k)} = aD + b\eta$ ($a = \sqrt{1 - \rho}$, $b = \sqrt{s}$) as Gaussian signal extraction of the log-concave signal D , the slope of the posterior mean equals the scaled posterior variance, $\frac{d}{dq}\mathbb{E}[D \mid Y = q] = a \text{Var}(D \mid Y = q)/b^2$ (Efron, 2011; Hansen and Tong, 2026). The dispersive monotonicity is equivalent to this slope being at most $(n-1)a/n$, so, using $\frac{n-1}{n}a^2 + b^2 = 1$, the entire transfer reduces to

$$\text{Var}(D \mid Y = q) \leq \frac{n-1}{n}b^2 \quad \text{for all } q, \quad (11)$$

a posterior-variance bound for log-concave Gaussian denoising, which the next lemma settles. The score-driven reading of the same identity (Hansen and Tong, 2026) is the econometric form of modelling the conditional spread, which is exactly what closes the information gap a conformal wrapper leaves (Cotton, a).

Lemma 2 (Strong log-concavity of D). $D = (\xi - \bar{\xi})_{(k)}$ is $\frac{n}{n-1}$ -strongly log-concave, $(\log g_D)'' \leq -n/(n-1)$.

Proof. The centred vector $w = \xi - \bar{\xi}$ has covariance $I - \frac{1}{n}J$, an orthogonal projection, so on the hyperplane $\{\sum_i w_i = 0\}$ it is a standard Gaussian: its $-\log$ density is $\frac{1}{2}\|w\|^2$, with Hessian the identity. The law of the ordered vector is this density restricted to the convex cone $\{w_1 \leq \dots \leq w_n\}$. The cone enters only through the indicator of a convex set, a log-concave factor; writing the restricted density as $\exp(-\frac{1}{2}w^\top(I - \frac{1}{n}J)^+w - \iota_C(w))$ with ι_C the convex indicator of the cone, the Hessian of the potential is $(I - \frac{1}{n}J)^+ + \nabla^2\iota_C \succeq (I - \frac{1}{n}J)^+$. So the strong-log-concavity constant is set by the Gaussian quadratic alone; the cone can only raise it, never lower it, and in particular does not interfere through its boundary. Strong log-concavity passes to marginals (Prékopa, 1973; Brascamp and Lieb, 1976; Saumard and Wellner, 2014), the marginal of a coordinate being strongly log-concave to the reciprocal of that coordinate's reference variance; for $w_{(k)}$ this is $\frac{n-1}{n}$ (the diagonal of $I - \frac{1}{n}J$), giving constant at least $\frac{n}{n-1}$ (the parameter bookkeeping is in the Appendix). $\square$

Proposition 4 (Gaussian-scale dispersive monotonicity). *For equicorrelated standard normals with $\rho \in (-\frac{1}{n-1}, 1)$, $Z_{(k)}$ is increasing in the dispersive order as ρ increases.*

Proof. For a smooth one-parameter family the quantile evolves by $\partial_\rho q_\rho(u) = \mathbb{E}[\dot{Y} \mid Y = q_\rho(u)]$ with $\dot{Y} = \partial_\rho Z_{(k)}$, and dispersive monotonicity is exactly $\partial_\rho q_\rho(u)$ increasing in u , i.e. increasing in q . Writing $Y = aD + b\eta$ and using $a\mathbb{E}[D \mid Y] + b\mathbb{E}[\eta \mid Y] = Y$ together with the budget $\frac{n-1}{n}a^2 + b^2 = 1$, this reduces to $m'(q) \leq (n-1)a/n$, where $m(q) = \mathbb{E}[D \mid Y = q]$ (the computation of $M'(q)$ is in the Appendix). Tweedie's identity for Gaussian signal extraction gives $m'(q) = a \text{Var}(D \mid Y = q)/b^2$. The posterior of D given Y has $-\log$ density with Hessian $(-\log g_D)'' + a^2/b^2 \geq \frac{n}{n-1} + \frac{a^2}{b^2} = \frac{n}{(n-1)b^2}$, the last equality by the budget and Lemma 2; the Brascamp–Lieb variance inequality (Brascamp and Lieb, 1976) then gives $\text{Var}(D \mid Y = q) \leq (n-1)b^2/n$, hence $m'(q) \leq (n-1)a/n$. Equivalently the quantile density $q'_\rho(u) = 1/f_\rho(q_\rho(u))$ increases in ρ at each level (the density at each quantile decreases), which is the dispersive monotonicity. At the open-interval ends the statement is read in the limit: $b \rightarrow 0$ at $\rho = -1/(n-1)$ (the degenerate floor, $Z_{(k)} \rightarrow \sqrt{n/(n-1)}D$) and $a \rightarrow 0$ at $\rho = 1$ ($Z_{(k)} \rightarrow \eta$). $\square$

This settles the Gaussian-scale per-level statement in the strong (dispersive, not merely variance) sense. The one step left for the uniform-marginal conjecture under the Gaussian copula is the passage through Φ : being S-shaped, Φ is neither convex nor concave, so it need not transmit the dispersive order, and $\text{Var}(U_{(k)}) = \text{Var}(\Phi(Z_{(k)}))$ monotone in ρ is confirmed numerically (`check_fan.py`) but not by a transfer theorem.

The Gaussian case fixes the copula and varies k ; the opposite slice, the top level $k = n$ for *any* negatively associated sample, is also exactly tractable and is where the conjecture becomes one-dimensional. There $U_{(n)} \leq u$ iff every score lies below u , so the law of $U_{(n)}$ is the copula diagonal $\delta(u) = C(u, \dots, u) = \mathbb{P}(N(u) = n)$, and $\text{Var}(U_{(n)}) = \int_0^1 \int_0^1 [\delta(u \wedge v) - \delta(u)\delta(v)] du dv$. Negative association enters only as the negative-orthant bound $\delta(u) \leq u^n$ (Joag-Dev and Proschan, 1983) — equivalently the convex-order image of the discretely convex indicator $\mathbf{1}\{N = n\}$. Writing the deficit $g(u) = u^n - \delta(u) \geq 0$ and $\mu^* = n/(n+1) = \mathbb{E}[U_{(n)}^{\text{iid}}]$, the variance has the exact closed form

$$\text{Var}(U_{(n)}) = \text{Var}^{\text{iid}}(U_{(n)}) + 2 \int_0^1 (u - \mu^*) g(u) du - \left(\int_0^1 g \right)^2, \quad (12)$$

so the per-level conjecture at $k = n$ is *equivalent* to a centroid inequality on the deficit,

$$\frac{\int_0^1 u g(u) du}{\int_0^1 g(u) du} \leq \mu^* + \frac{1}{2} \int_0^1 g : \quad (13)$$

the mass by which the diagonal falls short of independence must sit, on average, at most a half-deficit above the independent-maximum mean. This is *not* a consequence of the diagonal alone. A linear program over the functions meeting every condition a copula diagonal must satisfy (monotone, $0 \rightarrow 1$, $\delta \leq u$, n -Lipschitz, Fréchet lower bound; Nelsen, 2006) together with the negative-orthant bound $\delta \leq u^n$ returns a diagonal that violates (13) (`check_fan.py`). The single level is therefore irreducibly a property of the off-diagonal negative-orthant probabilities across thresholds, not of the diagonal that determines $U_{(n)}$'s law: exactly the content the aggregate convex-order bound uses in bulk and that a one-level statement cannot read off. It is the same obstruction that blocks localizing Theorem 4 to a single k , now in closed form. At $n = 2$ the LP-violating diagonal is realised by the genuine NA law of Proposition 2, so (13) fails there; for $n \geq 3$ the off-diagonal constraints appear to rule such diagonals out, and every negatively associated structure tested satisfies (13) with margin.

The same convex-order contraction $N(u) \leq_{\text{cx}} \text{Bin}(n, u)$ that drives the aggregate bound is available per level, but turning it into the single- k statement needs the tail $\mathbb{P}(N(u) \geq k)$ to cross the binomial tail once in u , which the total positivity of the binomial tail in (k, u) (Karlin, 1968)

makes plausible, followed by a single-crossing-implies-less-spread step. We claim only the variance statement, and deliberately not the stronger dispersive ordering: an order statistic of a dependent sample need not be dispersively smaller than the independent one even at fixed marginals (Panja et al., 2022), so the result is specific to the variance and is not a corollary of a dispersive theorem. The natural sufficient condition on the dependence is negative supermodular dependence (Christofides and Vaggelatou, 2004), which all the structures below satisfy. The table is illustrative, each row a single Monte Carlo estimate (6×10^4 draws, so the variance entries carry a relative Monte Carlo error of a few percent), fully reproduced by `check_fan.py` ($n = 20$, $\alpha = 0.1$, independent fan variance 3.92×10^{-3}); it is evidence for the conjecture, not a proof.

calibration dependence	$\mathbb{E}[c]$	$\text{Var}(c)$	ratio to fan	sign
independent ($\rho = 0$)	0.904	3.92×10^{-3}	1.00	—
exchangeable, $\rho = -1/(n-1)$	0.923	1.16×10^{-3}	0.29	neg. assoc.
MA(1), lag-1 corr $-\frac{1}{2}$	0.912	2.71×10^{-3}	0.69	neg. assoc.
random negatively-paired blocks	0.909	3.14×10^{-3}	0.80	neg. assoc.
contest / without replacement	0.905	0	0.00	floor
one-factor, $\rho = +0.2$	0.863	1.33×10^{-2}	3.41	positive

Every negatively associated structure, exchangeable or not, comes in under the independent fan; the positive control widens it. The mean of c also responds to dependence, drifting up under negative and down under positive (to 0.863 in the one-factor row), confirming it is not pinned; the variance is what we track here. A wider search agrees: over exchangeable, moving-average, block, and randomly-generated non-exchangeable negatively-correlated Gaussian copulas at $n \leq 8$ and every k , the ratio to the independent fan never exceeds one (it ranges from about 0.30 at the floor to 0.80 for weak dependence), with no counterexample. In all of these the order-statistic CDFs $\mathbb{P}(N(u) \geq k)$ cross the independent ones exactly once, the single crossing the remark below uses.

Remark 3 (A closed form, and what an exact proof must dominate). *The variance is an explicit functional of $g(u) = \mathbb{P}(N(u) \geq k)$,*

$$\text{Var}(c) = 2 \int_0^1 (1-u) g(u) du - \left(\int_0^1 g(u) du \right)^2, \quad (14)$$

so writing $\delta = g - g^{\text{iid}}$, $A = \int_0^1 g^{\text{iid}} = 1 - \mathbb{E}[c^{\text{iid}}]$, and $\Delta = \int_0^1 \delta$,

$$\text{Var}(c) - \text{Var}(c^{\text{iid}}) = 2 \int_0^1 [(1-u) - A] \delta(u) du - \Delta^2. \quad (15)$$

The $-\Delta^2$ term is non-positive. The weight $(1-u) - A$ changes sign at $u = \mathbb{E}[c^{\text{iid}}] = k/(n+1)$. Empirically (across every structure in the table, and in the wider search below) δ changes sign exactly once, near $u \approx k/n$; we emphasize this is an observed single-crossing, not a consequence of Lemma 1, since count-variance contraction alone does not force one-crossing of the tail curve g . Granting that single crossing, the two sign changes agree up to the conformal offset between k/n and $k/(n+1)$, a band of width $O(1/n)$ on which the integrand is positive and elsewhere negative, so the identity gives the contraction up to that $O(1/n)$ boundary term, a second route to the leading order of Theorem 1, and isolates what a finite-sample proof must control: the boundary remainder against the slack Δ^2 .

Remark 4 (Why the closed form does not close it, and the lemma that would). *The closed form is, by itself, circular. Substituting the moment identities $\int u \delta = \frac{1}{2}(\mathbb{E}[c_{\text{iid}}^2] - \mathbb{E}[c^2])$ and $\int \delta = \mathbb{E}[c_{\text{iid}}] - \mathbb{E}[c]$*

into any elementary rearrangement of the difference, single-crossing splits, Abel summation, the exact first-order perturbation, collapses it back to $\text{Var}(c) - \text{Var}(c_{\text{iid}})$ with the auxiliary crossing point cancelling: these structural facts carry no magnitude. Nor is the per-level convex order of Lemma-1 enough on its own. Because $\text{Var}(c)$ depends only on the single curve $g(u) = \mathbb{P}(N(u) \geq k)$, one might hope $N(u) \leq_{\text{cx}} \text{Bin}(n, u)$ at each u suffices; it does not, since a curve dominated pointwise can have larger variance (place an atom at the boundary), and such curves are excluded only because they are not realizable by a genuine sample. The content is therefore an extremal statement about the realizable curves, not their marginals:

Open lemma ($n \geq 3$). Among all $g(u) = \mathbb{P}(N(u) \geq k)$ realizable by a negatively associated uniform sample of size $n \geq 3$, the independent (binomial) curve maximizes $\text{Var}(c)$.

The restriction to $n \geq 3$ is necessary: Proposition 2 realizes an over-dispersing curve at $n = 2$. This is what a total-positivity or coupling argument would have to deliver. Proposition 1 and the contest endpoint are the cases settled by hand.

Remark 5 (The signed measure as a negative between-term). Theorem 2 reads the fan as within-dataset variance plus a non-negative between-dataset variance. In the signed-de-Finetti picture of the companion note, the negative corner is exactly where that between-dataset object turns negative, a Feynman–Wigner quasi-variance that subtracts from the within-dataset fan rather than adding to it, reaching the full cancellation $R = 0$ at the contest floor. This is heuristic, since a signed measure has no genuine conditional variance, but it is the right cartoon: the same sign that decides conditional adaptivity in Cotton (b) decides whether the fan is inflated or cancelled here.

6 A constructive corollary

If negative dependence narrows the fan, one can engineer it, and there is a clean continuous-marginal construction that stays inside the $U_i \sim \text{Unif}(0, 1)$ model (unlike the discrete grid of Theorem 3). Take a one-per-stratum design: with π a uniform random permutation of $\{1, \dots, n\}$,

$$U_i = \frac{\pi(i) - 1 + V_i}{n} \quad (\text{Latin hypercube, } V_i \stackrel{\text{iid}}{\sim} \text{Unif}(0, 1)), \quad \text{or} \quad U_i = \frac{\pi(i) - 1 + T}{n} \quad (\text{common jitter, } T \sim \text{Unif}(0, 1)) \quad (16)$$

Each U_i is exactly $\text{Unif}(0, 1)$ and the sample is exchangeable, with one point per stratum, so the k -th order statistic lies in the k -th stratum and $\mathbb{E}[U_{(k)}] = (k - \frac{1}{2})/n$, $\text{Var}(U_{(k)}) = 1/(12n^2)$, a fan of order $1/n^2$ rather than $1/n$. (Note these continuous designs give mean $(k - \frac{1}{2})/n$, not the grid's exact $k/(n + 1)$.) More generally the classical variance-reduction designs apply: stratification never increases variance (Cochran, 1977), antithetic pairing helps for the monotone empirical CDF, and determinantal or repulsive designs give negative association with faster-than-independent concentration (Hough et al., 2006; Bardenet and Hardy, 2020).

Remark 6 (Admissible designs). The marginal guarantee needs each calibration score exchangeable with the test score, not the calibration scores independent of one another, so negative association is in principle compatible. But the routes differ in what they cost. Score-dependent curation of the calibration set, the obvious way to balance it, breaks exchangeability with the test point and voids the guarantee. A pooling that uses all $n + 1$ scores together (pooled ranks or de-meaning) preserves rank validity, but it is transductive, full-conformal in flavour, not split conformal with a threshold fixed before the test point is seen, and once the test score enters the pooled construction the realized-coverage variable is no longer the fixed-threshold $c = U_{(k)}$ studied here. The safe message

is the design one: a pre-specified, exchangeability-preserving randomized design can lower the run-to-run variance of the threshold without disturbing validity. What makes such a design admissible is not that it is merely randomized and pre-specified, but that it preserves the exchangeability the validity argument needs; covariate stratification qualifies only when it is external to the responses and applied symmetrically across the $n + 1$ points. Exact zero fan is the discrete rank/grid idealization; continuous-marginal designs (Latin hypercube) give a small but nonzero fan, of order $1/n^2$. The tension is intrinsic: a generic negatively associated design lowers the dispersion only by also nudging $\mathbb{E}[c]$ off $1 - \alpha$ (the mean is not pinned), so coverage that is both reproducible and on-target demands a mean-preserving symmetric design. Distribution-free validity and minimal coverage dispersion cannot both be driven to their limit at once, save in the degenerate zero-fan grid. A deeper caveat: the dispersion is a within-model quantity, computed with the marginal F and the copula C held fixed. Shrinking it by design lowers sampling variability but does nothing for model risk (whether F is the right target at all) and removes the run-to-run scatter that is the cheapest diagnostic of misspecification. A vanishing fan against an assumed F certifies reproducibility, not correctness; the zero-fan grid reads $1 - \alpha$ every draw by construction, not because the world obliges.

Remark 7 (The copula viewpoint). Because the scores are read on the uniform PIT scale, the joint law of $(U_1, \dots, U_n)$ is, by Sklar’s theorem, a copula C , with the marginal F factored out entirely. So $\text{Var}(c)$ is a copula functional: the fan is a copula invariant, identical for any score whose calibration dependence is C . The fan coefficient is read off the bivariate copula diagonal, $\bar{c} = \frac{1}{n(n-1)} \sum_{i \neq j} (C_{ij}(p, p) - p^2)$ at the operating quantile $p = 1 - \alpha$; for the maximum ($k = n$) $\text{Var}(c)$ is exactly a functional of the diagonal $u \mapsto C(u, u)$, since $\mathbb{P}(U_{(n)} \leq u) = C(u, u)$. The endpoints are the Fréchet bounds: independence Π gives the Beta fan, the comonotone copula M ($U_1 = \dots = U_n$) the maximal fan $\frac{1}{12}$; the negative side has no dual, as the countermonotone lower bound is not a copula for $n > 2$, which is why the floor is only the most negative exchangeable copula, $\rho = -1/(n - 1)$. Theorem 2 then says that the extendable copulas, the exchangeable Archimedean families (Clayton, Gumbel, Frank), each with a de Finetti frailty W as directing variable, carry a non-negative between- Q term and widen the fan; the negative regime is the signed-de-Finetti corner with no such mixture, which is why its contraction needed convex order rather than conditioning. Finally, tail dependence acts through the copula’s tails: a copula dependent in one tail pins the order statistics there and fattens the opposite end, so lower-tail dependence (Clayton) inflates the high, conformal quantile more than upper-tail dependence (Gumbel) — the reverse of the naive guess, as the companion demonstration shows.

7 The time-series fan

When the calibration scores are a stationary segment of a series (the deployment mode priced by Barber and Pananjady, 2026), the coupled-test caution of the setup applies: the coverage statement for the next step of the same series is theirs, not ours. What the machinery above supplies is the dispersion of the threshold $U_{(k)}$ across calibration segments, and with it the second moment of realized coverage against the stationary law. The mean shifts by at most the switch-coefficient tax, $|\mathbb{E}[c] - (1 - \alpha)|$ bounded through $\min_{\tau} \{\tau/(n + 1) + 2\beta(\tau)\}$ by their two-sided bounds. The variance is governed by the same covariance kernel as Theorem 1: in the Bahadur regime,

$$\text{Var}(c) = \frac{\sigma_{\text{LR}}^2}{n} + o(1/n), \quad \sigma_{\text{LR}}^2(u) = \sum_{j \in \mathbb{Z}} \text{Cov}(\mathbf{1}\{U_0 \leq u\}, \mathbf{1}\{U_j \leq u\}),$$

the long-run variance of the sub-threshold indicator process at the operating quantile — equivalently 2π times its spectral density at zero. For positively serially dependent scores $\sigma_{\text{LR}}^2 \geq \alpha(1 - \alpha)$ and

the fan widens by the constant factor $\sigma_{\text{LR}}^2/(\alpha(1-\alpha))$, which does not shrink with n . Realized coverage on a time-series calibration set is thus approximately $\mathcal{N}(1-\alpha-\text{tax}, \sigma_{\text{LR}}^2/n)$: the mean damage vanishes at rate $1/n$ while the dispersion carries a permanent inflation factor. Dependence therefore presents three separate charges: the vanishing mean tax, the fixed fan inflation, and the fixed sharpness rent of the companion note (Cotton, 2026). Only the first is reduced by more calibration data.

8 Related work

The Beta($k, n-k+1$) law of calibration-conditional coverage and its variance are Vovk (2012) and Marques F. (2025); to our knowledge the papers cited here do not study the dependence among calibration scores as a control variable for the realized-coverage variance. The loss-side generalization of the calibration-conditional law is Snell and Griffiths (2025), who put Dirichlet($1, \dots, 1$) weights on the ordered calibration losses via the Aitchison–Dunsmore quantile-spacing result and report a posterior over realized risk: with the 0–1 miscoverage loss their construction collapses to the Beta fan above, their recovery of conformal risk control is the fan’s *mean*, and their highest-posterior-density rule is the prescription “report the fan’s quantile, not its mean.” Their guarantee is what they call data-conditional (conditional on the calibration draw, our axis) and, as they note, not input-conditional; it moves along the fan, not across the information gap. Negative association and its consequences are Joag-Dev and Proschan (1983); that negatively associated sums are dominated in convex order by independent ones is Shao (2000), and sampling without replacement concentrates at least as tightly as with replacement (Serfling, 1974). The total positivity behind the single-crossing step is Karlin (1968); that an order statistic of a dependent sample need not be dispersively smaller than the independent one, which is why we claim only the variance, is Panja et al. (2022); the closest prior comparison of order statistics across dependence, Hu and Hu (1998), orders their *distribution functions* between associated and independent samples rather than the *variance* that is our object; the connection between supermodular ordering and negative dependence is Christofides and Vaggelatos (2004); the Bahadur representation is Bahadur (1966). Determinantal and repulsive designs are Hough et al. (2006); Bardenet and Hardy (2020) and stratification is Cochran (1977); Wiecek (2023) uses “design-based” conformal prediction in the survey-sampling sense—randomization-based validity under a complex sampling design—distinct from the variance-reduction designs here. The finite signed de Finetti representation is Kerns and Székely (2006), and the within-dataset, across- x reading of its sign is the companion note (Cotton, b); the information gap is (Cotton, a). The de Finetti directing measure behind the positive side is a prior in all but name; the Bayesian–conformal interface, where priors meet finite-sample coverage and the trade is validity against efficiency, is surveyed by Deliu and Liseo (2025), and the dispersion-versus-guarantee tension above is a variance-side instance of it. The contribution here is to make the dependence the control variable for the *variance* of realized coverage: the leading-order fan coefficient, the exact law-of-total-variance decomposition on the positive side, the exact aggregate contraction under negative association (Theorem 4), and, for equicorrelated normals, the exact and dispersively-monotone per-level law. The general single-level negative-association contraction is left as a conjecture. The copula reading (Remark 7) rests on Sklar’s theorem (Sklar, 1959; Nelsen, 2006); copulas enter conformal prediction elsewhere through the multivariate *output* (e.g. Sun and Yu, 2024), a different axis from the calibration-set copula here.

Appendix: the quantile-slope reduction for Proposition 4

With $Z_{(k)} = aD + b\eta$, $a = \sqrt{1 - \rho}$, $b = \sqrt{(1 + (n - 1)\rho)/n}$, the budget $(n - 1)a^2 + nb^2 = n$ holds and $da/d\rho = -1/(2a)$, $db/d\rho = (n - 1)/(2nb)$. Under the natural coupling (the same D, η for all ρ), $\dot{Y} := \partial_\rho Z_{(k)} = -D/(2a) + (n - 1)\eta/(2nb)$, and the quantile evolves by $\partial_\rho q_\rho(u) = \mathbb{E}[\dot{Y} \mid Y = q] =: M(q)$. Writing $m(q) = \mathbb{E}[D \mid Y = q]$ and $b\mathbb{E}[\eta \mid Y = q] = q - am(q)$,

$$M(q) = -\frac{m(q)}{2a} + \frac{n - 1}{2nb^2} (q - am(q)). \quad (17)$$

Differentiating and using $\frac{1}{2a} + \frac{(n-1)a}{2nb^2} = \frac{nb^2 + (n-1)a^2}{2anb^2} = \frac{1}{2ab^2}$ (the budget),

$$M'(q) = \frac{n - 1}{2nb^2} - \frac{m'(q)}{2ab^2}, \quad (18)$$

so $M'(q) \geq 0$, dispersive monotonicity ($\partial_\rho q_\rho$ increasing in q), is equivalent to $m'(q) \leq (n - 1)a/n$. Tweedie's identity $m'(q) = a \text{Var}(D \mid Y = q)/b^2$ turns this into the posterior-variance bound $\text{Var}(D \mid Y = q) \leq (n - 1)b^2/n$, which Lemma 2 and the Brascamp–Lieb inequality deliver, as in Proposition 4.

The strong-log-concavity constant in Lemma 2. The constant $n/(n - 1)$ is tracked as follows. On the hyperplane $H = \{\sum_i w_i = 0\}$ the centred vector w has $-\log$ density $\frac{1}{2}\|w\|^2$ (the covariance $I - \frac{1}{n}J$ restricts to the identity on H), so the measure is 1-strongly-log-concave there; restricting to the convex cone $\{w_1 \leq \dots \leq w_n\} \cap H$ adds a convex term to $-\log$ and keeps it 1-strongly-log-concave. Strong log-concavity is preserved under marginalization along a linear functional (Prékopa, 1973; Brascamp and Lieb, 1976; Saumard and Wellner, 2014): for an m -strongly-log-concave measure and $\ell(x) = \langle v, x \rangle$, the law of ℓ is $(m/\|v\|^2)$ -strongly-log-concave (the marginal along the unit direction $v/\|v\|$ is m -slc, and rescaling the argument by $\|v\|$ scales the constant by $1/\|v\|^2$). Here $D = w_{(k)} = \langle e_k, w \rangle$ on the cone, with $v = \text{proj}_H(e_k) = e_k - \frac{1}{n}\mathbf{1}$ and $\|v\|^2 = \frac{n-1}{n}$, so $m = 1$ gives the constant $n/(n - 1)$ (the cone restriction only raises it). Numerically the realized constant is larger still, $(\log g_D)'' \leq -2.5$ in the cases checked, so the bound holds with room.

References

- R. R. Bahadur. A note on quantiles in large samples. *The Annals of Mathematical Statistics*, 37(3): 577–580, 1966.
- R. F. Barber and A. Pananjady. Predictive inference for time series: why is split conformal effective despite temporal dependence? In *Proceedings of the 37th International Conference on Algorithmic Learning Theory (ALT)*, volume 313 of *Proceedings of Machine Learning Research*, pages 1–24, 2026. arXiv:2510.02471.
- R. Bardenet and A. Hardy. Monte carlo with determinantal point processes. *The Annals of Applied Probability*, 30(1):368–417, 2020. arXiv:1605.00361.
- H. J. Brascamp and E. H. Lieb. On extensions of the brunn–minkowski and prékopa–leindler theorems, including inequalities for log concave functions, and with an application to the diffusion equation. *Journal of Functional Analysis*, 22(4):366–389, 1976.
- T. C. Christofides and E. Vaggelatou. A connection between supermodular ordering and positive/negative association. *Journal of Multivariate Analysis*, 88(1):138–151, 2004.

- W. G. Cochran. *Sampling Techniques*. Wiley, 3rd edition, 1977.
- P. Cotton. Marginally useful: Formalizing the information gap in conformal prediction, a. Companion paper.
- P. Cotton. A feynman–wigner-style diagnostic for the efficacy of conformal prediction via signed de finetti representations, b. Companion note.
- P. Cotton. The two prices of dependence: a vanishing coverage tax and a fixed sharpness rent, 2026. Technical note, conformalprediction.net/papers/two-prices/.
- B. de Finetti. La prévision: ses lois logiques, ses sources subjectives. *Annales de l’Institut Henri Poincaré*, 7(1):1–68, 1937.
- N. Deliu and B. Liseo. The interplay between bayesian inference and conformal prediction, 2025. arXiv:2510.26930.
- B. Efron. Tweedie’s formula and selection bias. *Journal of the American Statistical Association*, 106(496):1602–1614, 2011.
- P. R. Hansen and C. Tong. Tweedie’s formula and score-driven updating, 2026. Preprint, arXiv:2605.15902.
- J. B. Hough, M. Krishnapur, Y. Peres, and B. Virág. Determinantal processes and independence. *Probability Surveys*, 3:206–229, 2006.
- T. Hu and J. Hu. Comparison of order statistics between dependent and independent random variables. *Statistics & Probability Letters*, 37(1):1–6, 1998.
- K. Joag-Dev and F. Proschan. Negative association of random variables, with applications. *The Annals of Statistics*, 11(1):286–295, 1983.
- S. Karlin. *Total Positivity, Vol. 1*. Stanford University Press, 1968.
- G. J. Kerns and G. J. Székely. De finetti’s theorem for abstract finite exchangeable sequences. *Journal of Theoretical Probability*, 19(3):589–608, 2006.
- E. L. Lehmann. Some concepts of dependence. *The Annals of Mathematical Statistics*, 37(5):1137–1153, 1966.
- T. Lewis and J. W. Thompson. Dispersive distributions, and the connection between dispersivity and strong unimodality. *Journal of Applied Probability*, 18(1):76–90, 1981.
- P. C. Marques F. Universal distribution of the empirical coverage in split conformal prediction. *Statistics & Probability Letters*, 219:110350, 2025. arXiv:2303.02770.
- R. B. Nelsen. *An Introduction to Copulas*. Springer, 2nd edition, 2006.
- A. Panja, P. Kundu, and B. Pradhan. Dispersive and star ordering of sample extremes from dependent random variables following the proportional odds model, 2022. Preprint, arXiv:2006.04454.
- A. Prékopa. On logarithmic concave measures and functions. *Acta Scientiarum Mathematicarum*, 34:335–343, 1973.

- A. Saumard and J. A. Wellner. Log-concavity and strong log-concavity: a review. *Statistics Surveys*, 8:45–114, 2014.
- R. J. Serfling. Probability inequalities for the sum in sampling without replacement. *The Annals of Statistics*, 2(1):39–48, 1974.
- Q.-M. Shao. A comparison theorem on moment inequalities between negatively associated and independent random variables. *Journal of Theoretical Probability*, 13(2):343–356, 2000.
- A. Sklar. Fonctions de répartition à n dimensions et leurs marges. *Publications de l’Institut de Statistique de l’Université de Paris*, 8:229–231, 1959.
- J. C. Snell and T. L. Griffiths. Conformal prediction as bayesian quadrature. In *Proceedings of the 42nd International Conference on Machine Learning (ICML)*, 2025. arXiv:2502.13228.
- S. H. Sun and R. Yu. Copula conformal prediction for multi-step time series forecasting. In *ICLR*, 2024. arXiv:2212.03281.
- V. Vovk. Conditional validity of inductive conformal predictors. In *Asian Conference on Machine Learning*, volume 25 of *PMLR*, pages 475–490, 2012.
- J. Wiecek. Design-based conformal prediction. *Survey Methodology*, 49(2), 2023. arXiv:2303.01422.