

Marginally Useful

Conformal Prediction Certifies the Coverage of a Set,
Not the Quality of a Distribution

Peter Cotton*

June 1, 2026

Abstract

Conformal prediction supplies a genuine and valuable guarantee: under exchangeability it gives finite-sample, distribution-free *marginal* coverage for a set. This paper argues that this guarantee is often read as something it is not — evidence of distributional forecast quality. The two are different objects, graded by different instruments. We first review three limitations of the marginal guarantee: it is not conditional coverage, non-trivial conditional coverage is unavailable distribution-free without additional structure, and it depends on exchangeability. We then make the separation precise with three small contributions. First, an *orthogonality* statement: for a residual-based score the conformal set is determined by residual order statistics and is not a functional of any accompanying predictive density, so its coverage can be held fixed while the density’s log-score varies arbitrarily. Second, a *residual-information gap*. For a fixed location predictor, the expected log-score regret of a single-shape residual predictive system relative to the oracle is the expected Kullback–Leibler divergence of the conditional residual law from its marginal — equivalently the mutual information $I(R; X)$ — for a signed-residual conformal predictive system, and $I(R; X) + \text{KL}(\bar{r} \parallel h_{\text{sym}})$ for ordinary absolute-residual interval conformal, which additionally forfeits any marginal residual skewness. This gap cannot be reduced by any recalibration that ignores X ; only conditioning on X removes it. Third, a *coverage-score diagnostic* that displays conformalization as a correction of set coverage rather than an improvement of distributional sharpness. The conclusion is not that conformal prediction is weak, but that it is exact for a different objective: controlling whether the truth lies in a reported region.

1 Introduction

A practitioner maintains a probabilistic forecaster and is told, repeatedly, that conformal prediction is the principled, distribution-free way to “add uncertainty.”¹ They bolt it on. The number they actually report — a proper score such as the predictive log-likelihood — does not improve. The

*Microprediction. peter.cotton@microprediction.com. This paper accompanies the interactive demonstrations at *Marginally Useful*; every quantitative claim below has a slider-driven counterpart there.

¹The reading is pervasive, and not only in blogs. Applied and survey writing routinely calls the marginal-coverage guarantee “well-calibrated” or “reliable” uncertainty quantification — the word *calibrated* standing in for coverage (Bellotti and Zhao, 2025) — and tutorial expositions go further, presenting conformal prediction as a way to “add uncertainty” (tds, a), the one thing you need for it (tds, b), or even to “predict full probability distributions” (Manokhin). The core *methodological* literature is more careful — it separates *validity* from *efficiency* — and a growing critique strand makes the present point independently: that valid-but-uninformative sets have little practical value (Mehrtens et al., 2025), and that interval length can be gamed while marginal coverage is preserved (Min et al., 2026), a rediscovery of the trivial-validity argument of Section 3. Our quarrel is with the conflation, not with conformal prediction.

natural diagnosis is a tuning failure. The correct diagnosis, we argue, is a *category error*: conformal prediction is a method for *certifying the coverage of a set*, and it is being asked to *estimate a distribution*. These are different kinds of object, graded by different instruments, and no amount of tuning converts one into the other.

None of this impugns the mathematics. Given exchangeable data, a fitted predictor, and a nonconformity score, split conformal prediction returns a set $C_\alpha(x)$ with

$$\mathbb{P}(Y_{n+1} \in C_\alpha(X_{n+1})) \geq 1 - \alpha, \quad (1)$$

finite-sample and with no assumption on the data distribution (Vovk et al., 2005; Lei et al., 2018; Angelopoulos and Bates, 2023). The trouble is entirely in the reading of (1). The probability averages over X_{n+1} as well as Y_{n+1} : it is a statement about the *average* future input, not about the one in front of you. It is *marginal* coverage, and as the foundational analyses warned from the start, “a good estimator must satisfy something more than marginal coverage” (Lei and Wasserman, 2014).

Contributions and structure. Sections 2–3 are expository: the guarantee, and the three senses in which it is weaker than advertised. Our contributions are:

- **Orthogonality** (Section 4, Proposition 1). For a residual-based score the conformal set is not a functional of the predictive density, so marginal coverage places no constraint on the log-score of an accompanying density: one can pin coverage at $1 - \alpha$ and send the log-score to $-\infty$. (*Folklore made precise.*)
- **The residual-information gap** (Section 5, Propositions 2–3). A single-shape residual predictive system re-levels the base model’s residual shape; its log-score regret to the oracle is the mutual information $I(R; X) = \mathbb{E}_X \text{KL}(r(\cdot | X) \| \bar{r})$ for the signed-residual conformal predictive system, and $I(R; X) + \text{KL}(\bar{r} \| h_{\text{sym}})$ for the usual absolute-residual intervals. This gap is untouched by any recalibration that ignores X ; only conditioning on X reduces it. Within the single-shape class the conformal system is log-score optimal. (*A known re-leveling identity made two-case; a decomposition we have not seen stated this way.*)
- **The coverage–score plane** (Section 6). A two-axis diagnostic in which, for a report consisting of a set and an unchanged density, conformalization is a horizontal projection onto the zero-coverage-error line, leaving the score coordinate untouched. (*A presentation device we find clarifying.*)

Section 7 treats exchangeability and time series; Section 8 gives the honest converse — when coverage really is the objective — and Section 9 bounds the strong claims against the modern conformal literature (conformal predictive systems, conformal training, and information-theoretic and conditional-coverage advances), each of which buys its gains exactly by the conditional modeling our analysis identifies.

2 Setup: split conformal and the marginal guarantee

Let $(X_i, Y_i)_{i=1}^{n+1}$ be exchangeable. Fix a point predictor $\hat{\mu}$ trained on a disjoint set, and a *nonconformity score* $s(x, y)$; the canonical choice in regression is the absolute residual $s(x, y) = |y - \hat{\mu}(x)|$. Compute calibration scores $s_i = s(X_i, Y_i)$ for $i = 1, \dots, n$ and let

$$\hat{q} = s_{(k)}, \quad k = \lceil (n+1)(1 - \alpha) \rceil, \quad (2)$$

the k -th order statistic (with $\hat{q} = +\infty$ when $k > n$). The split-conformal set is

$$C_\alpha(x) = \{y : s(x, y) \leq \hat{q}\} = [\hat{\mu}(x) - \hat{q}, \hat{\mu}(x) + \hat{q}] \quad (3)$$

for the absolute-residual score. Exchangeability of the augmented scores makes the rank of s_{n+1} uniform, which yields (1) (Vovk et al., 2005; Lei et al., 2018). The construction is elegant and the guarantee is real; the demonstrations confirm that the empirical coverage tracks $1 - \alpha$ as the sample size, noise level, and α are varied. We do not dispute any of this. We dispute what it is taken to mean.

3 Three ways the marginal guarantee is weaker than it looks

(i) Marginal is not conditional, and conditional is impossible for free. One would prefer *conditional* coverage, $\mathbb{P}(Y \in C(x) | X = x) \geq 1 - \alpha$ for (almost) every x . Distribution-free, this cannot be bought. Lei and Wasserman (2014) show (their Lemma 1) that any band with non-trivial finite-sample conditional validity has *infinite* expected length at almost every point of a continuous distribution. Foygel Barber et al. (2021) restate and sharpen this: a conditionally valid C_n has $\mathbb{E} \text{leb}(C_n(x)) = \infty$ at almost all non-atomic x (their Prop. 2.2), and even the relaxed demand of coverage on every subgroup of probability $\geq \delta$ “is impossible to attain beyond the trivial solution” of inflating the marginal level to $1 - \alpha\delta$ (their Thm. 3.1). We collect both as the operative no-go result.

Lemma 1 (No-go for distribution-free conditional coverage). *Let P have non-atomic X . (a) (Lei and Wasserman, 2014; Foygel Barber et al., 2021) Any C_n with non-trivial finite-sample conditional validity has $\mathbb{E} \text{leb}(C_n(x)) = \infty$ at almost every x . (b) (Foygel Barber et al., 2021) Requiring $\mathbb{P}(Y \in C_n(X) | X \in G) \geq 1 - \alpha$ for every measurable G with $P(X \in G) \geq \delta$ is attainable only by a C_n whose marginal level is inflated to $1 - \alpha\delta$ — a uniformly wider, non-adaptive band.*

These are not loose bounds to be tightened; they are the reason the gap cannot be closed for free, and both surface as concrete finite-sample facts in the demonstrations. Part (a) drives Demo 7: localizing split conformal to recover per- x coverage shrinks each cell’s calibration count n_b until $[(n_b + 1)(1 - \alpha)] > n_b$, at which point (2) gives $\hat{q} = +\infty$ and the only valid cell interval is $(-\infty, \infty)$ — the infinite expected length of the lemma, arriving as arithmetic. Part (b) is Demo 8: the distribution-free protection of all δ -subgroups is exactly the flat band at level $1 - \alpha\delta$, whose width diverges as $\delta \rightarrow 0$ while never adapting to x , at a multiplicative width cost over an assumption-driven adaptive oracle.

So the guarantee you can have distribution-free is the marginal one, and a marginally valid band “tends to overestimate the set when x is in the high density area and to underestimate for low density x ” (Lei and Wasserman, 2014) — it over-covers the easy inputs and under-covers the hard ones, silent by construction about the cases the model was built to get right. The heteroscedastic demonstration (Demo 2) exhibits exactly this: 90% overall, near-100% where the noise is small, well below target where it is large.

To be clear, the conformal literature is not unaware of this distinction. The standard vocabulary already separates *validity* from *efficiency* and adaptivity, and modern methods such as conformalized quantile regression (Romano et al., 2019) deliberately use better conditional modeling to obtain more adaptive sets. The narrower claim here is that the distribution-free marginal coverage *certificate* should not be mistaken for distributional forecast quality — and that, as the next sections make precise, whatever adaptivity such methods gain is bought by conditional modeling, not by the conformal step itself.

(ii) **Validity is trivially satisfiable.** A predictor that returns the whole outcome space with probability $1 - \alpha$ and the empty set otherwise satisfies (1) exactly and conveys nothing. More practically, the *same* coverage attaches to an excellent $\hat{\mu}$ and to a deliberately terrible one — the latter merely yields a wider $\hat{q}$ in (2). Marginal validity is therefore not evidence of a good method; it certifies the fence, not the cattle. The next section makes this precise.

(iii) **Without exchangeability, even the marginal guarantee fails.** Equation (1) rests on exchangeability of the augmented sample. Under covariate shift, label shift, or temporal dependence it does not hold, and the patches that restore something do so by weakening what is guaranteed (Section 7).

4 Coverage and score are orthogonal

We first make “coverage is not a measure of forecast quality” a precise, constructive fact. Model a forecaster as a pair $(f(\cdot|x), s)$ of a predictive density and a nonconformity score. The point is conditional on the score: it is sharp precisely when the density is *not* used to build s (as for the canonical residual score), and it does not apply to genuinely density- or quantile-based scores.

Proposition 1 (Coverage does not constrain the log-score). *The split-conformal set (3) is a measurable function of the nonconformity scores $\{s_i\}$ and of $s(x, \cdot)$; it is not a functional of the predictive density f unless f is explicitly used in the score. Consequently, if two forecasters share the same score function s and the same calibration scores, their conformal sets — and hence their marginal coverage at every level α — are identical, while their expected log-scores $\mathbb{E}[\log f(Y|X)]$ may differ by an arbitrary amount.*

Proof. The first claim is immediate from (2)–(3): $\hat{q}$ is an order statistic of $\{s_i\}$ and $C_\alpha(x) = \{y : s(x, y) \leq \hat{q}\}$, so two forecasters agreeing on s and on $\{s_i\}$ produce the same set for every x, α , and (1) concerns only that set.

For the gap, take X degenerate and $Y \sim N(0, 1)$; let $\hat{\mu} \equiv 0$ with the absolute-residual score $s(x, y) = |y|$, which does not depend on the predictive density at all. For any claimed predictive density $f_\sigma = N(0, \sigma^2)$ the scores are $|Y_i|$, independent of σ , so the conformal set is the fixed interval $[-\hat{q}, \hat{q}]$ with $\hat{q}$ the empirical $(1 - \alpha)$ -quantile of $|Y|$. Yet

$$\mathbb{E}[\log f_\sigma(Y)] = -\frac{1}{2} \log(2\pi\sigma^2) - \frac{\mathbb{E}[Y^2]}{2\sigma^2} = -\frac{1}{2} \log(2\pi\sigma^2) - \frac{1}{2\sigma^2} \xrightarrow{\sigma \rightarrow 0^+} -\infty,$$

and likewise $\rightarrow -\infty$ as $\sigma \rightarrow \infty$. Hence for any $M > 0$ there are σ_1, σ_2 with identical conformal sets (identical marginal coverage at every α) yet log-scores differing by more than M . $\square$

Corollary 1. *For a residual-score conformal predictor, marginal validity certifies an order statistic of residual magnitudes. It does not identify, bound, or rank the scale, shape, or sharpness of any predictive density that was not used to construct the score; in particular, no inference about forecast quality (as graded by a strictly proper score) can be drawn from coverage alone.*

This is the rigorous form of a point made informally by several authors — that conformal validity “almost invites you to use garbage prediction functions” (Recht, 2024) — and it is what the orthogonality demonstration shows visually: the claimed density’s spread slides freely while the conformal interval and its 90% coverage do not move, because the spread never entered (2).

5 The category error, made precise: the sharpness gap

Forecast quality decomposes into *calibration* and *sharpness*, and the discipline’s stated goal is to maximize sharpness subject to calibration, assessed by a proper scoring rule (Gneiting et al., 2007; Gneiting and Raftery, 2007). We show where conformal prediction sits in this decomposition, and we do so fairly: within the class of objects it can produce, it is optimal; the loss is the class.

Conformal predictive systems. A conformal predictor can be made to output a distribution rather than a set — the *conformal predictive system* (CPS) of Vovk et al. (2019), built from *signed* residual scores. This is the fairest possible target for the criticism, because it is the mode in which conformal prediction estimates a full distribution. We must, however, distinguish it from the more common practice of reading the nested symmetric intervals of ordinary *absolute*-residual split conformal as if they were a distribution; the two re-level different objects, and the distinction matters for the gap below.

Proposition 2 (Re-leveling, two cases). *Fix a location predictor $\hat{\mu}$ with residual $R = Y - \hat{\mu}(X)$ of marginal density g and CDF G . In the large-calibration limit (under exchangeability):*

(A) **Signed-residual CPS.** *The CPS predictive distribution is $\Pi_x(y) = G(y - \hat{\mu}(x))$ (the calibration ECDF $\hat{G}_n \rightarrow G$ by Glivenko–Cantelli), with density $\pi_x(y) = g(y - \hat{\mu}(x))$: the base location forecast re-leveled by the marginal signed-residual law.*

(B) **Absolute-residual intervals.** *Ordinary split conformal at level α returns the symmetric interval $\hat{\mu}(x) \pm \hat{q}_\alpha$, with $\hat{q}_\alpha$ the $(1 - \alpha)$ -quantile of $|R|$. Read as a predictive distribution (by sweeping α) it is symmetric about $\hat{\mu}(x)$ with $|\cdot|$ distributed as $|R|$; its density is the symmetrized marginal residual law $h_{\text{sym}}(z) = \frac{1}{2}(g(z) + g(-z))$.*

Proof sketch. (A) The CPS at level α returns endpoints equal to the signed-residual order statistics; sweeping α traces $\hat{\mu}(x)$ plus the residual quantile function, i.e. $\hat{G}_n^{-1}$ shifted by $\hat{\mu}(x)$, so $\Pi_x(y) = \hat{G}_n(y - \hat{\mu}(x))$; take limits (Vovk et al., 2019). (B) The intervals depend on the data only through $|R|$ and are symmetric about $\hat{\mu}(x)$; the unique symmetric density whose absolute value has the law of $|R|$ (with signed marginal g) is $h_{\text{sym}}(z) = \frac{1}{2}(g(z) + g(-z))$. $\square$

In both cases the estimation was done by the base model and the residual sample; conformal supplies only the leveling. We now quantify the cost. Write the true conditional residual density as $r(\cdot | x)$, so the oracle density is $q^*(y | x) = r(y - \hat{\mu}(x) | x)$, and let $\bar{r} = \mathbb{E}_X r(\cdot | X) = g$ be the marginal residual density.

Proposition 3 (Residual-information gap). *Fix the location predictor $\hat{\mu}$ and write $I(R; X) = \mathbb{E}_X \text{KL}(r(\cdot | X) \| \bar{r}) \geq 0$, the mutual information between the residual and the input. The expected log-score regret relative to the oracle q^* is*

$$(A) \text{ signed CPS: } \mathbb{E}[\log q^*(Y | X)] - \mathbb{E}[\log \pi_X(Y)] = I(R; X), \quad (4)$$

$$(B) \text{ absolute intervals: } \mathbb{E}[\log q^*(Y | X)] - \mathbb{E}[\log h_{\text{sym}}(R)] = I(R; X) + \text{KL}(\bar{r} \| h_{\text{sym}}). \quad (5)$$

Both are ≥ 0 . The term $I(R; X)$ vanishes iff $R \perp X$; the extra term $\text{KL}(\bar{r} \| h_{\text{sym}})$ vanishes iff the marginal residual law is symmetric. Among all single-shape forecasters $y \mapsto h(y - \hat{\mu}(x))$, $\mathbb{E}[\log h(R)]$ is maximized at $h = \bar{r}$; the signed CPS attains this optimum. Hence no recalibration that ignores X can reduce $I(R; X)$ — it can at best replace the shape by $\bar{r}$, removing the skewness term in (5) and reducing case (B) to case (A). Reducing $I(R; X)$ itself requires conditioning on X .

Proof. With $R = Y - \hat{\mu}(X)$, $\log q^*(Y | X) = \log r(R | X)$. (A) With $\pi_X(Y) = g(R) = \bar{r}(R)$, $\mathbb{E}[\log r(R | X) - \log \bar{r}(R)] = \mathbb{E}_X \mathbb{E}_{R|X} \log \frac{r(R|X)}{\bar{r}(R)} = \mathbb{E}_X \text{KL}(r(\cdot | X) \| \bar{r}) = I(R; X)$, the definition of mutual information since $\bar{r}$ is the marginal of R . (B) Add and subtract $\log \bar{r}(R)$: $\mathbb{E}[\log r(R | X) - \log h_{\text{sym}}(R)] = I(R; X) + \mathbb{E}_{R \sim \bar{r}} \log \frac{\bar{r}(R)}{h_{\text{sym}}(R)} = I(R; X) + \text{KL}(\bar{r} \| h_{\text{sym}})$. For the optimum, $\mathbb{E}[\log \bar{r}(R)] - \mathbb{E}[\log h(R)] = \text{KL}(\bar{r} \| h) \geq 0$ (Gibbs), so $h = \bar{r}$ is best among shared shapes; a single shape cannot represent X -dependence in r , so it cannot change $I(R; X)$. $\square$

Remark 1 (What this says, and does not say). The right-hand side of (4) is exactly the information about the residual distribution that remains in X after the location predictor is fixed. Calling this the residual-information gap makes the content of “blind to sharpness” precise. Three honest caveats. (1) *Conformal is not dominated within its class.* The signed CPS is log-score optimal among single-shape location forecasters; the cost is the restriction, not the calibration step. The absolute-residual interval system is the one that pays the extra $\text{KL}(\bar{r} \| h_{\text{sym}})$, for discarding sign information. (2) *What recalibration can and cannot do.* A recalibration that ignores X (e.g. marginal PIT recalibration) can move the shape toward $\bar{r}$ and thereby erase the skewness term, i.e. turn case (B) into case (A); it cannot touch $I(R; X)$. Reducing $I(R; X)$ requires modeling conditional residual shape — heteroscedastic models, conditional recalibration (Kuleshov et al., 2018), or conformalized quantile regression (Romano et al., 2019), which is the conformal world’s way of leaving the single-shape class. (3) *The distribution-free guarantee is real.* Recalibration offers calibration only in expectation or asymptotically and carries no coverage certificate; conformal’s marginal guarantee is finite-sample and assumption-light. When the certificate itself is the deliverable, that is decisive (Section 8).

The recalibration demonstration realizes the special case in which $\hat{\mu}$ is correct and $r(\cdot | x)$ is a fixed Gaussian with the wrong scale ($I(R; X) = 0$): variance recalibration fit to the log-score recovers the shape and improves the score, while conformalization re-levels to exact coverage and returns a set.

6 The coverage–sharpness plane

The preceding results suggest a single diagnostic, provided we are careful about what is being plotted. Define a *reporting system* as a set-valued report $C_\alpha(x)$ and, optionally, a density-valued report $f(\cdot | x)$. Its coordinates are

$$(C, S), \quad C = |\mathbb{P}(Y \in C_\alpha(X)) - (1 - \alpha)|, \quad S = \mathbb{E}[\log f(Y | X)], \quad (6)$$

the marginal-coverage error of the *set* and the proper-score performance of the *density*. The decomposition of forecast quality into calibration and sharpness (Gneiting et al., 2007) is, in this plane, the statement that the two coordinates are distinct, and our results read geometrically:

- **Residual split conformal is a horizontal projection.** It maps a report $(C_\alpha^{\text{base}}, f)$ to $(C_\alpha^{\text{conf}}, f)$: it changes the set so that $C \rightarrow 0$ while leaving the density f — and hence S — untouched (Proposition 1). It moves a point left, never up.
- **Conformal predictive systems are different:** they supply their own density-valued report, so they do have a vertical coordinate — but that coordinate is governed by the single-shape restriction of Proposition 3, capped at the oracle minus $I(R; X)$.

- **Recalibration and better modeling move vertically.** Fitting shape to a proper score raises S ; conditioning on x is the only way to raise it past the ceiling set by the residual-information gap (4).

A system reported by its coverage alone is reported by one coordinate of a two-coordinate quality. The plane is a presentation device — adjacent to reliability and sharpness diagrams (Gneiting et al., 2007) — but the “conformal = move left, never up” reading makes the category distinction hard to miss, and it is the through-line of the demonstrations. The companion time-series benchmark plots methods in exactly this plane (worst-window coverage against interval score).

7 Exchangeability and the time-series case

Drop exchangeability and (1) is no longer guaranteed. The literature’s repairs are ingenious, and uniform in one respect: they survive by redefining what is guaranteed.

- **Weighted conformal prediction** (Tibshirani et al., 2019) restores marginal coverage under covariate shift when the likelihood ratio is known; with estimated weights coverage becomes approximate and the effective sample size drops.
- **Adaptive Conformal Inference** (Gibbs and Candès, 2021) updates the level online and “puts no constraints on the data generating distribution,” but guarantees a *long-run time-average* miscoverage frequency, $\frac{1}{T} \sum_t \text{err}_t \rightarrow \alpha$ — not a finite-sample statement about $\mathbb{P}(Y \in C(X))$, and certainly not conditional coverage.
- **EnbPI** (Xu and Xie, 2021) drops exchangeability for approximate marginal coverage under strong-mixing/stationarity surrogates; later online methods (Zaffran et al., 2022; Gibbs and Candès, 2024) retreat to regret bounds over local intervals. The general beyond-exchangeability analysis of Barber et al. (2023) bounds the coverage gap by a total-variation distance that is exactly what nonstationarity inflates.

It is worth separating three distinct guarantees that the word “conformal” now spans: (i) *split conformal under exchangeability* gives finite-sample marginal coverage; (ii) *adaptive conformal inference* gives long-run, time-average coverage control under distribution shift; (iii) *online conformal under arbitrary shifts* gives regret-style or local-interval performance rather than the original exchangeable guarantee. These are progressively weaker and more diffuse objects; none is the sharp predictive distribution *now*, for *this* step, that a proper score grades. The exchangeability required by guarantee (i) is typically unavailable in deployed time-series settings, though blocking or learned-transform constructions can sometimes recover an exchangeability-like assumption.

Our companion benchmark bears this out, and is fair to the methods. On a nonstationary series with known σ_t , fixed split conformal collapses far below target once drift sets in, but the genuinely adaptive methods — ACI, AgACI, conformal PID control, and weighted/NexCP — recover marginal coverage and are competitive on the interval (Winkler) score; conformal PID attains the smallest conditional-coverage gap of any method. They are not straw men. The honest finding is the one the theory predicts: a simple probabilistic volatility model (an EWMA-variance Gaussian) matches or beats them on the proper score *and* returns a full distribution, and where the conformal methods help they do so precisely by adapting the interval to recently realized residuals — that is, by local conditional modeling, with the conformal step supplying the coverage leveling on top.

8 When coverage *is* the objective

The argument is skeptical, not dismissive, and it comes with a precise converse. The litmus is one question:

*Is your loss a function of **whether** the truth lands in a region, or of **where** it lands?*

If the former, coverage is the native object and conformal prediction is sound and often ideal:

- **Selective prediction and risk control.** “Return a small set guaranteed to contain the label 95% of the time; a human checks it.” The deliverable is a set; risk-controlling prediction sets and conformal risk control formalize this (Angelopoulos et al., 2021; Bates et al., 2021; Angelopoulos et al., 2024).
- **Retrieval and shortlisting**, where the product is a candidate set and coverage is recall.
- **Anomaly and novelty detection via conformal p -values** — conformal prediction’s cleanest home, because here it is not a forecaster but a distribution-free hypothesis test and coverage is Type-I error control (Vovk et al., 2005).
- **Compliance and long-run frequency control**, where a certificate of the form “contained 95% of the time on average” *is* the product, and the distribution-free, finite-sample nature of (1) is exactly the value added.

Two caveats keep even these honest. If you trust a calibrated posterior you can form a sharper set yourself — the smallest region of mass $\geq 1 - \alpha$ — and recover the expectations a bare set discards; conformal’s marginal value is then purely the distribution-free guarantee, useful when you distrust the density and exchangeability holds. And even when coverage is the goal you usually want it conditionally, which returns us to the impossibility of Section 3.

9 Scope, and the constructive escapes

Our strong claims target the object that is actually over-sold: *post-hoc, marginal, split conformal prediction with a residual score*. A large and active literature builds conformal methods that do more, and it is worth saying precisely how they relate, because in every case they confirm the paper’s mechanism rather than contradict it: they buy sharpness or conditional coverage by *conditioning on X or optimizing a proper objective*, which is exactly what Propositions 3 say is required.

Conformal predictive systems that condition on X . Mondrian / binned conformal predictive systems (Boström et al., 2021) fit a separate residual law per region, and Toccaceli (2026) chooses the bins by minimizing a leave-one-out CRPS. These genuinely produce sharp, shape-adaptive predictive distributions — and they do so by leaving the single-shape class of Proposition 2 (a per-bin shape is X -dependent) and, in the CRPS case, by making the partition criterion a proper score. This is the most pointed case for the defence, and the honest concession is real: when the reference class is itself selected by a proper scoring rule, the conformal *construction* is performing conditional density estimation. The category distinction does not vanish; it relocates to “which component estimates the distribution,” and the answer is the conditional partition, not the coverage certificate — the value is in the estimation, as throughout. Conformalized quantile regression (Romano et al., 2019) is the same story with a quantile model in place of the bins.

End-to-end conformal training. The orthogonality of Proposition 1 is a statement about *post-hoc* conformalization of a fixed predictor. Conformal training (Stutz et al., 2022; Correia et al., 2024) deliberately differentiates through the conformalizer to minimize expected set size subject to coverage, which couples the coverage and efficiency axes by construction. This does not contradict the proposition; it illustrates its corollary — to obtain sharpness you must optimize for it, rather than read it off the coverage certificate.

The information in set size. Correia et al. (2024) show that the expected size of a calibrated conformal set lower-bounds the conditional entropy $H(Y | X)$. This is complementary to our Proposition 3: both say conformal *geometry* carries information. It also sharpens the two-coordinate plane of Section 6 — the *efficiency* (set-size / sharpness) coordinate is the information-bearing one, while the marginal *coverage* coordinate, by Proposition 1, is not. Reporting only the latter discards exactly the coordinate that encodes aleatoric uncertainty.

Approaching conditional coverage. The no-go result of Section 3 concerns *exact, distribution-free, finite-length* conditional coverage. It is not the end of the subject. Diagnostics such as the excess-risk-of-target-coverage family (Braun et al., 2025) quantify and decompose the conditional gap, and post-processing via pivotal scores and optimal transport (Laplante, 2026) attains *approximate* conditional coverage by estimating the conditional law of the score — again, conditioning on X , and explicitly under the structural assumptions the impossibility theorem rules out for the exact, assumption-free case. These approach the goalpost; they do not move the one the no-go result defends.

Modern time-series conformal. Beyond the adaptive methods benchmarked in Section 7, methods such as HopCPT (Auer et al., 2023) exploit temporal structure to tighten intervals and improve empirical adaptivity. Their guarantees remain of the approximate / long-run kind catalogued there; the improvement, again, comes from modeling the conditional error law.

Beyond the log-score. We used the log-score for the clean information-theoretic identities, but the coverage–score plane and the argument carry over to any strictly proper score; the companion benchmark in fact ranks methods by the interval (Winkler) score and CRPS (Gneiting and Raftery, 2007), on which the same orderings hold.

10 Conclusion

Conformal prediction certifies the coverage of a set, finite-sample and distribution-free, under exchangeability. That guarantee is genuine and, for the right objective, exact. The error is reading it as distributional forecast quality. For a fixed location predictor the guarantee you can have (marginal) is rarely the one you want (conditional); the conditional one is unavailable distribution-free without further structure; the output is a set, not a measure; its log-score loss relative to the oracle is precisely the residual conditional-information term $I(R; X)$ of (4); and the exchangeability the basic guarantee needs is typically unavailable in deployed time-series settings. Within the class of objects it produces it is optimal, and where a guarantee is the product it is the right tool. But coverage answers a question — *is the truth in this region?* — that, in forecasting, is often not the one you were asking. Conformal prediction is not weak uncertainty quantification; it is exact uncertainty quantification for a set-valued coverage objective, and mistaking a set-coverage certificate for evidence of distributional quality is the category error.

Reproducibility. Every numerical claim has an interactive counterpart in the accompanying *Marginally Useful* demonstrations; the orthogonality of Section 4, the gap of Section 5, and the projection of Section 6 are each a slider away.

References

- How to add uncertainty estimation to your models with conformal prediction. Towards Data Science, a. <https://towardsdatascience.com/how-to-add-uncertainty-estimation-to-your-models-with-conformal-prediction-a5acdb86ea05/>.
- All you need is conformal prediction. Towards Data Science, b. <https://towardsdatascience.com/all-you-need-is-conformal-prediction-726f18920241/>.
- Anastasios N. Angelopoulos and Stephen Bates. Conformal prediction: A gentle introduction. *Foundations and Trends in Machine Learning*, 16(4):494–591, 2023.
- Anastasios N. Angelopoulos, Stephen Bates, Jitendra Malik, and Michael I. Jordan. Uncertainty sets for image classifiers using conformal prediction. In *International Conference on Learning Representations (ICLR)*, 2021.
- Anastasios N. Angelopoulos, Stephen Bates, Adam Fisch, Lihua Lei, and Tal Schuster. Conformal risk control. In *International Conference on Learning Representations (ICLR)*, 2024.
- Andreas Auer, Martin Gauch, Daniel Klotz, and Sepp Hochreiter. Conformal prediction for time series with modern hopfield networks. In *Advances in Neural Information Processing Systems (NeurIPS)*, 2023. arXiv:2303.12783.
- Rina Foygel Barber, Emmanuel J. Candès, Aaditya Ramdas, and Ryan J. Tibshirani. Conformal prediction beyond exchangeability. *The Annals of Statistics*, 51(2):816–845, 2023.
- Stephen Bates, Anastasios Angelopoulos, Lihua Lei, Jitendra Malik, and Michael I. Jordan. Distribution-free, risk-controlling prediction sets. *Journal of the ACM*, 68(6):1–34, 2021.
- Anthony Bellotti and Xindi Zhao. Conformal prediction and trustworthy AI. In *Artificial Intelligence and Machine Learning (Springer)*. 2025. arXiv:2508.06885; “well-calibrated uncertainty quantification”.
- Henrik Boström, Ulf Johansson, and Tuwe Löfström. Mondrian conformal predictive distributions. In *Conformal and Probabilistic Prediction and Applications (COPA)*, PMLR, volume 152, pages 24–38, 2021.
- Sacha Braun, David Holzmüller, Michael I. Jordan, and Francis Bach. Conditional coverage diagnostics for conformal prediction. *arXiv preprint arXiv:2512.11779*, 2025.
- Alvaro H.C. Correia, Fabio Valerio Massoli, Christos Louizos, and Arash Behboodi. An information theoretic perspective on conformal prediction. In *Advances in Neural Information Processing Systems (NeurIPS)*, 2024. arXiv:2405.02140.
- Rina Foygel Barber, Emmanuel J. Candès, Aaditya Ramdas, and Ryan J. Tibshirani. The limits of distribution-free conditional predictive inference. *Information and Inference: A Journal of the IMA*, 10(2):455–482, 2021.
- Isaac Gibbs and Emmanuel J. Candès. Adaptive conformal inference under distribution shift. In *Advances in Neural Information Processing Systems (NeurIPS)*, volume 34, pages 1660–1672, 2021.
- Isaac Gibbs and Emmanuel J. Candès. Conformal inference for online prediction with arbitrary distribution shifts. *Journal of Machine Learning Research*, 25(162):1–36, 2024.
- Tilman Gneiting and Adrian E. Raftery. Strictly proper scoring rules, prediction, and estimation. *Journal of the American Statistical Association*, 102(477):359–378, 2007.

- Tilman Gneiting, Fadoua Balabdaoui, and Adrian E. Raftery. Probabilistic forecasts, calibration and sharpness. *Journal of the Royal Statistical Society: Series B*, 69(2):243–268, 2007.
- Volodymyr Kuleshov, Nathan Fenner, and Stefano Ermon. Accurate uncertainties for deep learning using calibrated regression. In *International Conference on Machine Learning (ICML)*, PMLR, volume 80, pages 2796–2804, 2018.
- Félix Laplante. A post-processing conformal prediction approach for conditional coverage via pivotal scores. *arXiv preprint arXiv:2605.25852*, 2026.
- Jing Lei and Larry Wasserman. Distribution-free prediction bands for non-parametric regression. *Journal of the Royal Statistical Society: Series B*, 76(1):71–96, 2014.
- Jing Lei, Max G’Sell, Alessandro Rinaldo, Ryan J. Tibshirani, and Larry Wasserman. Distribution-free predictive inference for regression. *Journal of the American Statistical Association*, 113(523):1094–1111, 2018.
- Valeriy Manokhin. Predicting full probability distributions with conformal prediction. Medium. <https://valeman.medium.com/predicting-full-probability-distributions-with-conformal-prediction-1dd4c1f26973>.
- Hendrik Meertens, Tabea Bucher, and Titus J. Brinker. Pitfalls of conformal predictions for medical image classification. *arXiv preprint arXiv:2506.18162*, 2025.
- Yizhou Min, Yizhou Lu, Lanqi Li, Zhen Zhang, and Jiaye Teng. Questioning the coverage-length metric in conformal prediction: When shorter intervals are not better. *arXiv preprint arXiv:2601.21455*, 2026.
- Benjamin Recht. Cover songs; and from intervals to bands. *arg min* blog, 2024. <https://www.argmin.net/>.
- Yaniv Romano, Evan Patterson, and Emmanuel J. Candès. Conformalized quantile regression. In *Advances in Neural Information Processing Systems (NeurIPS)*, volume 32, 2019.
- David Stutz, Krishnamurthy Dvijotham, Ali Taylan Cemgil, and Arnaud Doucet. Learning optimal conformal classifiers. In *International Conference on Learning Representations (ICLR)*, 2022. arXiv:2110.09192.
- Ryan J. Tibshirani, Rina Foygel Barber, Emmanuel J. Candès, and Aaditya Ramdas. Conformal prediction under covariate shift. In *Advances in Neural Information Processing Systems (NeurIPS)*, volume 32, 2019.
- Paolo Toccaceli. CRPS-optimal binning for univariate conformal regression. *arXiv preprint arXiv:2603.22000*, 2026.
- Vladimir Vovk, Alexander Gammerman, and Glenn Shafer. *Algorithmic Learning in a Random World*. Springer, New York, 2005.
- Vladimir Vovk, Jieli Shen, Valery Manokhin, and Min-ge Xie. Nonparametric predictive distributions based on conformal prediction. *Machine Learning*, 108(3):445–474, 2019.
- Chen Xu and Yao Xie. Conformal prediction interval for dynamic time-series. In *International Conference on Machine Learning (ICML)*, PMLR, volume 139, pages 11559–11569, 2021.
- Margaux Zaffran, Olivier Féron, Yannig Goude, Julie Josse, and Aymeric Dieuleveut. Adaptive conformal predictions for time series. In *International Conference on Machine Learning (ICML)*, PMLR, volume 162, pages 25834–25866, 2022.