

MACROSCOPIC

Two specimens received.

1. [REDACTED] Melanoma left mid back nick superior: An orientated skin ellipse measuring 70x35x15 mm. The skin surface reveals a tan centrally located macule measuring 16x10 mm. There is a nick superior which is orientated to 12 o'clock. The 3 o'clock margin is inked blue, the 9 o'clock margin is inked black. Sectioned from 12 o'clock to 6 o'clock. 1A-1F 1TS per block. Entire lesion submitted. Refer to photographic diagram.
2. [REDACTED] Right postauricular lesion: An unorientated skin ellipse measuring 22x8x3 mm. The skin surface reveals a tan macule measuring 10x3 mm. 2A 2TS; 2B 1TS. Entire lesion submitted. (L Whitham/jmt14).

MICROSCOPIC

1. Melanoma left mid back nick superior: Sections show a MELANOMA IN SITU of superficial spreading type which arises in mildly sun damaged skin. The lesion is broad and shows some architectural features of dysplastic naevus but single cells predominate over nests, there are zones of confluent lentiginous growth and scattered pagetoid areas. The lesion arises in association with a benign INTRADERMAL NAEVUS. A malignant dermal component is not seen. There is a zone of established regression up to 5 mm diameter. A separate intradermal naevus is also seen. In situ melanoma is well clear of all margins with peripheral clearances of at least 10 mm.
2. Right postauricular lesion: Sections show finely granular pigment which ranges from dusky yellow to brown to black encrusting elastotic fibres within the dermis and superficial subcutis. The pigment is not associated with a cellular proliferation and is most likely exogenous. There is no pigmentation is completely excised.

SUMMARY

1. Melanoma left mid back nick superior: MELANOMA IN SITU OF SUPERFICIAL SPREADING



[REDACTED]

[REDACTED]

TYPE (CLARK LEVEL 1) ARISING IN ASSOCIATION WITH AN INTRADERMAL NAEVUS. EXCISION COMPLETE.

2. Right postauricular lesion: PIGMENT ENCRUSTING ELASTIC FIBRES WITHIN THE DERM AND SUBCUTIS. EXCISION COMPLETE.

[REDACTED]

[REDACTED]