

DIAGNOSTIC SUMMARY:

1. LEFT CHEST SUPERIOR LATERAL, EXCISION - LOW GRADE DYSPLASTIC JUNCTIONAL NAEVUS, CLEAR OF THE MARGIN.
2. RIGHT NECK POSTERIOR LATERAL BASE 2, EXCISION - LOW GRADE DYSPLASTIC LENTIGINOUS JUNCTIONAL NAEVUS, CLEAR OF THE MARGIN.
3. RIGHT CHEST MID LATERAL, BIOPSY - SCAR, NO EVIDENCE OF MALIGNANCY.

Clinical Notes: 1) L chest sup lat; e/o PSL next to old M scar, dermatologist wants lesion removed for HD, ?LGDN - 12 nick; 2) R neck post lat base punch excision, hidden in creases, duration unknown, atypical, exclude M - 12 nicks; 3) R chest mid lat pbx, asymptomatic, ?BCC.

Macroscopic:

1. Labelled 'L chest sup lat'. A 16 x 5 x 4mm skin ellipse. A nick marks one transverse edge, orientated 12 o'clock (12 o'clock green, 6 o'clock black). There is a tan lesion, 2 x 2mm. 1A ITS 1LS 3 o'clock; 1B ITS 1LS 9 o'clock.
2. Labelled 'R neck post lat base 2'. A 10 x 7 x 3mm skin ellipse. A suture marks one tip, orientated 12 o'clock (3 o'clock green, 9 o'clock black). There is a tan crusted and raised lesion, 5 x 5mm. 2A ITS 1LS 12 o'clock; 2B ITS 1LS 6 o'clock.
3. Labelled 'R chest mid lat 3'. A 4 x 6mm skin punch with a tan surface. 3A bisected in total.

Microscopic:

1. Sections show a low grade dysplastic junctional naevus with regression, clear of the margin.
2. Sections show a low grade dysplastic lentiginous junctional naevus

[REDACTED]

with focal regression, clear of the margin.

3. Sections show a scar. I can see no evidence of malignancy.

[REDACTED]