Conditional probability

Practice problems

9/23/24

Please work on the practice problems in your group. At least one of the following problems will be assigned to the weekly problem set.

- 1. Suppose we have an event A from one random process and an event B from a second random process such that P(A) = 0.3, P(B) = 0.7, and $P(A \cap B) = 0.1$.
 - a. Are the random processes independent?
 - b. What is P(A|B)?
- 2. Assortative mating is a nonrandom mating pattern where individuals with similar genotypes and/or phenotypes mate with one another more frequently than what would be expected under a random mating pattern. Researchers studying this topic collected data on eye colors of 204 Scandinavian men and their female partners. The table below summarizes the results. For simplicity, we only include heterosexual relationships in this exercise.

		Partner (female)			
		Blue	Brown	Green	Total
Self (male)	Blue	78	23	13	114
	Brown	19	23	12	54
	Green	11	9	16	36
	Total	108	55	41	204

- a. What is the probability that a randomly chosen male respondent or his partner has blue eyes?
- b. What is the probability that a randomly chosen male respondent with blue eyes has a partner with blue eyes?
- c. What is the probability that a randomly chosen male respondent with brown eyes has a partner with blue eyes? What about the probability of a randomly chosen male respondent with green eyes having a partner with blue eyes?
- d. Does it appear that the eye colors of male respondents and their partners are independent? Explain your reasoning.

3. After an introductory statistics course, 80% of students can successfully construct box plots. Of those who can construct box plots, 86% passed, while only 65% of those students who could not construct box plots passed. Calculate the probability that a student is able to construct a box plot if it is known that he passed.