

## Conditional Variance

1. Suppose you see a total of  $N \sim \text{Geom}(s)$  movies in your lifetime. Further suppose that for each movie, you have a probability  $p$  of liking the movie, independently of other movies and of  $N$ . Let  $T$  be the number of movies you like in your lifetime.
  - (a) Find the mean of  $T$ .
  - (b) Find the variance of  $T$ .
2. Emails arrive one at a time in an inbox. Let  $T_j$  be the time at which the  $j$ -th email arrives. Suppose the waiting times between emails are iid  $\text{Exp}(\lambda)$  random variables (i.e.  $T_1, T_2 - T_1, T_3 - T_2, \dots$  are iid  $\text{Exp}(\lambda)$ ).

Each email is non-spam with probability  $p$  and spam with probability  $1 - p$ , independently of other emails and of the waiting times. Let  $X$  be the time at which the first non-spam email arrives (so  $X$  is a continuous random variable). For example,  $X = T_1$  if the first email is non-spam,  $X = T_2$  if the first email is spam and the second email is non-spam, etc.

*Hint for both parts: Let  $N$  be the number of emails we require to see the first non-spam (including that non-spam email) and write  $X$  as a sum of  $N$  terms. Hopefully, we believe  $N$  is independent of when a particular email arrives.*

- (a) Find the mean of  $X$ .
- (b) Find the variance of  $X$ .