

This problem set covers material from Week 9, dates 4/13- 4/15. Unless otherwise noted, all problems are taken from the textbook. Problems can be found at the end of the corresponding subsection.

Instructions: Write or type complete solutions to the following problems and submit answers to the corresponding Canvas assignment. Your solutions should be neatly-written, show all work and computations, include figures or graphs where appropriate, and include some written explanation of your method or process (enough that I can understand your reasoning without having to guess or make assumptions)

Monday 4/13

1. Section 9.1: problem 10 (note you did problem 8 on previous homework.)
2. Let $X_1, \dots, X_n | \theta \stackrel{\text{iid}}{\sim} \text{Exp}(\theta)$. We would like to test the hypotheses

$$H_0 : \theta \geq 0.01 \quad H_1 : \theta < 0.01$$

The rejection region is $R = \{\bar{X} \geq c\}$ for some $c > 0$. We observe $\bar{X} = 120$ from $n = 40$ data points. What is our p-value? Obtain this in two ways: exactly, and via a CLT approximation.

3. In the following, we are still interested in testing hypotheses about a parameter in a statistical model. But rather than try to identify where in Ω the parameter lives, we will test whether two models share the same parameter:

Let $X_1 \sim \text{Binom}(n_1, p_1)$ independent of $X_2 \sim \text{Binom}(n_2, p_2)$. We would like to test the hypotheses:

$$H_0 : p_1 = p_2 = p \quad \text{versus} \quad H_1 : p_1 < p_2$$

We will walk through obtaining the form of the p-value for this test.

- (a) Obtain the joint PMF of (X_1, X_2) under H_0 .
- (b) Define $T = X_1 + X_2$. Do smaller or larger values of X_1 (relative to the fixed total $T = t$) lend support in favor of H_1 ? Why?
- (c) Derive the form of the conditional distribution of $X_1 | T = t$. *You should be able to name it!*
- (d) Suppose we observe $X_1 = x_{obs}$. Find an explicit expression for the p-value.

Wednesday 4/15

TBD

General rubric

Points	Criteria
5	The solution is correct <i>and</i> well-written. The author leaves no doubt as to why the solution is valid.
4.5	The solution is well-written, and is correct except for some minor arithmetic or calculation mistake.
4	The solution is technically correct, but author has omitted some key justification for why the solution is valid. Alternatively, the solution is well-written, but is missing a small, but essential component.
3	The solution is well-written, but either overlooks a significant component of the problem or makes a significant mistake. Alternatively, in a multi-part problem, a majority of the solutions are correct and well-written, but one part is missing or is significantly incorrect.
2	The solution is either correct but not adequately written, or it is adequately written but overlooks a significant component of the problem or makes a significant mistake.
1	The solution is rudimentary, but contains some relevant ideas. Alternatively, the solution briefly indicates the correct answer, but provides no further justification.
0	Either the solution is missing entirely, or the author makes no non-trivial progress toward a solution (i.e. just writes the statement of the problem and/or restates given information).
Notes:	For problems with multiple parts, the score represents a holistic review of the entire problem. Additionally, half-points may be used if the solution falls between two point values above.
Notes:	For problems with code, well-written means only having lines of code that are necessary to solving the problem, as well as presenting the solution for the reader to easily see. It might also be worth adding comments to your code.