

# CSCI 570 - Fall 2021 - HW 10

Due November 18th

## Graded

1. (20 pts) Consider the partial satisfiability problem, denoted as 3-Sat( $\alpha$ ). We are given a collection of  $k$  clauses, each of which contains exactly three literals, and we are asked to determine whether there is an assignment of true/false values to the literals such that at least  $\alpha k$  clauses will be true. Note that 3-Sat(1) is exactly the 3-SAT problem from lecture.

Prove that 3-Sat(15/16) is NP-complete.

Hint: If  $x$ ,  $y$ , and  $z$  are literals, there are eight possible clauses containing them:  $(x \vee y \vee z)$ ,  $(\neg x \vee y \vee z)$ ,  $(x \vee \neg y \vee z)$ ,  $(x \vee y \vee \neg z)$ ,  $(\neg x \vee \neg y \vee z)$ ,  $(\neg x \vee y \vee \neg z)$ ,  $(x \vee \neg y \vee \neg z)$ ,  $(\neg x \vee \neg y \vee \neg z)$

**Solution:**

To prove it's in NP: given a truth value assignment, we can count how many clauses are satisfied and compare it to  $15k / 16$ .

To prove it's NP-hard:

We will show that  $3\text{-SAT} \leq_p 3\text{-SAT}(15/16)$ . For each set of 8 original clauses, create 8 new clauses using 3 new variables, and construct the clauses considering the collection of all possible clauses on the 3 new variables. If the number of clauses is a multiple of 8, then we are done: any assignment will satisfy 7/8 of the new clauses, so we must satisfy all of the original clauses in a valid solution to satisfy exactly 15/16 of the clauses. If the number of clauses is not a multiple of 8, we will satisfy more than 15/16 of the clauses, but if even one of the original clauses is not satisfied, we will have satisfied less than 15/16 of the clauses.

So, 3-Sat(15/16) is in the intersection of NP and NP-hard which is the class NP-Complete

2. (20 pts) Consider modified SAT problem, SAT' in which given a CNF formula having  $m$  clauses in  $n$  variables  $x_1, x_2, \dots, x_n$ , the output is YES if there is an assignment to the variables such that exactly  $m - 2$  clauses are satisfied, and NO otherwise. Prove that SAT' is also NP-Complete.

**Solution:**

To show that SAT' is NP-Complete,

First we will show that  $SAT' \in NP$ :

Given the assignment values as certificate, we can evaluate the SAT' instance and verify if it is satisfied. This is same as the SAT-verification. Moreover, we can count the number of satisfied clauses and check if it is equal to  $m - 2$  in linear time.

Next, we show that  $SAT \leq_p SAT'$ :

Construction: Add four more clauses  $x_1, x_2, \neg x_1, \neg x_2$  to the original SAT instance.

Claim: CNF formula obtained for SAT', F' has an assignment which satisfies SAT' iff CNF formula of SAT, F has an assignment which satisfies SAT.

$\Rightarrow$ ) if F has an assignment which satisfies SAT, then F' has an assignment which satisfies SAT'

Proof: If an assignment  $x_1 \dots x_n$  satisfies F, then it satisfies exactly two of the four extra clauses, giving exactly  $m + 2$ , which is nothing but  $m' - 2$  satisfied clauses for the F'.

$\Leftarrow$ ) if F' has an assignment which satisfies SAT', then F has an assignment which satisfies SAT

Proof: By construction, the only unsatisfied clauses for F must be one of  $x_1$  or  $\neg x_1$  and one of  $x_2$  or  $\neg x_2$ , so all the original  $m$  clauses are satisfied.

3. (20 pts) Given a graph  $G=(V,E)$  and two integers  $k, m$ , the *Dense Subgraph Problem* is to find a subset  $V'$  of  $V$ , whose size is at most  $k$  and are connected by at least  $m$  edges. Prove that the *Dense Subgraph Problem* is NP-Complete.

Solution:

Proving this problem NP is trivial, So here we show how to prove it's NP-Completeness.

We prove that the Independent set problem  $\leq_p$  Dense Subgraph Problem.

Given a graph  $G(V, E)$  and an integer  $k$ , an independent set decision problem outputs yes, if the graph contains an independent set of size  $k$ . For an arbitrary graph  $G=(V, E)$  of  $n$  vertices, we first get the complementary graph  $G_c$  of  $G$ .

A clique is a subset of vertices of an undirected graph  $G$  such that every two distinct vertices in the clique are adjacent; that is, its induced subgraph is

complete. We also know that an independent set in  $G$  is a clique in  $G_c$  (the complement graph of  $G$ ) and vice versa.

Then we set  $m$  to  $k*(k-1)/2$  and test with the dense subgraph problem.

Claim: There exists an independent set of size  $k$  in  $G$  (equivalently, a clique in  $G_c$  of size  $k$ ), iff there exists a subgraph of  $G_c$  with at most  $k$  vertices and at least  $m = k*(k-1)/2$  edges.

$\Rightarrow$ ) If there exists a clique in  $G_c$  of size at least  $k$ , then there exists a subgraph of  $G_c$  with at most  $k$  vertices and at least  $k*(k-1)/2$  edges.

If there is a clique of size at least  $k$  then there is a clique of size exactly  $k$ . Moreover, by definition, a clique of size  $k$  would have  $k * (k-1)/2$  edges.

$\Leftarrow$ ) If there exists a subgraph of  $G_c$  with at most  $k$  vertices and at least  $k*(k-1)/2$  edges, then there is a clique of size at least  $k$ .

For a subgraph to have  $k * (k-1)/2$  edges, implies there are  $k$  vertices. So this subset with  $k$  vertices forms a clique in  $G_c$  of size  $k$ .

## Ungraded

4. (20 pts) (Modified from Textbook 8.16) Consider the problem of reasoning about the identity of a set from the size of its intersections with other sets. You are given a finite set  $U$  of size  $n$ , and a collection  $A_1 \dots A_m$  of subsets of  $U$ . You are also given numbers  $c_1 \dots c_m$ , and numbers  $d_1 \dots d_m$ . The question is: Does there exist a set  $X \subseteq U$  so that for each  $i = 1 \dots m$ , the cardinality of  $X \cap A_i$  is larger than  $c_i$  but smaller than  $d_i$ ? We will call this an instance of the *Intersection Inference Problem*, with input  $U$ ,  $\{A_i\}$ , and  $\{c_i\}$ ,  $\{d_i\}$ . Prove that Intersection Inference is NP-complete.

**Solution:**

To show this problem is NP-Complete, we first prove it's NP.

Given the set  $X \subseteq U$ , we can enumerate all  $B_i$  in the collection, calculate the intersection with  $X$ , and compare its cardinality with  $c_i$  and  $d_i$ . So this problem is NP.

Then, we prove that the vertex cover  $\leq_P$  Intersection Inference.

For an arbitrary graph  $G = (V, E)$  of  $n$  vertices, we let  $V$  to be set  $U$ , and construct  $n+1$  subsets  $A_1 \dots A_{n+1}$  by:  $A_i$  ( $1 \leq i \leq n$ ) consists of all adjacent nodes of  $v_i \in V$ , and  $A_{n+1} = U$ . Then We set  $c_1 \dots c_n$  to 1,  $d_1 \dots d_n$  to infinity,  $c_{n+1}$  to 0,  $d_{n+1}$  to  $k$ . The solution of the intersection inference problem with this inputs is exactly the vertex cover of  $G$  of size no more than  $k$ .

5. (20 pts) (Textbook 8.28) The following is a version of the independent Set Problem. you are given a graph  $G = (V, E)$  and an integer  $k$ . For this problem, we will call a set  $I \subset V$  *strongly independent* if, for any two nodes  $v, u \in I$ , the edge  $(v, u)$  does not belong to  $E$ , and there is also no path of two edges from  $u$  to  $v$ , that is, there is no node  $w$  such that both  $(u, w) \in E$  and  $(w, v) \in E$ . The Strongly independent Set Problem is to decide whether  $G$  has a strongly independent set of size at least  $k$ .

Prove that the Strongly independent Set Problem is NP-complete.

**Solution:**

To show this problem is NP-Complete, we first prove it's NP.

Given a vertex subset  $V' \subseteq V$ , we can check in polynomial time if any pair of vertices  $u, v \in V'$  are either connected by an edge in  $E$  or by another node  $w \in V$ . Thus we can determine if a vertex subset is a strong independent set in polynomial time.

Then, We prove that Independent Set  $\leq_P$  Strong Independent Set.

For an arbitrary graph  $G = (V, E)$ , we split every edge in  $E$  into two nodes and connect them by a new node  $w_e$  (that means, an edge  $(u, v)$  in  $E$  becomes two edges  $(u, w_e)$  and  $(v, w_e)$ ). Then we connect all  $w_e$  nodes pairwise. Mathematically, we generate a new graph  $G_+ = (V_+, E_+)$  where  $V_+ = V \cup \{w_e \mid e \in E\}$ , and  $E_+ = \{(u, w_e), (v, w_e) \mid e = (u, v), e \in E\} \cup \{(w_e, w_{e'}) \mid e, e' \in E\}$ . To determine if there's an independent set in  $G$  of size at least  $k$ , we can check if there's any strong independent set in  $G_+$  with the same condition. Then:

1. If there is not such set: It's easy to prove that any independent set of  $G$  is also a strong independent set of  $G_+$ , so there's also no independent set of size at least  $k$  in  $G$ .

2. If there is such set: Let  $V_+'$  be that set. Since  $V_+'$  is a strong independent set of  $G_+$ , if any node in  $x \in \{w_e \mid e \in E\}$  is in  $V_+'$ , then  $V_+'$  can only contain  $x$  (because all other nodes in  $V_+$  are all not further than 2 to  $x$ ). In this case,  $|V_+'| = 1$ , and it's trivial to find an independent set of size 1 in  $G$ . On the other hand, if  $V_+'$  don't intersect with  $\{w_e \mid e \in E\}$ , then  $V_+' \subseteq V$ , making it an

independent set of  $V$ . In both cases, there's an independent set in  $G$  of size at least  $k$ .