

ICPSR 26841

**National Survey of Midlife in the
United States (MIDUS Refresher):
Daily Stress Project, 2012-2014**

Carol D. Ryff

University of Wisconsin-Madison

David M. Almeida

The Pennsylvania State University

Saliva Collection

Inter-university Consortium for
Political and Social Research
P.O. Box 1248
Ann Arbor, Michigan 48106
www.icpsr.umich.edu

Terms of Use

The terms of use for this study can be found at:
<http://www.icpsr.umich.edu/cocoon/ICPSR/TERMS/26841.xml>

Information about Copyrighted Content

Some instruments administered as part of this study may contain in whole or substantially in part contents from copyrighted instruments. Reproductions of the instruments are provided as documentation for the analysis of the data associated with this collection. Restrictions on "fair use" apply to all copyrighted content. More information about the reproduction of copyrighted works by educators and librarians is available from the United States Copyright Office.

NOTICE

WARNING CONCERNING COPYRIGHT RESTRICTIONS

The copyright law of the United States (Title 17, United States Code) governs the making of photocopies or other reproductions of copyrighted material. Under certain conditions specified in the law, libraries and archives are authorized to furnish a photocopy or other reproduction. One of these specified conditions is that the photocopy or reproduction is not to be "used for any purpose other than private study, scholarship, or research." If a user makes a request for, or later uses, a photocopy or reproduction for purposes in excess of "fair use," that user may be liable for copyright infringement.



MIDUS Refresher Project 2:

National Study of *Daily Experiences*

Wave 1: Saliva Collection

For file:
MR_P2_DAILYDATA_N782_20200702.sav

July, 2020

Address correspondence to David M. Almeida, Department of Human Development and Family Studies, The Pennsylvania State University, 403 Biobehavioral Health Building, University Park, Pennsylvania 16802 (email: dalmeida@psu.edu). This project was supported by the Network on Successful Mid-Life Development of the John D. and Catherine T. MacArthur Foundation and Grants R01 AG19239 and P01 AG0210166-02, from the National Institutes of Health.

National Study of *Daily Experiences*

Refresher Wave 1: Saliva Collection

Description

In the Refresher wave of NSDE, participants were asked to provide four saliva samples on days 2 through 5 in addition to completing short telephone interviews about their daily experiences across eight consecutive days. Saliva samples were collected at waking (before getting out of bed), 30 minutes after getting out of bed, before lunch, and before bed. Participants were instructed to collect samples before eating, drinking, brushing their teeth, or smoking. In addition, participants were asked not to consume any caffeinated products (e.g., coffee, tea, soda, or chocolate) before collecting samples. Data on the exact time respondents provide each saliva sample was obtained from the nightly telephone interviews and from a paper-pencil log sent with the collection kit. Prescription and over-the-counter medications taken during the collection period were recorded during the telephone interview of the last day of saliva collection.

When all 16 tubes were ready to be returned, participants used a pre-addressed, paid courier package for shipping to the MIDUS Biological Core at the University of Wisconsin, where they were stored in an ultracold freezer at -60° C. For analysis, the salivettes are thawed and centrifuged at 3000 rpm for 5 minutes, yielding a clear fluid with low viscosity. Cortisol concentrations are quantified with a commercially available luminescence immunoassay (IBL, Hamburg, Germany), with intra-assay and inter-assay coefficients of variation below 5%.

Of the 782 respondents who completed the Refresher Wave of the NSDE, 671 (85.8%) provided saliva samples.

Saliva and Cortisol Variables

In addition to the saliva and cortisol information described below, saliva and cortisol variables are also outlined in MR_P2_DailyDairyInstrument_20160906.pdf.

A. The first set of saliva- and cortisol-related variables allows for a descriptive examination of the saliva and cortisol data.

RA2DCORT: denotes whether a participant provide any saliva samples (1=yes, 2=no). Additionally, a value of 3 is denoted if a participant said they provided saliva but no samples were received. This is a between-person variable.

RA2DCDAY: denotes at least one valid cortisol sample on a given day for participants who provided saliva (1=yes, 2=no, 8=refused/missing). This is a daily variable.

RA2DCWDAY: denotes whether a participant provided saliva samples outside of the designated saliva collection days 2 through 5 (1=yes, 2=no, 8=refused/missing). For example, a participant could have started saliva collection on day 1, rather than day 2, and ended collection on day 4, rather than day 5. This is a between-person variable.

B. Raw Cortisol Values

The raw cortisol values for the four occasions are represented by RA2DCORW (cortisol at waking), RA2DCORA (30 minutes after waking), RA2DCORL (before lunch), and RA2DCORB (before bed). Cortisol values are in nanomoles per liter (nmol/l). No boundary restrictions are placed on the cortisol values, such that values as low as .00020 nmol/l and as high as 201241.60829 nmol/l are observed in the data.

In addition to actual raw values, the following values denote some type of non-response.

999994. UNRELIABLE
999995. NOT DONE
999996. EMPTY
999998. REFUSED/MISSING

Unreliable includes zero and unreliable values. Not done means the sample is missing. Empty reflects that a sample was sent to the lab, it was spun down, and there was little or no saliva in the bottom of the tube.

C. Cortisol Collection Time

Cortisol collection time variables (RA2DCORWT, RA2DCORAT, RA2DCORLT, and RA2DCORBT) appear in 24-hour military time. Collection times outside of 0-to-24 hour window occur in cases where a participant collects a saliva sample particularly early (i.e., prior to the beginning of interview day) or late (i.e., after the end of interview day). For example, an early awakening time of 11 p.m. would result in RA2DCORWT value of -1.00. Similarly, a relatively late bedtime of 2 a.m. would result in RA2DCORBT value of 26.00.

Two sources of data provide information on cortisol collection times: 1) paper-pencil log, as completed by the participant, and 2) cortisol collection times obtained by the interviewer from the participant during the telephone interview. Participants fill out the appropriate time for each cortisol collection occasion on the paper-pencil log and mail it along with their completed saliva samples. In addition, the interviewer records each cortisol collection time during interview days that are relevant to saliva collection.

Interviewer-obtained saliva collection time is a secondary source of cortisol collection time information, which is used to supplement paper-pencil log cortisol collection time. Specifically, participant-provided paper-pencil log times are first used to create RA2DCORWT, RA2DCORAT, RA2DCORLT, and RA2DCORBT variables. In cases where one or more paper-pencil log cortisol collection times are not available, the interviewer-obtained saliva collection times are entered into these variables instead.

Cortisol Collection Time Variables:

RA2DCORWT: CORTISOL COLLECTION TIME AT WAKING

RA2DCORAT: CORTISOL COLLECTION TIME 30 MINUTES AFTER WAKING

RA2DCORLT: CORTISOL COLLECTION TIME BEFORE LUNCH

RA2DCORBT: CORTISOL COLLECTION TIME BEFORE BED

98. REFUSED/MISSING

D. Medication Use During Saliva Collection

Information on use of over-the-counter and prescription medications relevant to cortisol levels is obtained on the last day of saliva collection. Participants endorse use of seven types of medication (see below) with a 1 (yes) or 2 (no) response. Typically, these questions are administered on day 5 of the NSDE interview, unless the participant finishes saliva collection early (e.g., day 4) or late (e.g., day 6).

Medication Use Variables:

During the days you provided saliva, did use you use any of the following medications?

RA2DMED1: STEROID NASAL SPRAY
(e.g., Flonase, Nasonex, Rinocort, Nasarel, Nasacort, Omnaris, Beconase, Veramyst)

RA2DMED2: INHALED STEROID MEDICATION
(e.g., Advair, Aerobid, Symbicort, Asmanax, Azmacort, Flovent, Pulmicort, Qvar)

RA2DMED3: ORAL STEROID MEDICATION
(E.G., PREDNISONE)

RA2DMED4: CREAM OR OINTMENT CONTAINING CORTISONE
(e.g., CORTAID)

RA2DMED5: CORTICOSTEROID INJECTION
(e.g., epidural corticosteroid injection, joint corticosteroid injection)

RA2DMED6: ANTI-DEPRESSANTS OR ANTI-ANXIETY MEDICATION

RA2DMED7: OTHER HORMONAL MEDICATIONS

1. YES
2. NO
7. DON'T KNOW
8. REFUSED/MISSING
9. INAPPLICABLE

E. Alpha-Amylase Values

RA2DAA: denotes whether a participant has any alpha-amylase data (1=yes, 2=no).

The raw alpha-amylase values for the four occasions are represented by *RA2DAA1W* (alpha-amylase at waking), *RA2DAA1A* (30 minutes after waking), *RA2DAA1L* (before lunch), and *RA2DAA1B* (before bed). No boundary restrictions are placed on the values.

In addition to actual raw values, the following values denote some type of non-response.

- 9995. NO SALIVA DATA
- 9998. MISSING
- 9999. INAPP

We also provide an adjusted value for total alpha-amylase for a minute of saliva flow. These variables are represented by *RA2DAA2W* (alpha-amylase at waking), *RA2DAA2A* (30 minutes after waking), *RA2DAA2L* (before lunch), and *RA2DAA2B* (before bed).