

A binomial count of meromorphic travelling waves

Alexander Migita

Abstract

We study meromorphic travelling waves of $u_t + uu_x + \sum_{j=1}^p b_j \partial_x^{j+1} u = 0$, a constant-coefficient family containing KdV–Burgers, Kuramoto–Sivashinsky and Kawahara. After integration and normalization, the profiles satisfy $P(D)v + v^2/2 = 0$, where $D = d/dz$ and P is monic of degree $p \geq 2$ with complex coefficients.

For evolution equations of even order (p odd), and for purely dispersive equations (only odd-order derivatives), we give a complete algebraic description of all nonconstant globally meromorphic travelling waves: they are rational, rational in an exponential, or elliptic, with one pole per period.

For $p \geq 3$ such waves are exceptional: a generic equation of the family has none. We count the exceptions with the periods of the wave fixed. Across the whole family there are $\binom{2p-1}{p-1}$ normalized one-pole pairs (P, v) with v rational in an exponential of given rate, and $2\binom{2p-2}{p-2}$ with v elliptic on a given generic lattice. We also count how many of these belong to the purely dispersive subfamily (p even) and to the purely dissipative one (only even-order derivatives in the linear part, p odd). Each contains $F_p = \binom{p-1}{\lfloor (p-1)/2 \rfloor}$ of the rational-exponential pairs: all solitary pulses in the former, all fronts in the latter. The purely dispersive subfamily also contains $2F_p$ of the elliptic pairs.

When $2p - 1$ is prime, these statements are proved and all pairs are distinct. Exact computer certificates establish the counts, with multiplicities, for $2 \leq p \leq 73$. We conjecture the same in every order.

Keywords. Meromorphic travelling waves; dispersive and dissipative equations; elliptic functions; polynomial convolution; algebraic multiplicity.

1 Introduction

Exact travelling waves of nonlinear evolution equations are usually found one equation at a time, and usually only for special values of the coefficients. The generalized Kuramoto–Sivashinsky equation is a classical example. It has nonconstant travelling waves that are polynomials in a hyperbolic tangent at exactly four values of its squared dimensionless dispersion parameter. Kuramoto and Tsuzuki found the dispersionless one [37], Kudryashov the other three [24, 25], and Eremenko proved that, together with Kudryashov’s elliptic solution and a single rational one, they exhaust the nonconstant meromorphic travelling waves [18]. Counted as we count below, the tanh list has ten members. This paper shows that ten is $\binom{5}{2}$: the classical list is the order-three case of a binomial enumeration, proved for infinitely many orders and conjectured for all.

We consider the evolution equations

$$u_t + uu_x + \sum_{j=1}^p b_j \partial_x^{j+1} u = 0, \quad b_p \neq 0,$$

which contain KdV–Burgers ($p = 2$), Kuramoto–Sivashinsky ($p = 3$) and Kawahara ($p = 4$), the sixth-order turbulence models of Kudryashov and Migita [30], Simbawa et al. [42], and the quadratic long-wave reductions of Kudryashov et al. [32]. A travelling wave $u = w(x - ct)$ satisfies, after one integration, $\sum_j b_j w^{(j)} - cw + w^2/2 + K = 0$. Shifting w by a constant solution and normalizing the leading coefficient gives

$$E_P(v) := P(D)v + \frac{1}{2}v^2 = 0, \quad P(\rho) = \rho^p + \sum_{j=0}^{p-1} a_j \rho^j, \quad p \geq 2, \quad (1.1)$$

with $D = d/dz$ and complex coefficients. The speed, the integration constant and the background are absorbed into a_0 (Section 2). Throughout, p is the order of (1.1), one less than the order of the evolution equation.

Two subfamilies play a special role. In the evolution equation, derivatives of odd order are dispersive terms and derivatives of even order are dissipative terms. We call an equation *purely dispersive* if it contains only odd-order derivatives, as the KdV and Kawahara equations do; then p is even and P is an even polynomial. We call it *purely dissipative* if its linear part contains only even-order derivatives, as for the Kuramoto–Sivashinsky equation without dispersion; then p is odd and $P - P(0)$ is an odd polynomial. All other equations, such as KdV–Burgers, are *mixed*. The two pure subfamilies are the equations that are invariant under a reflection of x , and for their travelling waves this reflection is $z \mapsto -z$.

Classification. Eremenko’s theorem states that an autonomous algebraic differential equation with a single monomial of top degree and finitely many Laurent expansions at a pole has only three kinds of nonconstant meromorphic solutions: rational, rational in an exponential, and elliptic [18]. Theorem 3.1 verifies these hypotheses for (1.1) in two cases. The first is p odd, that is, every evolution equation of even order: there the Laurent expansion at a pole is unique. The second is the purely dispersive case: there one Laurent coefficient is free, but (1.1) is invariant under $z \mapsto -z$ and conserves an energy, and fixing the energy fixes that coefficient. In both cases every nonconstant meromorphic travelling wave is of one of the three kinds and has one pole per period. A fixed equation has at most one such wave up to translation when p is odd, and at most one on each energy level when it is purely dispersive. The mixed equations of odd order are not covered, and cannot be: Section 10 recalls a second-order example with other meromorphic waves. Section 4 writes each of the three kinds as a finite polynomial system, following the principal-part construction of Demina and Kudryashov [15, 16].

The count. For $p \geq 3$ such waves are exceptional: a generic equation of the family has none (Section 8). We count the exceptions, with the periods of the wave fixed. For waves that are rational in an exponential, fix the rate to one, place the pole at $i\pi$, and require $v \rightarrow 0$ as $e^z \rightarrow 0$. Let P range over all monic polynomials of degree p and count the pairs (P, v) for which v is such a solution of (1.1). For elliptic waves, fix a nonsingular Weierstrass lattice and count the pairs (P, v) with v elliptic on that lattice and its only pole at zero. The answers are

$$\begin{aligned} N_p &= \binom{2p-1}{p-1} && \text{rational-exponential pairs,} \\ M_p &= 2 \binom{2p-2}{p-2} && \text{elliptic pairs on a generic lattice.} \end{aligned}$$

Whenever $2p - 1$ is prime, both formulas hold and every pair is simple, that is, nondegenerate, so that all pairs are distinct; in the elliptic case this is for a generic lattice (Theorems 5.4 and 6.1). For every $2 \leq p \leq 73$, the formulas hold as counts with multiplicity (Section 7). On special lattices elliptic pairs do merge, and Section 9.2 determines where this happens at order three. At $p = 5$, where $2p - 1 = 9$ is the first composite index, a separate exact computation shows that all 126 rational-exponential pairs are simple. We conjecture the same counting and simplicity statements in every order (Conjecture 10.1). The counts are over $\mathbb{C}$ and range over all monic P , mixed operators included. The classification is used only to know that, where it applies, there are no other meromorphic waves.

Purely dispersive and purely dissipative equations. The reflection $z \mapsto -z$ acts on the pairs, and the pairs that it fixes are exactly those whose equation is purely dispersive, for even p , or purely dissipative, for odd p (Theorem 5.8). Under the same hypotheses as for N_p , their number is

$$F_p = \binom{p-1}{\lfloor (p-1)/2 \rfloor},$$

and the purely dispersive equations also account for $2F_p$ of the elliptic pairs on a generic lattice (Corollary 6.2). The two subfamilies carry different kinds of rational-exponential waves. Those of a purely dispersive equation are all pulses, which return to the same constant at both ends. Those of a purely dissipative equation are all fronts, which connect two different constants; this is proved whenever all pairs are distinct, and for real waves in every order. For example, the Kawahara counts 3 and 6 are F_4 and $2F_4$; the seventh-order KdV family has ten pairs, two of them real; and the six pairs of the purely dissipative family of order five contain the four kink solutions of Kudryashov and Migita [30] (Section 9).

A single equation. The count is over operators, and Section 8 translates it back to one evolution equation. Rescaling x moves the coefficient vector of an equation along a weighted curve, and this is why the rate is fixed in the count. An equation has a one-pole travelling wave that is rational in an exponential if and only if its coefficient vector lies on one of at most $(N_p + F_p)/2$ such curves. Unless all the coefficients b_j with $j < p$ vanish, it then has finitely many such waves, up to translations and Galilean shifts (Corollary 8.2). For odd $5 \leq p \leq 73$, the equations with any nonconstant meromorphic travelling wave form a set of dimension at most two in the $(p-1)$ -dimensional space of coefficients, so a generic equation has none (Corollary 8.3). At $p = 3$ this set is the union of the four classical Kuramoto–Sivashinsky curves.

Method. Every rational-exponential pair has a profile of the form $v_A = s_p A(D)Q$, where $Q = 1/(1 + e^{-z})$, s_p is a fixed scale and A is a monic polynomial of degree $p-1$. The pair (P, v_A) solves (1.1) if and only if A divides its own discrete self-convolution $S_A(n) = \sum_{j=1}^{n-1} A(j)A(n-j)$, and then $P = s_p S_A/(2A)$ (Proposition 5.1). When $\ell = 2p - 1$ is prime, $\ell S_A \equiv \pm(\rho^\ell - \rho) \pmod{\ell}$, so modulo ℓ the condition becomes $A \mid \rho^\ell - \rho$, and its solutions are indexed by the $(p-1)$ -element subsets S of $\mathbb{F}_\ell$. Each of them lifts by Hensel's lemma to a simple solution, and a weighted Bézout bound shows that the lifts are all the solutions. The reflection acts on the subsets by $S \mapsto -S$, which gives F_p . For the elliptic pairs, the same Bézout argument gives the count with multiplicity on every nonsingular lattice, once a degeneration to the cusp excludes solutions at infinity. A degeneration to the node, where the elliptic problem becomes the pulse case of the exponential one with two choices of background, shows that the pairs are simple on a generic lattice.

Two ingredients resist a uniform proof. One is a rigidity statement for the continuous convolution $\int_0^p A(t)A(\rho - t) dt$, needed to exclude solutions at infinity; it is proved for prime and prime-power indices, certified by exact arithmetic through degree 72, and would give all counts with multiplicity in every order. The other is simplicity of every rational-exponential pair, proved for prime indices and, by exact computation, at $p = 5$, and open in general (Conjecture 10.1).

Relation to earlier work. The classification theorem is a uniform version of results that exist in pieces. For odd p the indicial polynomial has no positive integer zero, the Laurent expansion at a pole is unique, and Eremenko's theorem applies directly; this case is contained in Yuan et al. [45, Theorem 1], with related odd-order results in Huang et al. [21], Yuan et al. [46] and a fifth-order instance in Dang [10]. For even p the index $3p$ is resonant, and the conclusion need not hold for mixed operators: the Fisher equation has an elliptic-exponential family of solutions [1, 5, 28] whose generic members are meromorphic and of none of the three kinds. The familiar Ablowitz–Zeppetella front is a rational-exponential degeneration of this family. When P is an even polynomial the equation has a first integral, the energy, and fixing its value removes the resonant freedom. This is the device of Eremenko et al. [19, Lemmas 3–4] for $P = D^p + a_0$, stated as a general step by Conte et al. [6, 7] and applied to the Kawahara equation [14, 44], the generalized Bretherton equation [15, 36] and the seventh-order KdV equation [9]. Theorem 3.1 extends it uniformly to every even P , intermediate derivatives included. Equations with two dominant monomials, or without the finiteness property, are treated by other methods in Demina [12, 13]. The general enumeration has no precedent that we know of; its special cases at $p = 3$ and in the Kawahara family are classical, as the examples make explicit.

Organization. Section 2 fixes the normalizations and the vocabulary. Sections 3 and 4 contain the classification and the three polynomial systems. Sections 5 and 6 prove the two counts, Section 7 records which orders are settled, and Section 8 reads the count for a single equation. Section 9 works out the lowest orders and the two pure subfamilies, including the lattices on which elliptic pairs merge. Section 10 collects the open questions, and Appendix A describes the exact verification and the ancillary material. All counts are over $\mathbb{C}$; the examples identify the real profiles, and stability is not discussed.

2 Equations, waves and what is counted

This section fixes the normalizations and the vocabulary used in the rest of the paper. It contains no new result.

2.1 From the evolution equation to the profile equation

Let $u = w(\xi)$, $\xi = x - ct$, be a travelling wave of

$$u_t + uu_x + \sum_{j=1}^p b_j \partial_x^{j+1} u = 0, \quad b_p \neq 0. \quad (2.1)$$

One integration gives

$$\sum_{j=1}^p b_j w^{(j)} - cw + \frac{1}{2}w^2 + K = 0 \quad (2.2)$$

with a constant K . Let w_* be a constant solution of (2.2), that is, a root of $w^2/2 - cw + K$, and let $\lambda \neq 0$. The substitution

$$w(\xi) = w_* + b_p \lambda^p v(\lambda \xi) \quad (2.3)$$

turns (2.2) into the profile equation (1.1) with

$$a_j = \frac{b_j}{b_p} \lambda^{j-p} \quad (1 \leq j \leq p-1), \quad a_0 = \frac{w_* - c}{b_p \lambda^p}. \quad (2.4)$$

Thus $a_1, \dots, a_{p-1}$ record the equation, up to the scale λ , and a_0 records the speed of the wave relative to the background w_* . The scale is free at this stage. It will be fixed by the period of the wave.

The constant solutions of (1.1) are $v = 0$ and $v = -2a_0$; they are the two roots w_* . Choosing the other root replaces the pair (P, v) by

$$(P - 2P(0), v + 2P(0)), \quad (2.5)$$

which again satisfies (1.1), as a direct substitution shows. The two pairs describe the same wave of the same evolution equation.

2.2 Dispersive, dissipative and mixed equations

In (2.1) a derivative of odd order is a dispersive term, and a derivative of even order is a dissipative term: for real coefficients the first changes the phase of a Fourier mode and the second its amplitude. We use the following names.

- The equation is *purely dispersive* if it contains only odd-order derivatives, that is, $b_j = 0$ for odd j . Then p is even, P is an even polynomial, and the equation is invariant under $(x, t) \mapsto (-x, -t)$. Examples are the KdV equation ($p = 2$), the Kawahara equation ($p = 4$) and the seventh-order KdV equation ($p = 6$).
- The equation is *purely dissipative* if its linear part contains only even-order derivatives, that is, $b_j = 0$ for even j . Then p is odd, $P - P(0)$ is an odd polynomial, and the equation is invariant under $(x, u) \mapsto (-x, -u)$. Examples are the Kuramoto–Sivashinsky equation without dispersion ($p = 3$) and the Nikolaevskiy equation ($p = 5$); see Simbawa et al. [42].
- All other equations are *mixed*. Examples are the KdV–Burgers equation and the Kuramoto–Sivashinsky equation with dispersion.

The two pure subfamilies are described by one condition. We call a monic polynomial P of degree p *reflection-symmetric* if $P - P(0)$ has the parity of ρ^p :

$$P(-\rho) = P(\rho) \quad (p \text{ even}), \quad P(\rho) + P(-\rho) = 2P(0) \quad (p \text{ odd}). \quad (2.6)$$

In the travelling-wave variable both symmetries are the reflection $z \mapsto -z$. If P is even and $v(z)$ solves (1.1), so does $v(-z)$. If p is odd and $P - P(0)$ is odd, so does $-v(-z) - 2P(0)$; this is the reflection combined with the change of background (2.5).

2.3 Waves, poles, pulses and fronts

Throughout, a meromorphic function is single-valued and meromorphic on the whole complex plane. Three kinds of meromorphic travelling waves occur in this paper: rational in z , rational in an exponential $e^{\kappa z}$, and elliptic. For the equations covered by Theorem 3.1 there are no others; for the remaining equations these are the waves that we count. The period group of such a wave is its *full* group of periods: trivial in the first case, infinite cyclic in the second, a lattice in the third. We say that the wave has *one pole per period* if its poles form a single coset of this group. There is then one pole in the rational case, one pole modulo the primitive period in the second case, and one pole in each period parallelogram in the elliptic case.

A wave that is rational in $e^{\kappa z}$ and has one pole per period has finite limits as $e^{\kappa z} \rightarrow 0$ and ∞ (Section 4). Both limits are constant solutions. The wave is a *pulse* if the two limits coincide and a *front* otherwise. For a real rate and a real profile these are the solitary waves and the kinks of the evolution equation, and they are smooth and bounded on the real axis when the pole lies off it. In (2.3) we take w_* to be the limit as $e^{\kappa z} \rightarrow 0$. Then $v \rightarrow 0$ at that end, and the other limit is 0 for a pulse and $-2a_0$ for a front.

2.4 What is counted

We count pairs (P, v) : a monic polynomial P of degree p and a solution v of (1.1) of a given kind, with its periods fixed.

- A *rational-exponential pair* has v rational in e^z , with one pole modulo $2\pi i\mathbb{Z}$, placed at $i\pi$, and $v \rightarrow 0$ as $e^z \rightarrow 0$.
- An *elliptic pair* on a lattice Λ has v elliptic with period lattice Λ , including its scale, and its only pole modulo Λ at zero.

Fixing the period or the period lattice removes the scale λ in (2.3), and placing the pole removes translations. The operator P varies over *all* monic polynomials of degree p , and all counts are over $\mathbb{C}$. A wave and its mirror image are counted as two pairs when they are distinct; the pairs that coincide with their mirror image are the subject of Section 5.4. In the elliptic case both pairs (2.5) are counted; in the rational-exponential case the condition $v \rightarrow 0$ selects one of them. Section 8 translates the counts back to a single evolution equation.

3 The classification theorem

This section proves Theorem 3.1 below: for every evolution equation of even order (p odd), and for every purely dispersive equation (P even), a nonconstant meromorphic solution of (1.1) is rational, rational in an exponential, or elliptic, and has one pole per period. The proof has three steps. We first compute the Laurent expansion at a pole and show that for odd p it is unique. For even P it is not: one coefficient is left free, and we show that the energy, a first integral of (1.1), fixes it, so that Eremenko's theorem applies on each energy level. Finally we show that any two poles of one solution differ by a period.

Let $\mathcal{W}$ consist of rational functions of z , rational functions of $e^{\kappa z}$ with $\kappa \neq 0$, and elliptic functions. Period groups are full period groups, as in Section 2; for a rational function of $e^{\kappa z}$ we choose κ so that $2\pi i/\kappa$ generates the group. For the general analytic background, see Eremenko [17].

The theorem collects the analytic input to the enumeration. Its odd-order case is contained in Yuan et al. [45, Theorem 1]. For the even-order case, the first-integral mechanism is that of Eremenko et al. [19, Lemmas 3–4]; here it is written uniformly for every even polynomial P . We give the proofs in full because Sections 4 and 5 use the Laurent recurrence and the energy dependence in explicit form.

Theorem 3.1 (Classification). *Assume either that p is odd or that P is an even polynomial; that is, the evolution equation has even order or is purely dispersive. Every nonconstant meromorphic solution of (1.1) belongs to $\mathcal{W}$. Every pole has order p , and the pole set is one coset of the full period group: the solution has one pole per period. Consequently there is one pole in the rational case, one pole modulo the primitive period in the simply periodic case, and one pole on the full elliptic period torus.*

For fixed P of odd degree, there is at most one nonconstant solution with its pole at zero. For even P , the same statement holds after fixing the conserved energy. The only entire solutions are $v = 0$ and $v = -2a_0$.

An even polynomial is one in which every odd power of ρ has coefficient zero, so that $P(D)$ contains only even derivatives. The hypothesis does not cover the mixed equations of odd order, and it cannot: Section 10 gives a mixed second-order operator with a meromorphic solution outside $\mathcal{W}$.

3.1 Pole order and the Laurent recurrence

Write a pole expansion at zero as

$$v(z) = \sum_{n \geq 0} c_n z^{n-p}. \quad (3.1)$$

The pole order is forced by dominant balance. A pole of order r makes $P(D)v$ of order $r + p$ and v^2 of order $2r$; the two can cancel only if $r = p$. Comparing the leading coefficients gives

$$c_0 = c_* := -2(-1)^p R_p, \quad R_p = \frac{(2p-1)!}{(p-1)!}. \quad (3.2)$$

For the remaining coefficients, set

$$I_p(n) = \prod_{j=p}^{2p-1} (n-j) + c_*. \quad (3.3)$$

The product is the factor produced by D^p acting on z^{n-p} , and c_* is the factor produced by the cross term of $c_n z^{n-p}$ with the leading pole in $v^2/2$. With $c_n = 0$ for $n < 0$, comparison of the coefficients of z^{n-2p} in (1.1) gives

$$I_p(n)c_n = -\sum_{j=0}^{p-1} a_j (n-2p+j)^{\underline{j}} c_{n-p+j} - \frac{1}{2} \sum_{k=1}^{n-1} c_k c_{n-k}, \quad n \geq 1. \quad (3.4)$$

The underline denotes a falling factorial. The right side contains only coefficients with index less than n , so (3.4) determines c_n whenever $I_p(n) \neq 0$. The positive integer zeros of I_p are the *resonances* of the recurrence: at such an index the recurrence leaves c_n undetermined if its right side vanishes, and has no solution otherwise.

If p is odd, then $c_* = 2R_p$, and there is no positive integer resonance. For integers $1 \leq n < p$, the product in (3.3) is negative with absolute value less than R_p . For $p \leq n \leq 2p - 1$ it is zero, and for $n \geq 2p$ it is positive. Hence $I_p(n) \neq 0$ for every positive integer n , and there is exactly one formal Laurent expansion at the marked pole.

Theorem 2 of Eremenko [18] states that an autonomous polynomial differential equation with finitely many possible formal pole expansions and a single monomial of highest total degree has all its meromorphic solutions in $\mathcal{W}$. In (1.1) the only monomial of degree two is $v^2/2$, and for odd p we have just seen that there is exactly one formal pole expansion. Both hypotheses hold at every odd order, for every coefficient vector a .

3.2 The energy argument for an even operator

Let $p = 2m$ and

$$P(D) = D^{2m} + \sum_{j=0}^{m-1} b_j D^{2j}. \quad (3.5)$$

Now $c_* = -2R_p$, and there is exactly one positive integer resonance. Indeed the product in (3.3) is at most R_p for $0 \leq n < p$, is zero for $p \leq n \leq 2p - 1$, and is strictly increasing for $n \geq 2p$; its value at $3p$ is $2R_p = -c_*$. The only positive integer zero of I_p is therefore $n = 3p$, and the possible free Laurent coefficient is

$$h = c_{3p} = [z^{2p}]v.$$

If the recurrence is compatible at $n = 3p$, the equation has a one-parameter family of formal expansions at the pole, one for each value of h , and Eremenko's theorem does not apply to (1.1) itself. The remedy is a first integral whose value determines h .

For $j \geq 1$, define the quadratic differential expression

$$\mathcal{B}_j(v) = \sum_{\ell=0}^{j-2} (-1)^\ell v^{(\ell+1)} v^{(2j-1-\ell)} + \frac{(-1)^{j-1}}{2} (v^{(j)})^2, \quad (3.6)$$

where the sum is empty when $j = 1$. Termwise differentiation gives $D\mathcal{B}_j = v'v^{(2j)}$: every even derivative, multiplied by v' , is an exact derivative. Since P contains only even derivatives, it follows that

$$\mathcal{H}_P(v) = \mathcal{B}_m(v) + \sum_{j=1}^{m-1} b_j \mathcal{B}_j(v) + \frac{b_0}{2} v^2 + \frac{1}{6} v^3 \quad (3.7)$$

satisfies

$$D\mathcal{H}_P = v'E_P(v). \quad (3.8)$$

In particular $\mathcal{H}_P$, the energy, is constant on each solution. It is the energy in the usual sense: an even operator $P(D)$ is formally self-adjoint, (1.1) is then the Euler–Lagrange equation of an autonomous Lagrangian, and $\mathcal{H}_P$ is the conserved quantity that belongs to translation in z . The purely dispersive equations are also exactly those for which (1.1) is invariant under $z \mapsto -z$. The case with no intermediate derivatives is the first integral used by Eremenko et al. [19]. The familiar Kawahara first integral is the case $m = 2$ [14, 44]; the corresponding order-six construction appears in Dang [9].

The energy identity has two consequences for the Laurent expansion. The first is that the resonance is always compatible. To see this, construct (3.1) through index $3p - 1$. The first possible residual of (1.1) is $\Omega_p z^p$. Multiplication by v' gives residue $-pc_*\Omega_p$. By (3.8) this is the

residue of a derivative of a Laurent series, and is therefore zero. Hence $\Omega_p = 0$, and h may be chosen arbitrarily.

The second consequence is that the energy determines h . On the full formal solution the constant energy has the form

$$\mathcal{H}_P = \Lambda_p h + \eta_p(P), \quad \Lambda_p = -pc_* I'_p(3p) \neq 0. \quad (3.9)$$

To find the coefficient, linearize the highest-weight part of (3.8) about $c_* z^{-p}$, with perturbation z^{n-p} . If the corresponding energy term is $L(n)z^{n-3p}$, then

$$(n - 3p)L(n) = -pc_* I_p(n).$$

Taking $n \rightarrow 3p$ gives the coefficient in (3.9). Lower derivative terms have different powers of z and cannot contribute to this coefficient. Moreover,

$$I'_p(3p) = 2R_p \sum_{j=p+1}^{2p} \frac{1}{j} > 0.$$

Fixing the energy thus fixes h , and all subsequent Laurent coefficients follow from (3.4). This is the resonance-removal step in Eremenko et al. [19, Lemma 4], with an explicit coefficient for the present operator family; see also Conte et al. [6, Section 3.2] and Conte et al. [7, Section 5].

We can now apply Eremenko's theorem, not to (1.1) but to the level equation $\mathcal{H}_P(v) = H$ with P and H fixed. This equation is autonomous and polynomial, and its unique monomial of highest degree is $v^3/6$. It has exactly one formal pole expansion: differentiating it and dividing by the nonzero Laurent series v' returns $E_P(v) = 0$, so every expansion belongs to the family above, and (3.9) selects the value of h . Every meromorphic solution of (1.1) lies on one energy level and therefore belongs to $\mathcal{W}$.

3.3 The period group and the entire solutions

It remains to show that the poles of one solution form a single orbit under its periods. For either covered case, take two poles z_1, z_2 of the same solution. The functions $v(z + z_1)$ and $v(z + z_2)$ satisfy the same equation and, in the even case, have the same energy. Their Laurent series at zero coincide, so the identity theorem gives

$$v(z + z_1) = v(z + z_2).$$

Every difference of poles is a period. Conversely, every period translates a pole to a pole. The period group is discrete: a sequence of nonzero periods tending to zero would force $v' \equiv 0$. The pole set is therefore one coset of that group. The same argument proves the uniqueness statements of the theorem: two nonconstant solutions with the same operator, the same energy in the even case, and a pole at zero have the same Laurent series there.

Finally, an entire solution in $\mathcal{W}$ must be constant. In the rational case it is a polynomial, and the highest degree of its square cannot be cancelled in (1.1) unless the polynomial is constant. In the rational-exponential case it is a Laurent polynomial in $t = e^{\kappa z}$; the operator $P(\kappa t \partial_t)$ preserves each exponent, whereas squaring doubles an extreme nonzero exponent. An entire elliptic function is constant by compactness. The constants solve $a_0 v + v^2/2 = 0$, which gives $v = 0$ and $v = -2a_0$. This completes the proof of Theorem 3.1.

The same uniqueness shows that, in the two pure subfamilies of Section 2, every wave inherits the reflection symmetry of its equation.

Corollary 3.2 (Waves are symmetric about their poles). *Let v be a nonconstant meromorphic solution of (1.1) and let z_0 be one of its poles.*

- (i) *If P is even, then $v(z_0 + \zeta) = v(z_0 - \zeta)$.*
- (ii) *If p is odd and $P - P(0)$ is odd, then $v(z_0 + \zeta) + P(0) = -(v(z_0 - \zeta) + P(0))$.*

Consequently a solution that is rational in an exponential is a pulse in case (i). In case (ii) its two limits add up to $-2P(0)$, so it is a front unless $P(0) = 0$.

Proof. In case (i) the function $v(z_0 - \zeta)$ solves the same equation, has a pole at $\zeta = 0$, and has the same energy as v , because every term of (3.6) and (3.7) contains an even total number of derivatives. Fixed-energy uniqueness gives (i). In case (ii) the function $-v(z_0 - \zeta) - 2P(0)$ solves the same equation, by Section 2, and has a pole at $\zeta = 0$. For odd p the Laurent expansion at a pole is unique, which gives (ii). The statements about the limits follow by letting ζ tend to the two ends. $\square$

The finite systems of the next section are derived from the one-pole structure alone. The analytic facts of this section are used only to show that, in the covered cases, those systems miss no nonconstant meromorphic solution. The systems themselves never require the convergence of a formal series.

4 Three finite algebraic systems

Theorem 3.1 says that, in the covered cases, a nonconstant meromorphic solution is one of three kinds and has one pole per period. Each kind is a finite-dimensional family of functions: a polynomial in $1/z$, a polynomial in a fixed simply periodic function with one pole, or a constant plus a linear combination of $\wp$ and its derivatives. Substituting the general member of each family into (1.1) therefore gives finitely many polynomial equations in its coefficients. This section writes down the three systems, fixes their normalizations, and proves that they are exactly equivalent to the one-pole solutions (Theorem 4.1). They are instances of the principal-part construction of Demina and Kudryashov [15, 16].

The systems all have the same shape. Let V be a polynomial expression in some variables, and let δ be a differentiation rule for those variables, so that the polynomials in them form a differential algebra with derivation δ . Write

$$\mathcal{E}_P(V; \delta) = \delta^p V + \sum_{j=0}^{p-1} a_j \delta^j V + \frac{1}{2} V^2.$$

Each system below is the polynomial identity $\mathcal{E}_P(V; \delta) = 0$ for an appropriate V and δ . A polynomial identity means the vanishing of every coefficient.

4.1 Rational solutions

A rational solution whose only pole is at zero, of order p , and which has no nonconstant polynomial part at infinity is a polynomial of degree p in $T = 1/z$. Since $DT = -T^2$, the equation becomes

$$V_R(T) = b + \sum_{k=1}^p r_k T^k, \quad r_p = c_*, \quad \mathcal{E}_P(V_R; -T^2 \partial_T) = 0. \quad (4.1)$$

The left side of the identity is a polynomial of degree at most $2p$ in T , so this is a finite system in $b, r_1, \dots, r_{p-1}$. No rational solution has a nonconstant polynomial part at infinity: such a part would have its degree doubled by v^2 , while the linear differential expression cannot reach that degree. Consequently (4.1) is the full rational one-pole system.

4.2 Rational functions of an exponential

For $q = \kappa^2 \neq 0$, define

$$T_q(z) = \frac{\kappa}{2} \coth \frac{\kappa z}{2}, \quad DT_q = \frac{q}{4} - T_q^2. \quad (4.2)$$

The expression is independent of the choice of sign of κ . It has period $2\pi i/\kappa$ and one pole per period, at zero, with residue one; it plays the role that $1/z$ plays in the rational case. The finite system is

$$V_S(T) = \sum_{k=0}^p u_k T^k, \quad u_p = c_*, \quad q \neq 0, \quad \mathcal{E}_P \left(V_S; \left(\frac{q}{4} - T^2 \right) \partial_T \right) = 0. \quad (4.3)$$

To see that every one-pole solution rational in an exponential has this form, write $v = R(t)$, $t = e^{\kappa z}$. The rational function R has finite limits at both $t = 0$ and $t = \infty$. If it had a pole at either end, an extreme Laurent monomial would be doubled by R^2 and could not cancel against $P(\kappa t \partial_t)R$. After a translation, its only finite pole is at $t = 1$, of order p . Such a function is a degree- p polynomial in T_q . The logistic coordinate

$$Q = \frac{1}{1 + e^{-z}}, \quad DQ = Q(1 - Q),$$

is an equivalent coordinate after fixing the rate and translating the pole; $Q(z + i\pi) = T_1(z) + 1/2$. Polynomial logistic representations and their use in constructing travelling waves are developed by Kudryashov [29, 33, 34].

The rational system is the $q = 0$ specialization of (4.3), with $T_0 = 1/z$. We keep $q = 0$ and $q \neq 0$ distinct so that this degeneration is never counted twice.

4.3 Elliptic solutions

Use the Weierstrass algebra

$$Y^2 = 4X^3 - g_2X - g_3, \quad \mathcal{D}X = Y, \quad \mathcal{D}Y = 6X^2 - \frac{g_2}{2}, \quad \Delta = g_2^3 - 27g_3^2 \neq 0. \quad (4.4)$$

Its analytic realization is $X = \wp(z; g_2, g_3)$, $Y = \wp'(z; g_2, g_3)$; see NIST [41, Chapter 23]. Set

$$V_E = b + \sum_{j=0}^{p-2} t_j \mathcal{D}^j X, \quad t_{p-2} = -s_p, \quad s_p = \frac{2(2p-1)!}{((p-1)!)^2}, \quad \mathcal{E}_P(V_E; \mathcal{D}) = 0. \quad (4.5)$$

Using (4.4), every polynomial in X and Y reduces to the normal form $F(X) + YG(X)$. The system consists of the coefficients of F and G set equal to zero; all of them are polynomials with rational coefficients in the displayed unknowns.

The representation is the classical one. An elliptic function with one pole has zero residue there. The functions $1, \wp, \wp', \dots, \wp^{(p-2)}$ have distinct principal parts and span the space of elliptic functions whose only pole is at zero, of order at most p . Subtracting the indicated principal part

from a one-pole solution leaves an entire elliptic function, hence a constant. The leading coefficient in (4.5) follows from

$$D^{p-2}\wp = (-1)^{p-2}(p-1)!z^{-p} + O(1).$$

The nonsingularity condition $\Delta \neq 0$ is retained throughout; the singular cubics correspond to the rational and simply periodic systems, which we have already written down.

Theorem 4.1 (Finite matching and exhaustion). *For every $p \geq 2$, the systems (4.1), (4.3), and (4.5) are necessary and sufficient for the respective one-pole solutions of (1.1), with the pole at zero and with the stated normalizations. For the operator classes of Theorem 3.1, these systems, together with the constant solutions, exhaust all globally meromorphic solutions.*

The systems describe exceptional parameter values and continuous families as well as isolated solutions. Their sufficiency follows from substitution, without any genericity assumption.

Proof. Necessity is the principal-part argument given for each kind above. Conversely, a solution of a coefficient system gives a globally meromorphic function by its displayed realization, and the chain rule gives $E_P(v) = 0$. The leading coefficient $c_* \neq 0$ ensures a nonconstant function. In the elliptic case every nonsingular pair (g_2, g_3) has a Weierstrass realization, and the normal form $F(X) + YG(X)$ is unique. The exhaustion statement is Theorem 3.1. $\square$

4.4 Fixed operators

The systems above treat the operator and the profile coefficients as unknowns together, which is the form that the count requires. For a fixed operator they reduce further. The Laurent recurrence (3.4) determines the principal part of any solution from P , and the remaining unknowns are the rate q or the invariants g_2, g_3 . This is the coefficient-comparison construction of Demina and Kudryashov [11, 15]; for even P one appends the value of the free coefficient h prescribed by the energy through (3.9). We do not need this reduction below.

5 The count of rational-exponential pairs

In this section and the next, P ranges over *all* monic polynomials of degree p , mixed operators included, and we count the pairs (P, v) of Section 2. Nothing in the count requires the hypotheses of Theorem 3.1. Those hypotheses were needed to show that the one-pole solutions are all the meromorphic solutions; the matching equations and their solution counts make sense without them. The pairs that belong to the purely dispersive and purely dissipative subfamilies are counted in Section 5.4.

Recall the normalization: the rate is one, the pole is at $i\pi$ modulo $2\pi i\mathbb{Z}$, and $v \rightarrow 0$ as $e^z \rightarrow 0$. A wave and its mirror image are counted separately when they differ. The functions $Q, DQ, \dots, D^{p-1}Q$ span the degree- p polynomials in Q with zero constant term, since

$$D^j Q = (-1)^j j! Q^{j+1} + \text{lower powers of } Q.$$

The leading pole coefficient (3.2) therefore gives the unique form

$$v_A = s_p A(D)Q, \quad A(\rho) = \rho^d + \alpha_1 \rho^{d-1} + \dots + \alpha_d, \quad d = p - 1. \quad (5.1)$$

The scale s_p was defined in (4.5). This representation is the standard singular-part operator form: $Q = 1/2 + D \log \cosh(z/2)$, so (5.1) is a constant plus a linear differential operator applied to

$\log \cosh(z/2)$. Such forms, including the KS case, are discussed by Conte and Musette [3]. We use the coefficients $\alpha_1, \dots, \alpha_d$ of that operator as the coordinates for the enumeration.

The argument runs as follows. The profile v_A solves (1.1) for some monic P exactly when A divides a polynomial built from A by a discrete convolution (Proposition 5.1). The remainder of that division gives d polynomial equations in the d coordinates. We count their solutions with multiplicity by Bézout's theorem, using a weighted degree adapted to the equations (Theorem 5.3). When $2p - 1$ is prime, reduction modulo $2p - 1$ identifies the solutions one by one (Theorem 5.4). The reflection $z \mapsto -z$ then singles out the pairs of the two pure subfamilies (Theorem 5.8).

5.1 Discrete convolution and the matching equations

For polynomials F, G , define $F \star G$ by polynomial continuation of

$$(F \star G)(n) = \sum_{j=1}^{n-1} F(j)G(n-j), \quad n \geq 1.$$

The continuation exists by the power-sum formulas: for fixed F, G the right side is a polynomial in n . Write

$$S_A = A \star A.$$

For monic A of degree d ,

$$\deg S_A = 2d + 1, \quad [\rho^{2d+1}]S_A = \beta_d := \frac{(d!)^2}{(2d+1)!}. \quad (5.2)$$

The leading coefficient is the beta integral $\int_0^1 t^d(1-t)^d dt$.

Proposition 5.1 (Polynomial divisibility). *The profile (5.1) solves (1.1) for some monic P if and only if*

$$A \mid S_A. \quad (5.3)$$

The operator is unique and is

$$P_A = \frac{s_p S_A}{2A}. \quad (5.4)$$

Proof. In the half-plane $\operatorname{Re} z < 0$, the logistic function is a geometric series in e^z , and applying $A(D)$ multiplies the n th term by $A(n)$:

$$A(D)Q = \sum_{n \geq 1} (-1)^{n-1} A(n) e^{nz}.$$

Squaring gives the rational-function identity

$$(A(D)Q)^2 = -S_A(D)Q. \quad (5.5)$$

The map $B \mapsto B(D)Q$ is injective: its vanishing would give $B(n) = 0$ for every positive integer n . Consequently (1.1) is equivalent to $2P_A A = s_p S_A$, which has a polynomial solution P_A exactly when A divides S_A . Equation (5.2) and $s_p = 2/\beta_d$ show that P_A is monic of degree p . $\square$

The operator determines the profile. This holds for every p , without the hypotheses of Theorem 3.1, because it only uses the leading coefficient of a convolution.

Proposition 5.2 (One profile per operator). *Let A, B be monic polynomials of degree $p - 1$ satisfying $A \mid S_A$ and $B \mid S_B$. If $P_A = P_B$, then $A = B$. Consequently a monic operator of degree p admits at most one normalized rational-exponential one-pole profile. Any count of distinct pairs with this normalization is therefore also a count of distinct operators.*

Proof. Suppose $P_A = P_B = P$, and put $h = A - B$. If $h \neq 0$, its degree e is less than $d = p - 1$; write $f \neq 0$ for its leading coefficient. By (5.4) and the bilinearity and symmetry of the convolution,

$$S_A - S_B = h \star (A + B) = \frac{2}{s_p} P h.$$

The power-sum calculation in (5.2) gives the leading coefficient $2f \int_0^1 t^e (1-t)^d dt$ on the left. Since P is monic, the right side has leading coefficient $(2/s_p)f = \beta_d f$. Cancelling f would therefore give

$$2 \int_0^1 t^e (1-t)^d dt = \int_0^1 t^d (1-t)^d dt.$$

This is impossible: $e < d$ implies $t^e > t^d$ for $0 < t < 1$, so the left side exceeds twice the right side. Thus $h = 0$. $\square$

For odd p , Theorem 3.1 gives uniqueness among all nonconstant meromorphic solutions up to translation; for even P , it does so on a fixed energy level. Proposition 5.2 fixes the rational-exponential normalization and holds for every monic operator.

Divide S_A by the monic polynomial A :

$$S_A = AC_A + R_A, \quad R_A = \sum_{r=0}^{d-1} R_r(\alpha_1, \dots, \alpha_d) \rho^r. \quad (5.6)$$

The matching equations are $R_0 = \dots = R_{d-1} = 0$: d polynomial equations in the d unknowns $\alpha_1, \dots, \alpha_d$. Their solutions are in bijection with the normalized pairs (P_A, v_A) . We write $\mathcal{X}_p$ for the solution set together with the equations that define it; in algebraic language, $\mathcal{X}_p$ is the affine scheme defined by the R_r .

Three notions from algebra are used to count. A solution is *simple* if the Jacobian matrix of $(R_0, \dots, R_{d-1})$ with respect to $(\alpha_1, \dots, \alpha_d)$ is invertible there; equivalently, the equations admit no nonzero infinitesimal deformation of the solution. Every isolated solution has a *multiplicity*, the dimension over $\mathbb{C}$ of the local quotient algebra of the equations at that point. For a square system such as this one it may be pictured as the number of simple solutions into which the point splits under a generic small perturbation of the equations; it equals one exactly for simple solutions. When $\mathcal{X}_p$ is finite, its *length* is the sum of the multiplicities of its points, and $\mathcal{X}_p$ is *reduced* when every point is simple, so that the length is the number of points. An isolated solution of multiplicity greater than one is still the only solution near itself; local uniqueness is therefore weaker than simplicity.

5.2 Weighted degree and the role of continuous convolution

Bézout's theorem bounds the sum of the multiplicities of the isolated solutions of d polynomial equations in d unknowns by the product of their degrees. Equality holds when the system has no solutions at infinity, that is, when the highest-degree parts of the equations have no common zero other than the origin. The ordinary degrees of the matching equations give a poor bound. The

right measure of their size is a weighted degree that reflects how the α_j scale. Give α_j weight j , and ρ weight one. The remainder equations have highest weighted degrees

$$2d + 1 - r \quad (0 \leq r < d), \quad \text{namely } d + 2, \dots, 2d + 1. \quad (5.7)$$

Their highest weighted parts are the coefficients of $\text{rem}_A H_A$, where

$$H_A(\rho) = \int_0^\rho A(t)A(\rho - t) dt. \quad (5.8)$$

Indeed, if $A(\rho) = \sum a_i \rho^i$, then

$$H_A(\rho) = \sum_{i,j=0}^d \frac{i!j!}{(i+j+1)!} a_i a_j \rho^{i+j+1}.$$

These are precisely the highest weighted terms of the power sums defining S_A , and monic division preserves the highest weighted part. Indeed, A is weighted-homogeneous of weight d , and each step of long division subtracts a multiple of A of the same weight as the term being removed. In words, the discrete convolution S_A is a lower-order perturbation of the continuous convolution H_A , and the behaviour of the matching equations at infinity is governed by the continuous problem $A \mid H_A$.

For this degree, denote by (R_d) the condition

$$A \text{ monic}, \quad \deg A = d, \quad A \mid H_A \quad \implies \quad A(\rho) = \rho^d. \quad (5.9)$$

It says that the continuous problem has only the trivial solution. By the preceding paragraph, it is exactly the condition that the matching equations have no solution at weighted infinity.

Theorem 5.3 (Count with algebraic multiplicities). *For every $p \geq 2$, the sum of multiplicities of the isolated points of $\mathcal{X}_p$ is at most*

$$N_p = \binom{2p-1}{p-1}.$$

If (R_{p-1}) holds, then $\mathcal{X}_p$ is finite and

$$\text{length } \mathcal{X}_p = N_p. \quad (5.10)$$

This equality does not require simplicity.

Proof. Weighted degree is turned into ordinary degree by the finite substitution

$$\alpha_j = y_j^j, \quad 1 \leq j \leq d. \quad (5.11)$$

A monomial of weight w in the α_j becomes a monomial of ordinary degree w in the y_j , so the pulled-back equations have ordinary degrees at most $d + 2, \dots, 2d + 1$, and their highest-degree parts are the pull-backs of the highest weighted parts. Bézout's inequality for isolated solutions bounds the sum of their multiplicities by $(d + 2) \cdots (2d + 1)$; see Mondal [39, Chapter VIII]. If (R_d) holds, the leading homogeneous forms have no nonzero common zero. The homogenized system has no point at infinity, and ordinary Bézout gives equality [8].

It remains to return from the y_j to the α_j . A solution in the α_j with all $\alpha_j \neq 0$ has $1 \cdot 2 \cdots d = d!$ preimages in the y_j , one for each choice of the roots $y_j = \alpha_j^{1/j}$, each with the original multiplicity. In general, for each solution the sum of the multiplicities of its preimages is $d!$ times its original multiplicity, including at branch points where some α_j vanish. Thus the substitution multiplies the total length by $d!$. Algebraically, $\mathbb{C}[y_1, \dots, y_d]$ is a free module over $\mathbb{C}[\alpha_1, \dots, \alpha_d]$, with basis

$$y_1^{e_1} \cdots y_d^{e_d}, \quad 0 \leq e_j < j.$$

This remains true after imposing the equations and restricting to an isolated component. Dividing the Bézout count by $d!$ gives

$$\frac{(d+2) \cdots (2d+1)}{d!} = \binom{2d+1}{d}.$$

The no-infinity condition also gives finiteness of the whole affine scheme, completing the proof. $\square$

For $p = 3$, the variable weights are 1, 2 and the equation weights are 4, 5, so the calculation is $(4 \cdot 5)/(1 \cdot 2) = 10$.

5.3 The prime-index theorem

Theorem 5.4 (Prime-index enumeration). *If $\ell = 2p - 1$ is prime, then*

$$\#\mathcal{X}_p(\mathbb{C}) = \text{length } \mathcal{X}_p = \binom{\ell}{p-1},$$

and every point is simple. After a choice of an ℓ -adic embedding of the algebraic coefficients, reduction labels the points by the $(p-1)$ -element subsets $S \subset \mathbb{F}_\ell$:

$$\overline{A}(\rho) = \prod_{s \in S} (\rho - s). \quad (5.12)$$

Proof. Put $d = p - 1$, so $\ell = 2d + 1$, and work over $\mathbb{Z}_{(\ell)}$, the rationals integral at ℓ . The proof has four steps: a congruence for S_A modulo ℓ ; the solution of the reduced equations over $\mathbb{F}_\ell$; the exclusion of solutions at infinity, which proves (R_d) ; and the lifting of the reduced solutions back to characteristic zero.

Step 1: the congruence. Induction using $(D - n)Q^n = -nQ^{n+1}$ gives

$$Q^n = \frac{(-1)^{n-1}}{(n-1)!} \prod_{j=1}^{n-1} (D - j)Q.$$

Write $(A(D)Q)^2 = \sum_{n=2}^{2p} b_n Q^n$. Each b_n is an integer polynomial in the coefficients of A , and $b_{2p} = (d!)^2$. By (5.5),

$$S_A(\rho) = \sum_{n=2}^{2p} \frac{(-1)^n b_n}{(n-1)!} \prod_{j=1}^{n-1} (\rho - j).$$

Only the last denominator, $(2p-1)! = \ell!$, contains ℓ . Consequently ℓS_A is integral and

$$\ell S_A(\rho) \equiv \frac{(d!)^2}{(\ell-1)!} \prod_{j=1}^{\ell} (\rho - j) \equiv (-1)^d (\rho^\ell - \rho) \pmod{\ell}. \quad (5.13)$$

The scalar follows by pairing j with $\ell - j$ in $(\ell - 1)!$, and the product is $\rho^\ell - \rho$ because it runs over all residues modulo ℓ .

Step 2: the reduced equations. The scaled remainder equations $\ell R_r = 0$ therefore reduce exactly to $\overline{A} \mid \rho^\ell - \rho$. There are $\binom{\ell}{d}$ monic divisors of degree d , all defined over $\mathbb{F}_\ell$, and these exhaust the solutions over its algebraic closure. Each of them is a simple solution of the reduced equations. Write $\rho^\ell - \rho = \overline{A}C$. For a variation h with $\deg h < d$, the derivative of the remainder map is, up to the nonzero scalar $(-1)^d$,

$$h \longmapsto -\text{rem}_{\overline{A}}(Ch). \quad (5.14)$$

Since $\rho^\ell - \rho$ is squarefree, $\overline{A}$ and C are coprime, so multiplication by C is invertible modulo $\overline{A}$, and the map (5.14) is invertible.

Step 3: no solutions at infinity. We must also prove that characteristic-zero solutions cannot escape to infinity, that is, that (R_d) holds. Taking the highest weighted part of (5.13) gives

$$\ell H_A(\rho) \equiv (-1)^d \rho^\ell \pmod{\ell}. \quad (5.15)$$

Thus the reduced leading remainder equations have only $\overline{A} = \rho^d$. After (5.11), they are homogeneous equations with no nonzero common zero over $\overline{\mathbb{F}}_\ell$. The same is true in characteristic zero. If the equations had a complex projective zero, they would also have one with algebraic coordinates, since their coefficients are rational. Normalization at an ℓ -adic place would then produce a nonzero reduced point. The valuation argument is stated precisely in Appendix A. We have proved (R_d) , so Theorem 5.3 gives total length $\binom{\ell}{d}$.

Step 4: lifting. The multivariable simple-root Hensel lemma lifts each of the $\binom{\ell}{d}$ reduced points to a unique point over $\mathbb{Z}_\ell$; see Conrad [2]. The lifts are distinct and simple. Their number already equals the characteristic-zero total length, so they exhaust the scheme. The equations define a finite algebra over $\mathbb{Q}$, hence every coordinate of a solution is algebraic over $\mathbb{Q}$, and the number of geometric points and their local lengths are the same over $\overline{\mathbb{Q}}_\ell$ and over $\mathbb{C}$. This proves the assertion and the labelling. $\square$

Remark 5.5 (Formal verification). The count and simplicity assertions of Theorem 5.4, and (R_{p-1}) for $2p - 1$ prime, have been verified in Lean 4 against the Mathlib library; Appendix A describes the files. The formal proof of the count does not use Theorem 5.3. It shows directly that every complex matching point is integral at any valuation ring of $\mathbb{C}$ whose maximal ideal contains ℓ : the roots of A are scaled by a root of smallest valuation, and ℓS_A evaluated at that root has one unit term, all others lying in the maximal ideal. Reduction of integral points is injective, which gives finiteness and the upper bound $\binom{\ell}{d}$, and the same normalization gives simplicity. For the lower bound, each reduced point is lifted over $\mathbb{Z}_\ell$ by a multivariable Hensel lemma derived from formal smoothness, and the resulting family of distinct $\mathbb{Q}_\ell$ -points is transferred to $\mathbb{C}$ through the Nullstellensatz, without identifying $\mathbb{Q}_\ell$ with a subfield of $\mathbb{C}$. The labelling (5.12) is the injective reduction map of that proof.

The labels (5.12) describe reductions of algebraic coefficients; they are not assertions that the complex roots of A , or the characteristic roots of P_A , are integers. For example, $A = \rho^2 + 1/11$ at $p = 3$ reduces to $(\rho - 2)(\rho - 3)$ modulo five.

The endpoints of v_A are its limits as $z \rightarrow \mp\infty$, namely 0 and $s_p A(0)$. Pairs with $A(0) = 0$ have equal endpoints and are pulses; the others are fronts. The pulses are needed separately, because the elliptic count of Section 6 degenerates onto them.

Proposition 5.6 (Equal endpoints). *Assume (R_{p-1}) , and let $\mathcal{Z}_p = \mathcal{X}_p \cap \{A(0) = 0\}$, with its induced scheme structure. Then*

$$\text{length } \mathcal{Z}_p = S_p := \binom{2p-2}{p-2}. \quad (5.16)$$

If $\mathcal{X}_p$ is reduced, these are S_p distinct pairs and $P_A(0) \neq 0$ at each of them. In particular, under the prime-index hypothesis, exactly

$$\binom{2p-2}{p-2}$$

normalized pairs have equal zero endpoints. The remaining $\binom{2p-2}{p-1}$ have unequal endpoints. These are counts over $\mathbb{C}$.

Proof. Polynomial summation gives $S_A(0) = -A(0)^2$, so the constant remainder satisfies the polynomial identity

$$R_0 = -A(0)(A(0) + C_A(0)). \quad (5.17)$$

Set $\alpha_d = A(0) = 0$. The equation $R_0 = 0$ disappears, leaving $d-1$ equations of weights $d+2, \dots, 2d$ in variables of weights $1, \dots, d-1$. Their leading forms have no nonzero common zero by (R_d) . The weighted calculation in Theorem 5.3 gives

$$\text{length } \mathcal{Z}_p = \frac{(d+2) \cdots (2d)}{(d-1)!} = \binom{2d}{d-1}.$$

For $d=1$ the system is empty and its length is one.

If $\mathcal{X}_p$ is finite and reduced, its coordinate algebra is a product of copies of $\mathbb{C}$; the same is true of any quotient, in particular that of $\mathcal{Z}_p$, so $\mathcal{Z}_p$ is reduced as well. At a point with $A(0) = C_A(0) = 0$, the differential of (5.17) vanishes. Such a point cannot be simple in the square matching system. Therefore $C_A(0) \neq 0$, and $P_A(0) = s_p C_A(0)/2 \neq 0$, at every point of $\mathcal{Z}_p$ when $\mathcal{X}_p$ is reduced. The prime-index assertions now follow from Theorem 5.4 and $N_p - S_p = \binom{2p-2}{p-1}$. $\square$

Under the labelling (5.12) of Theorem 5.4, a pair is a pulse exactly when its label set contains 0. Indeed $A(0) = 0$ forces 0 to be a root of the reduced polynomial $\bar{A}$, and the two counts agree: the $(p-1)$ -element subsets of $\mathbb{F}_\ell$ containing 0 number $\binom{2p-2}{p-2} = S_p$, and those avoiding 0 number $\binom{2p-2}{p-1} = N_p - S_p$.

5.4 Reflection: the two pure subfamilies

The numbers N_p and M_p refer to the full monic family. We now count the pairs whose equation is purely dispersive or purely dissipative. These are the fixed points of the reflection $z \mapsto -z$ acting on the pairs, and their number is again a binomial coefficient.

For a monic polynomial A of degree $d = p-1$ put

$$(\iota A)(\rho) = (-1)^d A(-\rho), \quad \text{that is,} \quad \alpha_j \mapsto (-1)^j \alpha_j. \quad (5.18)$$

Proposition 5.7 (Reflection). *For every monic A of degree d ,*

$$S_{\iota A}(\rho) = -S_A(-\rho) - 2A(0)A(-\rho), \quad R_{\iota A}(\rho) = -R_A(-\rho). \quad (5.19)$$

Consequently ι is an involution of the scheme $\mathcal{X}_p$. At a point of $\mathcal{X}_p$,

$$v_{\iota A}(z) = (-1)^p (v_A(-z) - s_p A(0)), \quad P_{\iota A}(\rho) = (-1)^p (P_A(-\rho) + s_p A(0)). \quad (5.20)$$

In words, ι reads a wave in the opposite direction and then restores the normalization: it moves the zero limit back to the end $e^z \rightarrow 0$, and for odd p it changes the sign of the profile. The pairs A and ιA are the mirror images that N_p counts separately. For $p = 3$ the involution is $(a, b) \mapsto (-a, b)$ in Table 3.

Proof. Since $Q(-z) = 1 - Q(z)$, and D changes sign under $z \mapsto -z$,

$$(A(D)Q)(-z) = A(0) - A(-D)Q = A(0) - (-1)^d(\iota A)(D)Q.$$

This is the first formula in (5.20). Evaluate (5.5) at $-z$, and use (5.5) for ιA on the left:

$$A(0)^2 - 2A(0)A(-D)Q - S_{\iota A}(D)Q = -S_A(0) + S_A(-D)Q.$$

Every $B(D)Q$ is a polynomial in Q without constant term, so letting $z \rightarrow -\infty$ gives $S_A(0) = -A(0)^2$. The remaining terms, and the injectivity of $B \mapsto B(D)Q$, give the first identity in (5.19). Now substitute $S_A = AC_A + R_A$ at $-\rho$:

$$S_{\iota A}(\rho) = (\iota A)(\rho) (-1)^p (C_A(-\rho) + 2A(0)) - R_A(-\rho).$$

Division by the monic polynomial ιA is unique. Hence

$$R_{\iota A}(\rho) = -R_A(-\rho), \quad C_{\iota A}(\rho) = (-1)^p (C_A(-\rho) + 2A(0)).$$

The first relation reads $R_r(\iota\alpha) = (-1)^{r+1}R_r(\alpha)$, so the linear map (5.18) preserves the ideal of the matching equations. The second relation and (5.4) give the formula for $P_{\iota A}$. $\square$

Let

$$\mathcal{X}_p^{\text{sym}} = \mathcal{X}_p \cap \{\alpha_j = 0 \text{ for odd } j\}$$

be the fixed locus of ι , with its induced scheme structure, and put

$$F_p = \binom{p-1}{\lfloor (p-1)/2 \rfloor}. \quad (5.21)$$

Table 2 in Section 9 lists the first values, with the equations to which they belong.

Theorem 5.8 (Count in the two pure subfamilies). (i) *A point A of $\mathcal{X}_p$ lies in $\mathcal{X}_p^{\text{sym}}$ if and only if P_A is reflection-symmetric. Thus $\mathcal{X}_p^{\text{sym}}$ consists of the rational-exponential pairs of the purely dispersive equations when p is even, and of the purely dissipative equations when p is odd.*

(ii) *If (R_{p-1}) holds, then $\text{length } \mathcal{X}_p^{\text{sym}} = F_p$.*

(iii) *If, in addition, $\mathcal{X}_p$ is reduced, then $\mathcal{X}_p^{\text{sym}}$ consists of F_p distinct pairs, each a simple solution of the matching equations of its subfamily. If $2p-1$ is prime, they are the points whose label in (5.12) satisfies $S = -S$.*

(iv) *For even p , every pair of $\mathcal{X}_p^{\text{sym}}$ is a pulse. For odd p , every pair of $\mathcal{X}_p^{\text{sym}}$ is a front if $\mathcal{X}_p$ is reduced, and every pair with real coefficients is a front without any hypothesis.*

Thus the rational-exponential waves of a purely dispersive equation are pulses, and those of a purely dissipative equation are fronts. The first statement holds in every even order. The second is proved whenever all pairs are distinct, and for real waves in every odd order. Elliptic waves occur in both subfamilies and are not concerned.

Proof. (i). If $\iota A = A$, then (5.20) gives $P_A(\rho) = (-1)^p(P_A(-\rho) + s_p A(0))$. For even p the polynomial A is odd, so $A(0) = 0$ and P_A is even. For odd p the sum $P_A(\rho) + P_A(-\rho)$ is the constant $-s_p A(0)$. In both cases P_A is reflection-symmetric, and

$$s_p A(0) = -2P_A(0) \quad (p \text{ odd}). \quad (5.22)$$

Conversely, let $P = P_A$ be reflection-symmetric, and apply Corollary 3.2 at the pole $i\pi$, using the period $2\pi i$. If p is even, then $v_A(-z) = v_A(z)$. Letting $z \rightarrow -\infty$ gives $A(0) = 0$, and (5.20) becomes $v_{\iota A} = v_A$. If p is odd, then $-v_A(-z) = v_A(z) + 2P(0)$, and (5.20) becomes $v_{\iota A}(z) = v_A(z) + 2P(0) + s_p A(0)$. Letting $z \rightarrow -\infty$ shows that the constant vanishes. In both cases $v_{\iota A} = v_A$, hence $\iota A = A$ by the uniqueness of the form (5.1).

(ii). On the subspace $\{\alpha_j = 0, j \text{ odd}\}$ we have $\iota A = A$, and (5.19) shows that R_A is an odd polynomial. The equations $R_r = 0$ with r even therefore hold identically there. With $m = \lfloor d/2 \rfloor$, there remain the m equations $R_1, R_3, \dots, R_{2m-1}$ in the m unknowns $\alpha_2, \alpha_4, \dots, \alpha_{2m}$. By (5.7) their weights are

$$\begin{array}{c|c} \text{variables} & 2, 4, \dots, 2m \\ \text{equations} & 2d, 2d-2, \dots, 2(d-m+1). \end{array}$$

Their highest weighted parts are the restrictions of those of the full system. The highest weighted parts of the omitted equations vanish identically on the subspace, because H_A is an odd polynomial when $\iota A = A$. A common zero of the restricted leading forms is therefore a polynomial A with $A \mid H_A$, and (R_d) gives $A = \rho^d$. The finite-cover argument of Theorem 5.3, with $\alpha_{2i} = y_i^{2i}$, gives

$$\text{length } \mathcal{X}_p^{\text{sym}} = \prod_{i=1}^m \frac{2(d-i+1)}{2i} = \binom{d}{m}.$$

(iii). A closed subscheme of a finite reduced scheme is reduced, as in the proof of Proposition 5.6, and a reduced point of a square system is a simple solution of it. The map (5.18) negates the roots of $\bar{A}$, so it acts on the labels by $S \mapsto -S$, and the labelling is a bijection.

(iv). For even p the polynomial A is odd, so $A(0) = 0$. Let p be odd, and suppose that a point of $\mathcal{X}_p^{\text{sym}}$ is a pulse, $A(0) = 0$. Then $P_A(0) = 0$ by (5.22). If $\mathcal{X}_p$ is reduced, this contradicts Proposition 5.6. If A has real coefficients, then v_A is real on the real axis, and it tends to zero with all its derivatives at both ends. Integrating (1.1) along the real axis, and using $P_A(0) = 0$, gives $\int v_A^2 dz = 0$. Hence $v_A = 0$, which is impossible. $\square$

The label count agrees with (5.21). The nonzero residues modulo $\ell = 2d + 1$ form d pairs $\{s, -s\}$. A set $S = -S$ of d elements consists of $d/2$ pairs if d is even, and of 0 and $(d-1)/2$ pairs if d is odd, which gives $\binom{d}{\lfloor d/2 \rfloor}$ sets in both cases. It contains 0 exactly when d is odd, in agreement with part (iv).

6 The count of elliptic pairs

We now count the elliptic pairs of Section 2. Fix a lattice $\Lambda \subset \mathbb{C}$, including its scale, and put $g_i = g_i(\Lambda)$. We place the pole at zero, allow P to vary over all monic polynomials of degree p , and retain the constant term of the profile. Neither reflection nor a lattice automorphism is factored out. By Section 4, every profile has the form

$$V = -s_p \mathcal{D}^{p-2} X + \sum_{j=0}^{p-3} u_j \mathcal{D}^j X + b. \quad (6.1)$$

Its only pole on $\mathbb{C}/\Lambda$ is zero, of order p . In particular Λ is its full period lattice. Let $\mathcal{E}_p(g_2, g_3)$ denote the matching equations obtained by substituting (6.1) in (1.1), in the unknowns $a_0, \dots, a_{p-1}, u_0, \dots, u_{p-3}$ and b , together with their solution set. The operator coefficients may equivalently be eliminated first, as in the proof below. Length, simplicity and reducedness are understood as in Section 5.

Theorem 6.1 (Fixed-lattice elliptic enumeration). *Assume (R_{p-1}) . For every fixed nonsingular lattice,*

$$\text{length } \mathcal{E}_p(g_2, g_3) = M_p := 2 \binom{2p-2}{p-2}. \quad (6.2)$$

If, in addition, $\mathcal{X}_p$ is reduced, then $\mathcal{E}_p(g_2, g_3)$ is reduced for (g_2, g_3) in a nonempty Zariski-open subset of $\{g_2^3 - 27g_3^2 \neq 0\}$. In particular, if $2p-1$ is prime, a generic fixed lattice admits exactly M_p normalized elliptic operator-profile pairs, all simple.

Thus the elliptic assertion on an arbitrary lattice is a count with multiplicities. Exceptional lattices, on which distinct pairs merge, do occur and cannot be discarded when counting distinct solutions; Section 9.2 determines them at order three.

The proof has five steps. We eliminate the operator, which leaves $p-1$ equations in the $p-1$ profile unknowns. We assign weights to the unknowns, under which the highest weighted parts of the equations are obtained by setting $g_2 = g_3 = 0$. We show that at this cusp the equations have only the trivial common zero; this is where (R_{p-1}) enters, and it means that there are no solutions at weighted infinity for any g_2, g_3 . Weighted Bézout then gives the length M_p on every lattice. Finally, at a nodal cubic the elliptic problem becomes the equal-endpoint case of the rational-exponential problem, whose solutions are simple by Proposition 5.6 when $\mathcal{X}_p$ is reduced, and simplicity spreads from the nodal parameter value to nearby nonsingular parameters.

Proof. Step 1: eliminating the operator. The coefficient at pole order $2p$ vanishes by the choice of s_p . The coefficients at orders $2p-1, \dots, p$ determine $a_{p-1}, \dots, a_0$ successively. At each step the coefficient of the new a_j is the nonzero rational leading coefficient of $D^j V$. This is a polynomial elimination, valid over any $\mathbb{Q}$ -algebra. The residual has pole order at most $p-1$ and zero residue. It lies in the span of

$$\{1\} \cup \{\mathcal{D}^j X : 0 \leq j \leq p-3\}.$$

Hence $p-1$ equations remain in the $p-1$ profile coefficients u_j, b .

Step 2: weights. Give D, X, g_2, g_3 weights $1, 2, 4, 6$, respectively, and give the profile weight p . Then u_j has weight $p-j-2$, and b has weight p . After the triangular elimination, the residual principal-part coefficients and the residual Laurent constant have weights

$$\begin{array}{l|l} \text{variables} & 1, 2, \dots, p-2, p \\ \text{equations} & p+1, p+2, \dots, 2p-2, 2p. \end{array} \quad (6.3)$$

The Laurent expansions and the eliminations are polynomial in g_2, g_3 , so these statements extend to the singular cubics. For fixed g_2, g_3 , the highest weighted parts of the equations in the profile variables are obtained by setting $g_2 = g_3 = 0$, since g_2 and g_3 carry positive weight.

Step 3: the cusp. At the cusp $g_2 = g_3 = 0$, $X = z^{-2}$ and $Y = -2z^{-3}$, so the profile is a rational function. Write $V = b + f(z)$, where f has its sole pole at zero and vanishes at infinity. The constant and decaying parts of the equation give

$$a_0 b + \frac{1}{2} b^2 = 0, \quad (P+b)(D)f + \frac{1}{2} f^2 = 0.$$

The leading term of f at infinity forces $a_0 + b = 0$. It follows that $b = a_0 = 0$. Under the polynomial Laplace correspondence

$$\sum_k A_k \rho^k \mapsto \sum_k k! A_k z^{-k-1},$$

differentiation corresponds to multiplication by $-\rho$, and multiplication of the rational functions corresponds to continuous convolution. After a nonzero scalar normalization, the equation for f therefore gives $A \mid H_A$ for a monic A of degree $p-1$. By (R_{p-1}) , $A = \rho^{p-1}$. Thus $f = c_* z^{-p}$, and all the variables in (6.1) vanish. The leading equations have no nonzero common zero.

Step 4: the length. The same finite-cover Bézout argument as in Theorem 5.3, now with the weights (6.3), gives

$$\text{length } \mathcal{E}_p(g_2, g_3) = \frac{(p+1)(p+2) \cdots (2p-2)(2p)}{(p-2)!p} = 2 \binom{2p-2}{p-2}.$$

For $p = 3$, this is $(4 \cdot 6)/(1 \cdot 3) = 8$. Empty products are one, so the calculation includes $p = 2$. It proves this length at every (g_2, g_3) , including the singular parameters used below; at the cusp itself the whole length sits at the single solution where every unknown vanishes. It also makes the whole coefficient family finite over $\mathbb{C}[g_2, g_3]$: sufficiently large weighted monomials reduce to bounded weight using the leading equations, which are independent of g_2, g_3 . This finiteness is what allows simplicity to be transferred from the nodal parameter value to nearby nonsingular parameters in the last step.

Step 5: the node. To prove generic simplicity, consider the nodal parameters

$$g_2 = \frac{1}{12}, \quad g_3 = -\frac{1}{216}. \quad (6.4)$$

In the logistic coordinate they are realized by

$$X = Q^2 - Q + \frac{1}{12}, \quad Y = (2Q - 1)Q(1 - Q), \quad DQ = Q(1 - Q).$$

The nodal coordinate ring is

$$\mathbb{C}[X, Y] = \{f \in \mathbb{C}[Q] : f(0) = f(1)\}.$$

Indeed the right side is $\mathbb{C} + Q(1 - Q)\mathbb{C}[Q]$; it is generated by X, Y , since, with $S = Q(1 - Q)$, $S = 1/12 - X$, $SQ = (Y + S)/2$, and $Q^2 = Q - S$. The pole filtration is the ordinary polynomial degree in Q . Consequently the nodal specialization of (6.1) has the unique form

$$V = \beta + v_A, \quad A \text{ monic of degree } p-1, \quad A(0) = 0,$$

where β is the common endpoint. The coefficient change is invertible and affine over $\mathbb{Q}$.

The matching equations become, as equations of schemes,

$$A \in \mathcal{Z}_p, \quad P = P_A - \beta, \quad \beta(\beta - 2P_A(0)) = 0. \quad (6.5)$$

To see this directly, the constant equation is $P(0)\beta + \beta^2/2 = 0$. The remaining polynomial identity is $(P + \beta)(D)v_A + v_A^2/2 = 0$, equivalent by Proposition 5.1 to $A \mid S_A$ and $P + \beta = P_A$. The coefficient changes and monic division used here work over arbitrary $\mathbb{C}$ -algebras, so the correspondence preserves local multiplicities.

If $\mathcal{X}_p$ is reduced, Proposition 5.6 gives S_p simple points of $\mathcal{Z}_p$, with $P_A(0) \neq 0$ at every point. Each quadratic in (6.5) then has two distinct roots. The nodal fiber has $2S_p = M_p$ simple points. In the finite coefficient family, the image of the closed locus where the matching Jacobian vanishes is closed. Its complement contains (6.4), and hence meets the nonsingular-lattice locus in a nonempty open set. Every fiber in that open set is reduced, proving the theorem. $\square$

The factor two in M_p retains the two choices of background in (2.5). That transformation preserves the evolution equation and the lattice; it changes the operator P and the speed. At the nodal fiber it exchanges the two roots in (6.5), which are distinct when $\mathcal{X}_p$ is reduced. In that case the two backgrounds are distinct on a generic lattice, and the M_p pairs represent $M_p/2$ elliptic waves, up to translations and Galilean shifts, each counted with both of its backgrounds. On special nonsingular lattices the two backgrounds may coalesce, as Section 9.2 shows.

6.1 Elliptic pairs of purely dispersive equations

Corollary 6.2. *Let p be even and assume (R_{p-1}) . On every fixed nonsingular lattice, the elliptic pairs whose operator is an even polynomial form a closed subscheme $\mathcal{E}_p^{\text{sym}}(g_2, g_3)$ of $\mathcal{E}_p(g_2, g_3)$, defined by $u_j = 0$ for odd j , and*

$$\text{length } \mathcal{E}_p^{\text{sym}}(g_2, g_3) = 2F_p. \quad (6.6)$$

If $\mathcal{X}_p$ is reduced, $\mathcal{E}_p^{\text{sym}}(g_2, g_3)$ is reduced for (g_2, g_3) in a nonempty Zariski-open set of nonsingular parameters.

Proof. An even operator has an even profile, by Corollary 3.2. Conversely, if the profile (6.1) is even, the odd part of (1.1) is $\sum_{j \text{ odd}} a_j D^j V = 0$; the functions $D^j V$ have distinct pole orders, so $a_j = 0$ for odd j . On the subspace $u_j = 0$, j odd, the residual of Step 1 is even, and there remain the equations and unknowns of even weight in (6.3):

$$\begin{array}{c|c} \text{variables} & 2, 4, \dots, p-2, p \\ \text{equations} & p+2, p+4, \dots, 2p-2, 2p. \end{array}$$

The omitted leading forms vanish identically on the subspace, so the remaining ones have only the trivial common zero, by Step 3. Weighted Bézout gives

$$\frac{(p+2)(p+4) \cdots (2p-2)(2p)}{2 \cdot 4 \cdots (p-2)p} = 2 \binom{p-1}{p/2-1} = 2F_p.$$

At the node, reflection is $Q \mapsto 1 - Q$. An even nodal profile is $\beta + v_A$ with $\iota A = A$, so the nodal fiber is (6.5) with $A \in \mathcal{X}_p^{\text{sym}}$. If $\mathcal{X}_p$ is reduced, it has $2F_p$ simple points, by Theorem 5.8 and Proposition 5.6, and Step 5 applies without change. $\square$

For odd p there is no fixed-lattice analogue. A purely dissipative equation has a profile that is $-P(0)$ plus a function odd about the pole, and its matching system has one equation more than it has unknowns. Such elliptic pairs are therefore not expected on a generic lattice. At $p = 3$ they occur exactly when $g_2 = 0$, as (9.7) shows for $B = 0$.

7 Which orders are settled

The formulas count pairs with multiplicity, and in that form they follow from the rigidity condition (R_{p-1}). When the rational-exponential matching scheme $\mathcal{X}_p$ is reduced, its pairs are distinct, and the elliptic pairs are distinct on a generic lattice. On a special lattice several elliptic pairs can merge, while the count with multiplicity stays the same; Section 9.2 shows this at order three. Table 1 records what is known.

Table 1: Status of the counts N_p , M_p , F_p and $2F_p$. For elliptic pairs, distinctness is asserted only on a generic lattice.

Orders	Counts with multiplicity	Pairs distinct
$2p - 1$ prime	proved	proved
$2p - 1$ a prime power	proved	open, except $p = 5$
$2 \leq p \leq 73$	exact certificate	open in general
all p	conjectured	conjectured

The prime-index argument excluded solutions at infinity by reducing the continuous convolution modulo ℓ . The same argument proves (5.10) whenever $2p - 1 = \ell^e$ is an odd prime power. For $\deg A = (\ell^e - 1)/2$, the normalized continuous convolution satisfies

$$\ell^e H_A(\rho) \equiv u \rho^{\ell^e} \pmod{\ell}, \quad u \neq 0.$$

Every bilinear coefficient other than the leading one has denominator valuation at most $e - 1$, whereas the leading coefficient has valuation $-e$. This follows either from the factorial formula or from

$$\frac{i!j!}{(i+j+1)!} = \sum_{r=0}^j \frac{(-1)^r \binom{j}{r}}{i+r+1}.$$

This argument controls only the leading system. It gives the length, but no simple-root lifting statement for the discrete matching system.

More generally, the exact arithmetic certificates supplied with the paper establish (R_d) for every $d \leq 72$, hence

$$\text{length } \mathcal{X}_p = \binom{2p-1}{p-1}, \quad \text{length } \mathcal{E}_p(g_2, g_3) = 2 \binom{2p-2}{p-2} \quad (2 \leq p \leq 73). \quad (7.1)$$

In the same range, Theorem 5.8 and Corollary 6.2 give $\text{length } \mathcal{X}_p^{\text{sym}} = F_p$ and, for even p , $\text{length } \mathcal{E}_p^{\text{sym}}(g_2, g_3) = 2F_p$. Together with the ancillary proof, the certificates also cover infinite families of higher orders. They cover all complex coefficient vectors in these degrees. The ancillary proof and the full certificate are described in Appendix A.

Equation (7.1) is a statement about lengths only. At composite indices we establish simplicity here for $p = 5$ only, where $2p - 1 = 9$ is the first composite index: a separate exact calculation proves that all 126 rational-exponential points are simple; the input and its exact output are supplied in the ancillary material. Theorem 6.1 then gives 112 distinct elliptic pairs on a generic fixed lattice at that order. Inside the two pure subfamilies a direct exact computation is much smaller: it shows that $\mathcal{X}_p^{\text{sym}}$ is reduced for every $p \leq 8$, which includes the composite indices 9 and 15.

Order seven illustrates the prime-index case, since $2p - 1 = 13$. Among all monic operators of order seven there are exactly 1716 rational-exponential pairs, all simple; twenty of them belong to purely dissipative equations, and these are fronts. On every fixed nonsingular lattice the elliptic matching scheme has length 1584, and on a generic lattice its 1584 points are distinct and simple.

8 What the count means for a single equation

The count lets the operator vary. We now read it for one evolution equation (2.1) with fixed complex coefficients. Recall that (2.1) is invariant under translations and under the Galilean shifts $u \mapsto u + k$, $x \mapsto x - kt$.

Proposition 8.1 (The waves of a fixed equation). *Let $\lambda \neq 0$. Up to a translation and a Galilean shift, the travelling waves of (2.1) that are rational in $e^{\lambda(x-ct)}$, with one pole modulo $(2\pi i/\lambda)\mathbb{Z}$, are*

$$u = b_p \lambda^p v_A(\lambda(x - ct)), \quad c = -b_p \lambda^p P_A(0), \quad (8.1)$$

where A ranges over the points of $\mathcal{X}_p$ that satisfy

$$\frac{b_j}{b_p} = \lambda^{p-j} [\rho^j] P_A, \quad 1 \leq j \leq p-1. \quad (8.2)$$

The data (A, λ) and $(\iota A, -\lambda)$ describe the same wave.

Proof. Such a wave has a finite limit as $e^{\lambda(x-ct)} \rightarrow 0$, by the argument of Section 4, and a Galilean shift makes this limit zero. Then $w_* = 0$ in (2.3), and (2.4) becomes (8.2) together with the formula for c . After a translation v has the normalization of Section 2, so $v = v_A$ and $P = P_A$ by Proposition 5.1. The last assertion follows from (5.20): the two waves differ by the constant $b_p \lambda^p s_p A(0)$, which is a Galilean shift. $\square$

Let $\pi(A) = ([\rho^{p-1}]P_A, \dots, [\rho^1]P_A) \in \mathbb{C}^{p-1}$, and let $\lambda \in \mathbb{C}^*$ act on $\mathbb{C}^{p-1}$ with weights $1, 2, \dots, p-1$. Condition (8.2) says that the coefficient vector $\beta = (b_{p-1}/b_p, \dots, b_1/b_p)$ lies in the orbit of $\pi(A)$. Rescaling x moves an equation along such an orbit, which is a weighted curve. This is the reason for fixing the rate in the count: N_p counts these curves together with the waves that they carry.

Corollary 8.2 (Solvable equations). *Assume that $\mathcal{X}_p$ is finite, for instance that (R_{p-1}) holds.*

- (i) *Equation (2.1) has a travelling wave that is rational in an exponential, with one pole per period, if and only if β lies in one of the orbits $\mathbb{C}^* \pi(A)$, $A \in \mathcal{X}_p$. There are at most $(N_p + F_p)/2$ such orbits.*
- (ii) *If $\beta \neq 0$, there are finitely many such waves up to translations and Galilean shifts.*
- (iii) *If $p \geq 3$, the orbits form a set of dimension at most one in $\mathbb{C}^{p-1}$. A generic equation (2.1) has no such wave.*

Proof. Part (i) restates Proposition 8.1. For the bound, $\pi(\iota A) = (-1) \cdot \pi(A)$ by (5.20), so the orbit depends only on the ι -orbit of A . The number of ι -orbits of points is half the number of points plus half the number of fixed points, and these two numbers are at most length $\mathcal{X}_p \leq N_p$ and length $\mathcal{X}_p^{\text{sym}} \leq F_p$; the last inequality is the Bézout bound in the proof of Theorem 5.8, which does not require (R_{p-1}) . For (ii), if $\beta \neq 0$ and $\beta \in \mathbb{C}^* \pi(A)$, some coordinate $[\rho^j]P_A$ is nonzero, and (8.2) leaves at most $p - j$ values of λ . Part (iii) is a dimension count. $\square$

When $\beta = 0$ the equation is invariant under the scaling itself, and its waves come in one-parameter families of rates. Order two shows what the orbits mean. There are two of them, and both carry waves. The orbit $\beta = 0$ is the KdV equation, with its family of solitons of all rates. The orbit $\beta \neq 0$ consists of the equations with a nonzero Burgers coefficient, and each of them has the two fronts of Section 9.1, at the rates fixed by that coefficient. At $p = 3$, with $\beta = (B, \mu)$, the orbits are the curves $B^2 = \sigma\mu$ with the origin removed, and Table 3 gives the four classical values

$$\sigma = \frac{16a^2}{a^2 + 19b} \in \left\{0, 16, \frac{144}{47}, \frac{256}{73}\right\}. \quad (8.3)$$

Here the bound $(N_3 + F_3)/2 = 6$ is not attained, because the two pairs with $a = 0$ share $\sigma = 0$, and $(\pm 1, 0)$ and $(\pm i, 0)$ share $\sigma = 16$.

For odd p the classification turns Corollary 8.2 into a statement about all meromorphic travelling waves.

Corollary 8.3 (Generic equations of even order). *Let $p \geq 5$ be odd and assume (R_{p-1}) . The coefficient vectors $\beta \in \mathbb{C}^{p-1}$ for which (2.1) has a nonconstant meromorphic travelling wave lie in an algebraic subset of dimension at most two. In particular a generic equation has none. For $p = 3$ this set is the union of the four curves $B^2 = \sigma\mu$ of (8.3).*

Proof. By Theorem 3.1, such a wave is rational, rational in an exponential, or elliptic, with one pole per period. Take $\lambda = 1$ in (2.3), so that $[\rho^j]P = b_j/b_p$ for $1 \leq j \leq p-1$. The rational-exponential waves give a set of dimension at most one, by Corollary 8.2. For a rational wave (4.1), the argument of Step 3 in the proof of Theorem 6.1 applies verbatim, with the simple-pole term $r_1 z^{-1}$ included: it gives $b = a_0 = 0$, and the Laplace correspondence gives $A \mid H_A$ for a monic A of degree $p-1$, now without the restriction $A(0) = 0$. By (R_{p-1}) , $A = \rho^{p-1}$, hence $P = D^p$ and $\beta = 0$. For the elliptic waves, the family of all $\mathcal{E}_p(g_2, g_3)$ is finite over the (g_2, g_3) -plane, by Step 4 of the same proof. It has dimension two, so its image under $P \mapsto \beta$ has dimension at most two. For $p = 3$, Section 9.2 gives the elliptic condition $B^2 = 16\mu$, which is one of the four curves; the point $\beta = 0$ lies on all of them. $\square$

The same dimension count applies to the one-pole waves of every order $p \geq 3$ for which (R_{p-1}) holds: a generic equation has no travelling wave that is rational, rational in an exponential, or elliptic with one pole per period. For even p , however, Theorem 3.1 excludes other meromorphic waves only when the equation is purely dispersive.

9 Examples

The examples have one purpose: to show the counts at work, and to connect them with the classical lists. Table 2 collects the numbers for the lowest orders. Orders two and three are small enough for the full family to be treated by hand (Sections 9.1 and 9.2); the second of them also shows what multiplicity means for elliptic pairs. At higher orders we list the pairs of the two pure subfamilies (Sections 9.3 and 9.4). The normalizations are those of Section 2.

9.1 Order two: KdV–Burgers

The integrated travelling-wave equation for

$$u_t + uu_x + \nu u_{xx} + \delta u_{xxx} = 0, \quad \delta \neq 0,$$

Table 2: Counts of complex pairs. N_p and F_p count rational-exponential pairs at unit rate, in the full family and in the pure subfamily of that order. M_p and $2F_p$ count elliptic pairs on a generic fixed lattice, and are the total multiplicities on every nonsingular lattice. The status of each number is given in Table 1.

p	N_p	M_p	Pure subfamily of this order	F_p	$2F_p$
2	3	2	dispersive, $D^2 + a_0$ (KdV)	1	2
3	10	8	dissipative, $D^3 + a_1 D + a_0$ (KS without dispersion)	2	–
4	35	30	dispersive, $D^4 + a_2 D^2 + a_0$ (Kawahara)	3	6
5	126	112	dissipative, $D^5 + a_3 D^3 + a_1 D + a_0$ (Nikolaevskiy)	6	–
6	462	420	dispersive, $D^6 + a_4 D^4 + a_2 D^2 + a_0$ (seventh-order KdV)	10	20
7	1716	1584	dissipative	20	–

has order $p = 2$. Its exact front solutions have been studied in several equivalent forms [22, 27, 43]. The long-wave reductions in Kudryashov et al. [32] also contain this balance of dissipation and dispersion.

Here $A(\rho) = \rho + a$ has a single coefficient, and the matching system is one equation in one unknown. Direct polynomial division gives

$$\text{rem}_A S_A = -\frac{a(a-1)(a+1)}{6}, \quad P_A = \rho^2 + 5a\rho + a^2 - 6a - 1, \quad v_A = 12((a+1)Q - Q^2). \quad (9.1)$$

Thus there are exactly $N_2 = 3$ simple pairs:

$$\begin{array}{c|c|c} a & P_A(\rho) & v_A \\ \hline -1 & (\rho-2)(\rho-3) & -12Q^2 \\ 0 & \rho^2 - 1 & 12Q(1-Q) = 3\text{sech}^2(z/2) \\ 1 & (\rho-1)(\rho+6) & 12Q(2-Q). \end{array} \quad (9.2)$$

The pulse $a = 0$ is the KdV soliton. It is the single pair of the purely dispersive subfamily, $F_2 = 1$. The two fronts belong to mixed equations, and the reflection ι exchanges them. In the original variables, (2.3) and (2.4) read

$$u(x, t) = u_* + \delta\kappa^2 v_A(\kappa(x - ct)), \quad \nu = \delta\kappa(5a), \quad u_* - c = \delta\kappa^2(a^2 - 6a - 1).$$

In particular, a nonzero ν fixes the front rate to $\kappa = \nu/(5a\delta)$, $a = \pm 1$, in agreement with the classical parameter restriction.

For a fixed lattice the elliptic candidate is $V = -12X + b$. Its residual is

$$-12a_1 Y - 12(a_0 + b)X + 6g_2 + a_0 b + \frac{1}{2}b^2.$$

Consequently the complete elliptic system is

$$a_1 = 0, \quad v = -12\wp - a_0, \quad a_0^2 = 12g_2. \quad (9.3)$$

There is no one-pole elliptic member with nonzero Burgers coefficient: both elliptic pairs belong to the KdV equation, in agreement with $M_2 = 2F_2 = 2$. They are the two backgrounds (2.5) of one cnoidal wave. For $g_2 \neq 0$ they are distinct and simple. For $g_2 = 0$, $g_3 \neq 0$, they coincide, and the lattice carries one pair of multiplicity two.

Since $2p - 1 = 3$ is prime, the counts at order two are unconditional. The classification of Theorem 3.1 applies to the KdV subfamily $a_1 = 0$. For the mixed operators, the Fisher example in Section 10 explains why the same conclusion cannot be expected.

9.2 Order three: Kuramoto–Sivashinsky

Let $p = 3$, $s_3 = 60$, and $A(\rho) = \rho^2 + a\rho + b$. The profile and monic operator are

$$\frac{v_A}{60} = 2Q^3 - (a+3)Q^2 + (a+b+1)Q, \quad (9.4)$$

$$P_A(\rho) = \rho^3 + 4a\rho^2 + (a^2 + 19b)\rho - a^3 + 7ab - 5a - 30b. \quad (9.5)$$

The remainder of S_A is $(E_1\rho + E_0)/30$, where

$$E_0 = ab(a^2 - 7b + 5), \quad E_1 = a^4 - 8a^2b + 11b^2 + 10b - 1. \quad (9.6)$$

The system $E_0 = E_1 = 0$ is solved by cases on the factors of E_0 . If $b = 0$, then $a^4 = 1$. If $a = 0$, then $(b+1)(11b-1) = 0$. In the remaining case $a^2 = 7b - 5$, and the second equation becomes $4(b-2)(b-3) = 0$. This exhausts the system and gives Table 3. The Jacobian determinant is nonzero at every point, so all $N_3 = 10$ pairs are simple, as Theorem 5.4 predicts for $2p - 1 = 5$.

Table 3: All ten rational-exponential pairs at order three. The determinant is taken in the order (E_0, E_1) against (a, b) . Reality refers to the unit-rate logistic representatives.

a	b	Number	$\det \partial(E_0, E_1)/\partial(a, b)$	Real profiles
± 1	0	2	-24	Pulses
$\pm i$	0	2	-16	Not real
0	-1	1	144	Front
0	1/11	1	576/121	Front
± 3	2	2	-144	Fronts
± 4	3	2	384	Fronts

There are ten complex pairs, eight with real coefficients. All eight have smooth bounded representatives on the real axis: two pulses and six fronts. For example,

$$A = (\rho - 1)(\rho - 2), \quad v_A = 120Q^3, \quad P_A = (\rho - 3)(\rho - 4)(\rho - 5).$$

These belong to the classical hyperbolic-function solution sector [24, 25, 37]. The complete parameter list is tabulated by Kudryashov and Zargaryan [31, Table 1] and Conte and Musette [4, Table 1]. Their tables contain six parameter cases because they identify the two signs of the dispersion coefficient. These six cases are the orbits of the reflection ι , which is $(a, b) \mapsto (-a, b)$ here: four orbits of two pairs, and the two fixed pairs with $a = 0$. Exhaustion of this classical list follows from Eremenko [18].

The two pairs with $a = 0$ are the $F_3 = 2$ pairs of the purely dissipative subfamily, the Kuramoto–Sivashinsky equation without dispersion. Both are fronts, as Theorem 5.8 requires; the pulses $(\pm 1, 0)$ need dispersion. By Corollary 8.2, an equation with $P = D^3 + BD^2 + \mu D + a_0$ has a wave of this kind exactly when $B^2 = \sigma\mu$ for one of the four values (8.3). The pairs $(\pm 1, 0)$ and $(\pm i, 0)$ belong to the same equations, with $\sigma = 16$; their rates differ by a factor i , so for a real equation the second wave is periodic in x , and its real form has poles on the real axis.

Elliptic pairs, and the lattices on which they merge. An elliptic solution must have zero residue at its pole. For $P = D^3 + BD^2 + \mu D + a_0$ the Laurent residue is $60\mu/19 - 15B^2/76$, so

$B^2 = 16\mu$. With this restriction the entire elliptic matching scheme is

$$\begin{aligned} P &= D^3 + BD^2 + \frac{B^2}{16}D + a_0, \\ v &= -60\wp' - 15B\wp - \frac{B^3}{64} - a_0, \\ B^4 &= 3072g_2, \quad a_0^2 = \frac{13B^6}{2048} - 2160g_3. \end{aligned} \tag{9.7}$$

Read with the operator fixed, this is the classical solution of Kudryashov [25]: $w = v + a_0$ solves the usual integrated equation $w''' + Bw'' + \mu w' + w^2/2 + K = 0$ with $K = -a_0^2/2$. Necessity of the restriction $B^2 = 16\mu$ is proved by Hone [20], and the classification of all meromorphic KS solutions is completed by Eremenko [18]. For the count we read (9.7) the other way: fix g_2, g_3 and solve for B and a_0 . The coefficients eliminated along the way are determined linearly with nonzero rational pivots, so the last two equations carry the full matching system, multiplicities included. Their quotient algebra has basis $B^i a_0^j$, $0 \leq i < 4$, $0 \leq j < 2$, and length eight, which is M_3 .

On most lattices these are eight distinct pairs: four values of B , and for each of them the two backgrounds $\pm a_0$ of (2.5). Pairs merge exactly where the Jacobian of the last two equations vanishes, and that Jacobian is a nonzero constant times $B^3 a_0$. Let

$$j(\Lambda) = \frac{1728g_2^3}{g_2^3 - 27g_3^2}.$$

If $B = 0$, then $g_2 = 0$, hence $j = 0$. If $a_0 = 0$ and the lattice is nonsingular, elimination gives

$$675g_3^2 = 169g_2^3, \quad j = -300.$$

These conditions are sufficient as well. Thus:

$j(\Lambda)$	distinct pairs	multiplicities
$j \notin \{0, -300\}$	8	1, 1, 1, 1, 1, 1, 1, 1
-300	6	2, 2, 1, 1, 1, 1
0	2	4, 4.

(9.8)

The two exceptional cases show the two ways in which pairs can merge. For $j = -300$, all four values of B are distinct, and exactly two of them give $a_0 = 0$: there the two backgrounds of one wave coincide, and two double points result. For instance, $g_2 = 1/12$, $g_3 = 13/1080$ gives the double points $(B, a_0) = (\pm 4, 0)$. For $j = 0$, the four values of B collapse to $B = 0$, the operator becomes the purely dissipative $D^3 + a_0$ with $a_0^2 = -2160g_3$, and two points of multiplicity four result. On every nonsingular lattice the multiplicities add up to eight. This is the form in which Theorem 6.1 holds on an arbitrary lattice, and (9.8) is the exceptional set of that theorem at order three.

9.3 Purely dispersive equations: Kawahara and seventh-order KdV

Order four. For the Kawahara equation [14, 23], the integrated, shifted and monic travelling-wave operator is

$$P = D^4 + bD^2 + a_0. \tag{9.9}$$

The classification for this equation, using its first integral, is already established by Demina and Kudryashov [14]; see also Ye et al. [44]. Their first integral is the order-four instance of (3.7).

By Theorem 5.8, the pairs with an even operator are those with $\iota A = A$, that is, $A = \rho^3 + c\rho$. All of them are pulses: the profile is even about its pole, $v(Q) = v(1 - Q)$. The quotient and remainder are

$$\begin{aligned} P_A &= \rho^4 + 13c\rho^2 + \frac{31c^2 - 70c + 7}{3}, \\ v_A &= 280(1 + c)Q(1 - Q) - 1680Q^2(1 - Q)^2, \\ \text{rem}_A S_A &= -\frac{\rho}{420}(c + 1)(31c^2 - 31c + 10). \end{aligned} \tag{9.10}$$

The three distinct roots are

$$c = -1, \quad c = \frac{1}{2} \pm \frac{3i}{2\sqrt{31}}.$$

They give $F_4 = \binom{3}{1} = 3$ simple pairs. These are the three simply periodic parameter choices of Demina and Kudryashov [14, Section 3], in the present normalization: the parameter in their Eq. (31) equals $1/c$ here. At $c = -1$,

$$v = -105 \text{sech}^4(z/2), \quad P = (D^2 - 4)(D^2 - 9). \tag{9.11}$$

This is the familiar solitary wave of the Kawahara equation; the other two pairs are complex at unit real rate. The unrestricted order-four count is $N_4 = 35$.

All elliptic members of (9.9) have the form

$$v = -1680\wp^2 - \frac{280b}{13}\wp + 168g_2 + \frac{31b^2}{507} - a_0. \tag{9.12}$$

Substitution gives exactly two conditions:

$$F(b; g_2, g_3) := 31b^3 - 42588bg_2 - 4745520g_3 = 0, \tag{9.13}$$

$$a_0^2 = 23184g_2^2 + \frac{7980}{169}b^2g_2 + \frac{101520}{13}bg_3. \tag{9.14}$$

These are the Weierstrass profiles and parameter relations obtained from Laurent series by Demina and Kudryashov [14, Section 3], expressed in the normalization (1.1). In their notation, $b = -\beta$, $a_0^2 = C_0^2 + 12C_1$; the profile change is $v = -6(w - w_*)$, where w_* is a constant equilibrium and $a_0 = C_0 - 6w_*$. The first equation is cubic in b , and the second is monic quadratic in a_0 , so the quotient algebra has dimension six on every fixed lattice. This is the length $2F_4$ of Corollary 6.2: three values of b , each with its two backgrounds. At $(g_2, g_3) = (1, 0)$, the cubic has three distinct roots and the right side of (9.14) is nonzero at all of them, so the six pairs are simple on a nonempty open set of lattices. The unrestricted elliptic count at this order is $M_4 = 30$; elliptic formulas with nonzero odd derivative coefficients in the full fifth-order evolution equation are given by Kudryashov and Sinelshchikov [35].

Order six. The seventh-order KdV equation has the operator $P = D^6 + a_4D^4 + a_2D^2 + a_0$; its meromorphic travelling waves are studied by Dang [9]. Here $A = \rho^5 + c_1\rho^3 + c_2\rho$, $s_6 = 5544$, and the remainder of S_A has two coefficients, at ρ^3 and at ρ . Up to constant factors they are

$$\begin{aligned} 49c_1^4 - 730c_1^2c_2 + 1655c_2^2 - 1100c_1 + 2310c_2 + 1155, \\ 49c_1^3c_2 - 681c_1c_2^2 - 330c_1^2 + 693c_1c_2 + 924c_1 - 550c_2 - 1050. \end{aligned} \tag{9.15}$$

Eliminating c_2 , which is determined by c_1 at every solution, gives

$$(c_1 + 5)(c_1 + 10)(49c_1^2 - 5c_1 + 10)g_6(c_1) = 0, \quad (9.16)$$

where g_6 is an irreducible sextic without real roots; it is printed by the ancillary script. The ten roots are distinct, in agreement with $F_6 = \binom{5}{2} = 10$. Two pairs are real:

$$\begin{array}{c|c|c} A & P_A & v_A \\ \hline \rho(\rho^2 - 1)(\rho^2 - 4) & (\rho^2 - 9)(\rho^2 - 16)(\rho^2 - 25) & 10395 \operatorname{sech}^6(z/2) \\ \rho(\rho^2 - 1)(\rho^2 - 9) & (\rho^2 - 4)(\rho^2 - 25)(\rho^2 - 71) & 10395 \operatorname{sech}^4(z/2)(1 + \operatorname{sech}^2(z/2)). \end{array} \quad (9.17)$$

All ten pairs are pulses. An exact computation on a sample lattice gives $2F_6 = 20$ for the elliptic pairs of this family.

9.4 Purely dissipative equations of order five

At $p = 5$ the purely dissipative operators are $P = D^5 + a_3D^3 + a_1D + a_0$, and $A = \rho^4 + c_1\rho^2 + c_2$. The scheme $\mathcal{X}_5$ is reduced (Section 7), so there are exactly $F_5 = 6$ pairs. They are

$$A = (\rho^2 - 1)(\rho^2 - 4), \quad A = (\rho^2 - 1)(\rho^2 - 9), \quad A = (\rho^2 - 1)\left(\rho^2 + \frac{1}{5}\right), \quad (9.18)$$

and the three pairs with

$$1105c_1^3 - 2969c_1^2 + 5570c_1 - 4000 = 0, \quad c_2 = \frac{5c_1^3 + 21c_1 - 10}{79c_1 + 210}.$$

The cubic has one real root. All six pairs are fronts, as Theorem 5.8 requires, and the four with real coefficients are, after the scales are restored, the solutions (1.4) of Kudryashov and Migita [30]. For instance, $A = (\rho^2 - 1)(\rho^2 - 4)$ gives

$$v_A = 5040Q^3(10 - 15Q + 6Q^2), \quad P_A = \rho^5 - 55\rho^3 + 1254\rho - 2520,$$

a front from 0 to $5040 = -2P_A(0)$.

10 Open questions and the boundary of the classification

Two questions remain open in the enumeration, and one limitation of the classification deserves a precise statement.

Rigidity. The first question is the continuous-convolution rigidity assertion

$$0 \neq A \in \mathbb{C}[\rho], \quad A \mid \int_0^\rho A(t)A(\rho - t) dt \implies A = c\rho^{\deg A}.$$

Through the polynomial Laplace correspondence used in the proof of Theorem 6.1, it says that, after fixing the pole at zero, any rational one-pole solution of (1.1) vanishing at infinity would have to be c_*z^{-p} , with operator $P = D^p$. Algebraic and p -adic arguments prove it in the range $\deg A \leq 72$, corresponding to $2 \leq p \leq 73$, and in infinite arithmetic families of higher degrees. Rigidity in arbitrary degree remains a conjecture. By Theorems 5.3 and 6.1, it would establish both enumeration formulas with algebraic multiplicities at every order.

Simplicity. The second question is whether every point of $\mathcal{X}_p$, the rational-exponential matching system, is simple. The elliptic simplicity assertion is generic in the lattice and is derived from this one in Theorem 6.1. At a solution $S_A = AC_A$, the tangent map of the system is

$$J_A(h) = \text{rem}_A(2(A \star h) - C_A h), \quad \deg h < d. \quad (10.1)$$

The question is whether this map always has zero kernel. The prime-index argument proves this after reduction modulo ℓ , where the map becomes multiplication by a polynomial coprime to $\bar{A}$; a uniform replacement is missing at composite indices. Laurent uniqueness concerns a fixed operator, whereas (10.1) allows the operator to vary. It therefore does not by itself prove simplicity of $\mathcal{X}_p$.

Conjecture 10.1. Continuous-convolution rigidity holds in every degree, and the matching Jacobian (10.1) is invertible at every point of $\mathcal{X}_p$. Consequently $\#\mathcal{X}_p(\mathbb{C}) = \text{length } \mathcal{X}_p = \binom{2p-1}{p-1}$ for every $p \geq 2$.

The conjecture would also imply M_p simple elliptic pairs on a generic fixed lattice at every order, F_p distinct pairs in the two pure subfamilies, and that the rational-exponential waves of every purely dissipative equation are fronts (Theorem 5.8). Simplicity on every lattice is false already at order two; (9.8) describes the full exceptional set at order three.

Mixed equations of odd order. The hypothesis of Theorem 3.1 is a separate issue. For even p it requires a purely dispersive equation, and it cannot simply be dropped. The Fisher equation supplies a known illustration of the limitation: special-speed solutions were obtained by Ablowitz and Zeppetella [1], and exact solitary representations were also studied by Kudryashov [26]. Its elliptic-exponential family is also given in Kudryashov [28], Conte et al. [5, Lemma 4.4], and McCue et al. [38]. In our normalization it is

$$v(z) = -12\lambda^2 e^{2\lambda z} \wp(e^{\lambda z} - 1; 0, g_3), \quad P(D) = D^2 - 5\lambda D + 6\lambda^2, \quad \lambda \neq 0.$$

For generic $g_3 \neq 0$, this is globally meromorphic and lies outside $\mathcal{W}$. Further examples of meromorphic solutions outside $\mathcal{W}$ are discussed by Ng and Wu [40, Remark 1.11]. Substitution follows from $(D^2 - 5\lambda D + 6\lambda^2)[t^2 U(t)] = \lambda^2 t^4 U''(t)$, $t = e^{\lambda z}$. Thus a classification of the mixed equations of odd order requires a broader conclusion.

For mixed operators of even degree $p \geq 4$ we do not know whether meromorphic solutions outside $\mathcal{W}$ occur. Methods for constructing elliptic and rational-exponential solutions without the finiteness property are developed by Demina [13].

A Exact verification and ancillary material

This appendix records one lemma used in the proof of Theorem 5.4 and describes the computer checks that accompany the paper.

The lemma transfers the absence of solutions at infinity from characteristic ℓ to characteristic zero. Let homogeneous polynomials $F_1, \dots, F_m$, with coefficients in $\mathbb{Z}_{(\ell)}$, have no nonzero common zero over $\overline{\mathbb{F}}_\ell$. Then they have no nonzero common zero over $\mathbb{C}$. Indeed a complex projective zero yields a zero over $\overline{\mathbb{Q}}$ after fixing one nonzero coordinate. Choose a place above ℓ , extend it if necessary, and scale all coordinates by the inverse of a coordinate of smallest valuation. The coordinates are integral and one is a unit. Their reductions give a nonzero common zero over

$\overline{\mathbb{F}}_\ell$, contradicting the hypothesis. The argument needs the algebraic closure of the residue field; a search only over $\mathbb{F}_\ell$ would not suffice.

The ancillary material contains four kinds of checks.

1. `verify_manuscript.py` reconstructs the discrete and continuous convolutions, the low-order Laurent coefficients, and the Weierstrass differential algebra. It also checks the leading-term comparison used in Proposition 5.2. It checks the displayed order-two, order-three and order-four formulas by exact substitution, including the correspondence with the KS tables of Conte and Musette [4], Kudryashov and Zargaryan [31], the KS collision multiplicities, the Kawahara coefficient restriction and its literature normalization, the nodal correspondence, and the Fisher identity. These finite substitutions check the displayed formulas; the uniform proofs are the arguments in the text.
2. `p5_count.sing` is a self-contained rational SINGULAR calculation for the full order-five matching ideal. Its quotient dimension is 126. Adjoining the matching Jacobian determinant gives the unit ideal, so all 126 points are simple. The input is reconstructed from the original polynomial sum by the Python checker; the saved transcript states the two exact outputs.
3. `rigidity/` supplies the proof and exact certificate for (R_d) , $d \leq 72$. For $2d + 1 = \ell^e + g$, $g \in \{0, 2, 4, 6\}$, the proof reduces the normalized continuous convolution to a residual system of degree at most g . The certificate retains every residual degree, with integer ideal-membership identities and certified prime factors. The replay reconstructs the inputs, multiplies out every identity, and checks all 72 arithmetic choices. It does not infer characteristic-zero nonexistence from a list of residue-field points.
4. `symmetry_checks/` contains the exact checks for the two pure subfamilies and for single equations: the identities of Proposition 5.7 with symbolic coefficients for $d \leq 5$; the lengths F_p and the reducedness of $\mathcal{X}_p^{\text{sym}}$ for $p \leq 8$; the lengths $2F_p$ of Corollary 6.2 for $p \leq 6$ on a sample lattice; the six pairs (9.18) and the ten pairs (9.16), with the sextic factor in full; and the scaling (8.1) by substitution in the evolution equation.

A Lean 4 formalization accompanies the paper at <https://github.com/migita/binomial-count-meromorphic-waves>, directory `lean/`, with Lean and Mathlib pinned to version 4.33.1. It proves (R_d) for $2d + 1$ prime (`rigidity_prime_index`) and the count and simplicity statements of Theorem 5.4, both for the matching system (5.6) (`primeIndexCountAndSimplicity`) and for the normalized pairs (P, v) as solutions of (1.1) in the field of rational functions of e^z (`primeIndexRationalExponentialCountAndSimplicity`), together with the correspondence between the two formulations for every $p \geq 2$. The discrete convolution is defined there by the Bernoulli-polynomial power sums, and the remainder equations and their Jacobian are those of Section 5. The proof route is described in Remark 5.5. The files `EnumerationCheck.lean` and `Check.lean` verify that the proofs have exactly the stated propositions as their types and that the theorems depend only on the axioms `propext`, `Classical.choice` and `Quot.sound`; the project contains no `sorry`, added axiom or `native_decide`.

The scripts use exact rational or finite-field arithmetic. Instructions, dependency versions and checksums are included with the source package. The prime-index theorem does not depend on the computer calculations.

Tool and computational resource disclosure

This paper was written with substantial use of generative AI, which is disclosed here in the form recommended by the Leiden Declaration on Artificial Intelligence and Mathematics¹ and required by arXiv. No automated system is an author; credit and responsibility for the results, and for the attribution of prior work, rest with the author.

- **gpt-6-astra** (OpenAI), August–September 2026. The main mathematical ideas of the paper, the proofs of Theorems 3.1, 5.3, 5.4 and 6.1, the rigidity condition (R_d) and its certificate, the exact verification scripts, the SINGULAR input, the first complete draft of the text, and the Lean 4 formalization of Remark 5.5 were produced by this model in dialogue with the author.
- OpenAI Codex v0.153.4, September 2026. Independent mathematical audit of the final revision, integration of the edits, and rebuilding and validation of the source package.
- Claude Code (Anthropic, Claude Fable 5.1), September 2026. The literature search behind the attributions in Section 1, the readability revision of all sections, and the leading-coefficient argument of Proposition 5.2, which Codex then checked independently. In the final revision it proposed and proved the results on the two pure subfamilies and on single equations (Corollary 3.2, Section 5.4, Corollary 6.2 and Section 8), with their exact checks, and reorganized the manuscript around them. It also audited the Lean formalization independently: the formal statements against the manuscript, a rebuild from source, and the axiom audit.

References

- [1] Mark J. Ablowitz and Anthony Zeppetella. Explicit solutions of Fisher’s equation for a special wave speed. *Bulletin of Mathematical Biology*, 41(6):835–840, 1979. doi:[10.1007/BF02462380](https://doi.org/10.1007/BF02462380).
- [2] Keith Conrad. A multivariable Hensel’s lemma. University of Connecticut, expository notes, n.d. URL <https://kconrad.math.uconn.edu/blurbs/gradnumthy/multivarhensel.pdf>. Theorem 3.3.
- [3] Robert Conte and Micheline Musette. The Painlevé methods. In A. Kundu, editor, *Classical and Quantum Nonlinear Integrable Systems: Theory and Application*, pages 39–63. IOP Publishing, Bristol, 2003. doi:[10.1201/9781420034615.ch2](https://doi.org/10.1201/9781420034615.ch2). URL <https://arxiv.org/abs/nlin/0211048>.
- [4] Robert Conte and Micheline Musette. Elliptic general analytic solutions. *Studies in Applied Mathematics*, 123(1):63–81, 2009. doi:[10.1111/j.1467-9590.2009.00447.x](https://doi.org/10.1111/j.1467-9590.2009.00447.x).
- [5] Robert Conte, Tuen-Wai Ng, and Cheng-Fa Wu. Hayman’s classical conjecture on some nonlinear second-order algebraic ODEs. *Complex Variables and Elliptic Equations*, 60(11):1539–1552, 2015. doi:[10.1080/17476933.2015.1033414](https://doi.org/10.1080/17476933.2015.1033414). URL <https://arxiv.org/abs/1503.07074>.
- [6] Robert Conte, Tuen Wai Ng, and Chengfa Wu. Singularity methods for meromorphic solutions of differential equations. In N. Euler, editor, *Nonlinear Systems and Their Remarkable Mathematical Structures*, volume 1, pages 159–186. CRC Press, Boca Raton, 2018. URL <https://arxiv.org/abs/1806.03177>.
- [7] Robert Conte, Micheline Musette, Tuen Wai Ng, and Chengfa Wu. Methods for exact solutions of nonlinear ordinary differential equations. *Advances in Mathematics (China)*, 54(2):379–389, 2025. doi:[10.11845/sxjz.2024010a](https://doi.org/10.11845/sxjz.2024010a). URL <https://arxiv.org/abs/2503.00834>.

¹<https://leidendeclaration.ai/>, 2 June 2026.

- [8] David A. Cox, John Little, and Donal O'Shea. *Using Algebraic Geometry*, volume 185 of *Graduate Texts in Mathematics*. Springer, New York, second edition, 2005. doi:[10.1007/b138611](https://doi.org/10.1007/b138611).
- [9] Guoqiang Dang. Meromorphic solutions of the seventh-order KdV equation by using an extended complex method and Painlevé analysis. *ScienceAsia*, 49(1):108–115, 2023. doi:[10.2306/scienceasia1513-1874.2022.133](https://doi.org/10.2306/scienceasia1513-1874.2022.133).
- [10] Guoqiang Dang. Elliptic and multiple-valued solutions of some higher order ordinary differential equations. *Electronic Research Archive*, 31(10):5946–5958, 2023. doi:[10.3934/era.2023302](https://doi.org/10.3934/era.2023302).
- [11] M. V. Demina and N. A. Kudryashov. Doubly periodic meromorphic solutions of autonomous nonlinear differential equations. *Automatic Control and Computer Sciences*, 48(7):633–641, 2014. doi:[10.3103/S0146411614070207](https://doi.org/10.3103/S0146411614070207). Russian original: Modeling and Analysis of Information Systems 21(5) (2014), 49–60.
- [12] Maria V. Demina. Classification of meromorphic integrals for autonomous nonlinear ordinary differential equations with two dominant monomials. *Journal of Mathematical Analysis and Applications*, 479(2):1851–1862, 2019. doi:[10.1016/j.jmaa.2019.07.026](https://doi.org/10.1016/j.jmaa.2019.07.026).
- [13] Maria V. Demina. Meromorphic solutions of autonomous ordinary differential equations without the finiteness property. *Journal of Mathematical Analysis and Applications*, 516(2):126516, 2022. doi:[10.1016/j.jmaa.2022.126516](https://doi.org/10.1016/j.jmaa.2022.126516).
- [14] Maria V. Demina and Nikolay A. Kudryashov. From Laurent series to exact meromorphic solutions: The Kawahara equation. *Physics Letters A*, 374(39):4023–4029, 2010. doi:[10.1016/j.physleta.2010.08.013](https://doi.org/10.1016/j.physleta.2010.08.013).
- [15] Maria V. Demina and Nikolay A. Kudryashov. Explicit expressions for meromorphic solutions of autonomous nonlinear ordinary differential equations. *Communications in Nonlinear Science and Numerical Simulation*, 16(3):1127–1134, 2011. doi:[10.1016/j.cnsns.2010.06.035](https://doi.org/10.1016/j.cnsns.2010.06.035).
- [16] Maria V. Demina and Nikolay A. Kudryashov. On elliptic solutions of nonlinear ordinary differential equations. *Applied Mathematics and Computation*, 217(23):9849–9853, 2011. doi:[10.1016/j.amc.2011.04.043](https://doi.org/10.1016/j.amc.2011.04.043).
- [17] A. E. Eremenko. Meromorphic solutions of algebraic differential equations. *Russian Mathematical Surveys*, 37(4):61–95, 1982. doi:[10.1070/RM1982v037n04ABEH003967](https://doi.org/10.1070/RM1982v037n04ABEH003967). Erratum: *Russian Mathematical Surveys* 38(6) (1983).
- [18] Alexandre Eremenko. Meromorphic traveling wave solutions of the Kuramoto–Sivashinsky equation. *Journal of Mathematical Physics, Analysis, Geometry*, 2(3):278–286, 2006. URL <https://arxiv.org/abs/nlin/0504053>.
- [19] Alexandre E. Eremenko, Liangwen Liao, and Tuen Wai Ng. Meromorphic solutions of higher order Briot–Bouquet differential equations. *Mathematical Proceedings of the Cambridge Philosophical Society*, 146(1):197–206, 2009. doi:[10.1017/S030500410800176X](https://doi.org/10.1017/S030500410800176X). URL <https://arxiv.org/abs/0707.2898>.
- [20] A. N. W. Hone. Non-existence of elliptic travelling wave solutions of the complex Ginzburg–Landau equation. *Physica D: Nonlinear Phenomena*, 205(1–4):292–306, 2005. doi:[10.1016/j.physd.2004.10.011](https://doi.org/10.1016/j.physd.2004.10.011).
- [21] Zifeng Huang, Liming Zhang, Qiuhui Chen, and Wenjun Yuan. The representation of meromorphic solutions for a class of odd order algebraic differential equations and its applications. *Mathematical Methods in the Applied Sciences*, 37(10):1553–1560, 2014. doi:[10.1002/mma.3053](https://doi.org/10.1002/mma.3053).
- [22] Alan Jeffrey and Siqing Xu. Exact solutions to the Korteweg–de Vries–Burgers equation. *Wave Motion*, 11(6):559–564, 1989. doi:[10.1016/0165-2125\(89\)90026-7](https://doi.org/10.1016/0165-2125(89)90026-7).
- [23] Takuji Kawahara. Oscillatory solitary waves in dispersive media. *Journal of the Physical Society of Japan*, 33(1):260–264, 1972. doi:[10.1143/JPSJ.33.260](https://doi.org/10.1143/JPSJ.33.260).
- [24] N. A. Kudryashov. Exact soliton solutions of the generalized evolution equation of wave dynamics. *Journal of Applied Mathematics and Mechanics*, 52(3):361–365, 1988. doi:[10.1016/0021-8928\(88\)90090-1](https://doi.org/10.1016/0021-8928(88)90090-1).

- [25] N. A. Kudryashov. Exact solutions of the generalized Kuramoto–Sivashinsky equation. *Physics Letters A*, 147(5–6):287–291, 1990. doi:[10.1016/0375-9601\(90\)90449-X](https://doi.org/10.1016/0375-9601(90)90449-X).
- [26] N. A. Kudryashov. Exact solitary waves of the Fisher equation. *Physics Letters A*, 342(1–2):99–106, 2005. doi:[10.1016/j.physleta.2005.05.025](https://doi.org/10.1016/j.physleta.2005.05.025).
- [27] N. A. Kudryashov. Seven common errors in finding exact solutions of nonlinear differential equations. *Communications in Nonlinear Science and Numerical Simulation*, 14(9–10):3507–3529, 2009. doi:[10.1016/j.cnsns.2009.01.023](https://doi.org/10.1016/j.cnsns.2009.01.023).
- [28] N. A. Kudryashov. Meromorphic solutions of nonlinear ordinary differential equations. *Communications in Nonlinear Science and Numerical Simulation*, 15(10):2778–2790, 2010. doi:[10.1016/j.cnsns.2009.11.013](https://doi.org/10.1016/j.cnsns.2009.11.013). URL <https://arxiv.org/abs/1201.0126>.
- [29] N. A. Kudryashov. Method of the logistic function for finding analytical solutions of nonlinear differential equations. *Modeling and Analysis of Information Systems*, 22(1):23–37, 2015. doi:[10.18255/1818-1015-2015-1-23-37](https://doi.org/10.18255/1818-1015-2015-1-23-37). URL <https://www.mathnet.ru/eng/mais426>.
- [30] N. A. Kudryashov and A. V. Migita. Periodic structures developing with account for dispersion in a turbulence model. *Fluid Dynamics*, 42(3):463–471, 2007. doi:[10.1134/S0015462807030143](https://doi.org/10.1134/S0015462807030143).
- [31] N. A. Kudryashov and E. D. Zargaryan. Solitary waves in active-dissipative dispersive media. *Journal of Physics A: Mathematical and General*, 29(24):8067–8077, 1996. doi:[10.1088/0305-4470/29/24/029](https://doi.org/10.1088/0305-4470/29/24/029).
- [32] N. A. Kudryashov, D. I. Sinelshchikov, and I. L. Chernyavsky. Nonlinear evolution equations for description of perturbations in a viscoelastic tube. *Russian Journal of Nonlinear Dynamics*, 4(1):69–86, 2008. doi:[10.20537/nd0801004](https://doi.org/10.20537/nd0801004). In Russian.
- [33] Nikolay A. Kudryashov. One method for finding exact solutions of nonlinear differential equations. *Communications in Nonlinear Science and Numerical Simulation*, 17(6):2248–2253, 2012. doi:[10.1016/j.cnsns.2011.10.016](https://doi.org/10.1016/j.cnsns.2011.10.016).
- [34] Nikolay A. Kudryashov. Polynomials in logistic function and solitary waves of nonlinear differential equations. *Applied Mathematics and Computation*, 219(17):9245–9253, 2013. doi:[10.1016/j.amc.2013.03.032](https://doi.org/10.1016/j.amc.2013.03.032).
- [35] Nikolay A. Kudryashov and Dmitry I. Sinelshchikov. Elliptic solutions for a family of fifth order nonlinear evolution equations. *Applied Mathematics and Computation*, 218(12):6991–6997, 2012. doi:[10.1016/j.amc.2011.12.058](https://doi.org/10.1016/j.amc.2011.12.058).
- [36] Nikolay A. Kudryashov, Dmitry I. Sinelshchikov, and Maria V. Demina. Exact solutions of the generalized Bretherton equation. *Physics Letters A*, 375(7):1074–1079, 2011. doi:[10.1016/j.physleta.2011.01.010](https://doi.org/10.1016/j.physleta.2011.01.010).
- [37] Y. Kuramoto and T. Tsuzuki. Persistent propagation of concentration waves in dissipative media far from thermal equilibrium. *Progress of Theoretical Physics*, 55(2):356–369, 1976. doi:[10.1143/PTP.55.356](https://doi.org/10.1143/PTP.55.356).
- [38] Scott W. McCue, Maud El-Hachem, and Matthew J. Simpson. Exact sharp-fronted travelling wave solutions of the Fisher–KPP equation. *Applied Mathematics Letters*, 114:106918, 2021. doi:[10.1016/j.aml.2020.106918](https://doi.org/10.1016/j.aml.2020.106918).
- [39] Pinaki Mondal. *How Many Zeroes? Counting Solutions of Systems of Polynomials via Toric Geometry at Infinity*, volume 2 of *CMS/CAIMS Books in Mathematics*. Springer, Cham, 2021. doi:[10.1007/978-3-030-75174-6](https://doi.org/10.1007/978-3-030-75174-6).
- [40] Tuen-Wai Ng and Cheng-Fa Wu. Nonlinear Loewy factorizable algebraic ODEs and Hayman’s conjecture. *Israel Journal of Mathematics*, 229(1):1–38, 2019. doi:[10.1007/s11856-018-1791-0](https://doi.org/10.1007/s11856-018-1791-0). URL <https://arxiv.org/abs/1710.08593>.
- [41] NIST. Digital library of mathematical functions, n.d. URL <https://dlmf.nist.gov/>. Chapters 22 and 23.
- [42] Eman Simbawa, Paul C. Matthews, and Stephen M. Cox. Nikolaevskiy equation with dispersion. *Physical Review E*, 81(3):036220, 2010. doi:[10.1103/PhysRevE.81.036220](https://doi.org/10.1103/PhysRevE.81.036220).

- [43] M. Vlieg-Hulstman and W. D. Halford. The Korteweg–de Vries–Burgers’ equation: a reconstruction of exact solutions. *Wave Motion*, 14(3):267–271, 1991. doi:[10.1016/0165-2125\(91\)90041-L](https://doi.org/10.1016/0165-2125(91)90041-L).
- [44] Feng Ye, Jian Tian, Xiaoting Zhang, Chunling Jiang, Tong Ouyang, and Yongyi Gu. All traveling wave exact solutions of the Kawahara equation using the complex method. *Axioms*, 11(7):330, 2022. doi:[10.3390/axioms11070330](https://doi.org/10.3390/axioms11070330).
- [45] Wenjun Yuan, Yezhou Li, and Jianming Qi. All meromorphic solutions of some algebraic differential equations and their applications. *Advances in Difference Equations*, 2014:105, 2014. doi:[10.1186/1687-1847-2014-105](https://doi.org/10.1186/1687-1847-2014-105).
- [46] Wenjun Yuan, Yonghong Wu, Qiuhui Chen, and Yong Huang. All meromorphic solutions for two forms of odd order algebraic differential equations and its applications. *Applied Mathematics and Computation*, 240:240–251, 2014. doi:[10.1016/j.amc.2014.04.099](https://doi.org/10.1016/j.amc.2014.04.099).