
Exposed Maxwell mechanism experiments: physical materials selected by the original objective
Same saved AP pair for all three continuation branches; only the initial normal memory differs. No corrector and no GT selection.
Truth
Original material
AP shadow
Ordinary DR
Fractional event
Integer event
A: slotted shell
Offline reference
NRMSE 0.5151
residual / δ = 56.255
NRMSE 0.5211
residual / δ = 62.301
NRMSE 0.1871
residual / δ = 0.958
NRMSE 0.1778
residual / δ = 0.973
NRMSE 0.1861
residual / δ = 0.955
B: persistent failure
Offline reference
NRMSE 0.3658
residual / δ = 30.080
NRMSE 0.3680
residual / δ = 41.173
NRMSE 0.3658
residual / δ = 30.080
initial incumbent retained
NRMSE 0.3658
residual / δ = 30.080
initial incumbent retained
NRMSE 0.3658
residual / δ = 30.080
initial incumbent retained
C: exposed NN case
Offline reference
NRMSE 0.2739
residual / δ = 83.510
NRMSE 0.3349
residual / δ = 24.096
NRMSE 0.3349
residual / δ = 24.096
NRMSE 0.1332
residual / δ = 0.930
NRMSE 0.1288
residual / δ = 0.930
Real material contrast
20.0
17.5
15.0
12.5
10.0
7.5
5.0
2.5
0.0
Every continuation starts after 3000 paid AP steps. Both C event results are weaker than published DR from the original NN material (NRMSE 0.07799);
this figure supports a mechanism comparison, not superiority to the original DR pipeline.