
Brosimum utile (Kunth) Oken

MORACEAE

DESCRIPTION:

- **Tree:** Up to 50 m tall. Straight and cylindrical trunk, with low buttress roots. White latex.
- **Bark:** Reddish-brown color with abundant prominent lenticels; smooth or finely cracked appearance.
- **Leaves:** Simple alternate with terminal stipule, commonly large: 12 cm long × 6 cm wide.
- **Flowers:** Bisexual axillary inflorescence, rarely unisexual, usually solitary.
- **Fruits:** Globose infrutescences in syconia, brown when mature, approximately 3 to 4 cm in diameter.

PHENOLOGY:

- **Flowering:** October.
- **Fruiting:** December to January in trees with more than 35 cm DBH.
- **Sowing:** February to April.

The data presented are average values that may vary according to environmental conditions.

USES:

Wood: Light construction, moldings, plywood veneers, medicinal.

Latex: Waterproofing boats and canoes, medicinal.

Fruits: Human consumption.



SILVICULTURE:

- **Seed:** Small, 1.5 cm, recalcitrant; approx. 400 seeds per kilogram, with low viability.
- **Propagation:** Exclusively by seeds, reaches 70 % germination in nursery with high humidity levels.
- **Rotation and Growth:** Grows slowly after the first year. In the first year, it reaches 4.4 cm DBH and 2.17 m in height. At 15 years, it can reach 40 cm DBH and at 20 years, 60 cm DBH.

BIBLIOGRAPHY:

- Castaño, N.; Cárdenas, D.; Otavio, E. (2007). SINCHI.
- Comafors (1999). ITTO & OIMT. Ecuador.

VISUAL REFERENCE:



Prioria copaifera Griseb.

LEGUMINOSAE – FABACEAE

DESCRIPTION:

- **Tree:** 40 m tall and 1.5 m DBH; open rounded crown; cylindrical and straight trunk, without buttresses.
- **Bark:** Rough, gray or reddish-brown, thick.
- **Leaves:** Compound, alternate with 2 leathery leaflets, with translucent dots, dark green color, asymmetrical base.
- **Flowers:** Without petals, creamy or white, fragrant, sessile, terminal.
- **Fruits:** Brown pods, rounded, woody, do not open at maturity; a single large, flattened seed.

PHENOLOGY:

- **Flowering** February to March
- **Fruiting** September to November
- **Sowing** September to December

The data presented are average values that may vary according to environmental conditions.

USES:

Wood: Lightweight, easy to work, veneers.

Indigenous use: Repellent.

SILVICULTURE:

- **Seed:** Does not resist storage; must be sown quickly after collection. One kilogram contains 30–35 fresh seeds.
- **Propagation:** If fresh, it germinates quickly without pre-germination treatments, reaching high percentages in 30 days. Adapts well to bag sowing.
- **Rotation and Growth:** In natural stands, DBH increments of 3.6 to 9.3 cm per year have been reported.

BIBLIOGRAPHY:

- Jiménez, M. Q. (1999). INBio. San José, Costa Rica. 186 p.

VISUAL REFERENCE:

