



Gamma-ray emission from the coalescence of binary neutron stars: an electromagnetic counterpart of gravitational radiation

Néstor Ortiz



Guadalajara, México. Noviembre, 2016.


In collaboration with:



Luis Lehner



Stephen Green



Ryan Westernacher-Schneider

Plan of the talk

1. Motivation

- The role of electromagnetic (EM) counterparts
- EM counterparts of binary systems mergers
- Short Gamma-ray Bursts afterglows

Plan of the talk

1. Motivation

- The role of electromagnetic (EM) counterparts
- EM counterparts of binary systems mergers
- Short Gamma-ray Bursts afterglows

2. An alternative EM counterpart

Plan of the talk

1. Motivation

- The role of electromagnetic (EM) counterparts
- EM counterparts of binary systems mergers
- Short Gamma-ray Bursts afterglows

2. An alternative EM counterpart

3. Preamble: the single pulsar case

- Light curves
- Pulsar configuration
- Magnetosphere models

Plan of the talk

1. Motivation

- The role of electromagnetic (EM) counterparts
- EM counterparts of binary systems mergers
- Short Gamma-ray Bursts afterglows

2. An alternative EM counterpart

3. Preamble: the single pulsar case

- Light curves
- Pulsar configuration
- Magnetosphere models

4. The binary neutron star case

Plan of the talk

1. Motivation

- The role of electromagnetic (EM) counterparts
- EM counterparts of binary systems mergers
- Short Gamma-ray Bursts afterglows

2. An alternative EM counterpart

3. Preamble: the single pulsar case

- Light curves
- Pulsar configuration
- Magnetosphere models

4. The binary neutron star case


5. Summary

1. Motivation

- The role of electromagnetic (EM) counterparts

Gravitational/EM radiation sources

LIGO Scientific Collaboration (LSC) started the era of gravitational wave (GW) astronomy.




Credit: LSC

1. Motivation

- The role of electromagnetic (EM) counterparts

Gravitational/EM radiation sources

LIGO Scientific Collaboration (LSC) started the era of gravitational wave (GW) astronomy.



Credit: LSC


1. Motivation

- The role of electromagnetic (EM) counterparts

Gravitational/EM radiation sources

LIGO Scientific Collaboration (LSC) started the era of gravitational wave (GW) astronomy.

In particular, it will probe gravity in the highly dynamic, strong field regime.



Credit: LSC


1. Motivation

- The role of electromagnetic (EM) counterparts


Gravitational/EM radiation sources

LIGO Scientific Collaboration (LSC) started the era of gravitational wave (GW) astronomy.

In particular, it will probe gravity in the highly dynamic, strong field regime.



Credit: LSC




- The GW spectrum is wide
- More detectors needed
- Complementary data needed

1. Motivation

- The role of electromagnetic (EM) counterparts

The location problem of binary systems mergers

Two interferometers are not enough to precisely locate sources




Credit: LSC/A. Mellinger

1. Motivation


- The role of electromagnetic (EM) counterparts

The location problem of binary systems mergers

Two interferometers are not enough to precisely locate sources



Credit: LSC/A. Mellinger




[Abbott et al, 2016]

1. Motivation


- The role of electromagnetic (EM) counterparts

The location problem of binary systems mergers

Two interferometers are not enough to precisely locate sources



Credit: LSC/A. Mellinger



[Abbott et al, 2016]


EM counterparts to gravitational radiation will definitely help characterize/identify/locate sources

(see next talk by Ramiro Franco)

1. Motivation

- The role of electromagnetic (EM) counterparts

Binary NS-NS/BH-NS mergers



Prime EM counterpart candidates:



In the target of aLIGO interferometers

1. Motivation

- The role of electromagnetic (EM) counterparts

Binary NS-NS/BH-NS mergers



Prime EM counterpart candidates:

1. Short Gamma-Ray Bursts (SGRBs)
2. Orphan afterglows in radio/optical bands
3. Kilonovae



Associated afterglows


- 1.1 X-Ray
- 1.2 Optical
- 1.3 Radio

(Will discuss some pros and cons in a bit)

1. Motivation

- The role of electromagnetic (EM) counterparts

Binary NS-NS/BH-NS mergers



Prime EM counterpart candidates:

1. Short Gamma-Ray Bursts (SGRBs)
2. Orphan afterglows in radio/optical bands
3. Kilonovae



(Will discuss some pros and cons in a bit)

Part of the relevance:

- Astrophysical environment information
- Heavy r-process ion nucleosynthesis
- Independent constraints of system parameters
- Constraints on NS Equation of State (EoS)

1. Motivation


- EM counterparts of binary mergers

1. Short Gamma-Ray Bursts (SGRBs)

- Relativistic ejecta along rotation axis
- Less than 2 sec duration

Leading idea: After NS-NS/BH-NS merger, accretion onto newly formed BH can lead to bright Gamma-ray emission. Possible similarity with Supernovae Ia mechanism. [Rezzolla et al 2010]

Credit: NASA



1. Motivation

- EM counterparts of binary mergers

1. Short Gamma-Ray Bursts (SGRBs)


- Relativistic ejecta along rotation axis
- Less than 2 sec duration

Leading idea: After NS-NS/BH-NS merger, accretion onto newly formed BH can lead to bright Gamma-ray emission. Possible similarity with Supernovae Ia mechanism. [Rezzolla et al 2010]

Credit: NASA

(+)

- Fairly coincident with GW emission
- Extremely bright
- Well known associated afterglows



1. Motivation

- EM counterparts of binary mergers

1. Short Gamma-Ray Bursts (SGRBs)

- Relativistic ejecta along rotation axis
- Less than 2 sec duration

Leading idea: After NS-NS/BH-NS merger, accretion onto newly formed BH can lead to bright Gamma-ray emission. Possible similarity with Supernovae Ia mechanism. [Rezzolla et al 2010]


Credit: NASA

(+)

- Fairly coincident with GW emission
- Extremely bright
- Well known associated afterglows

(-)

- Highly collimated beams: only 1 in ~ 25 events would be oriented towards Earth
- Models are still uncertain:
No clear link between merger and SGRBs



1. Motivation

- EM counterparts of binary mergers

1. Short Gamma-Ray Bursts (SGRBs)

- Relativistic ejecta along rotation axis
- Less than 2 sec duration

Leading idea: After NS-NS/BH-NS merger, accretion onto newly formed BH can lead to bright Gamma-ray emission. Possible similarity with Supernovae Ia mechanism. [Rezzolla et al 2010]

Credit: NASA



(+)

- Fairly coincident with GW emission
- Extremely bright
- Well known associated afterglows

(-)

- Highly collimated beams: only 1 in ~ 25 events would be oriented towards Earth
- Models are still uncertain:
No clear link between merger and SGRBs

However: some promising NS-NS simulations by the Illinois Relativity Group [Ruiz et al, 2016]



1. Motivation

- EM counterparts of binary mergers
[SGRB Afterglows]

1.1 X-Rays

- Believed to originate from SGRBs outflow interaction with surrounding medium
- Off-axis models show timescales of 10 hours to a few days

1. Motivation

- EM counterparts of binary mergers
[SGRB Afterglows]

1.1 X-Rays

- Believed to originate from SGRBs outflow interaction with surrounding medium
- Off-axis models show timescales of 10 hours to a few days

1.2 Optical

- Also from interactions of relativistic outflows with surrounding medium
- Multiple observed counterparts to SGRBs
- Off-axis models show afterglows much fainter and longer timescales

[van Eerten and MacFadyen, 2011]

1. Motivation

- EM counterparts of binary mergers
[SGRB Afterglows]

1.1 X-Rays

- Believed to originate from SGRBs outflow interaction with surrounding medium
- Off-axis models show timescales of 10 hours to a few days

1.2 Optical

- Also from interactions of relativistic outflows with surrounding medium
- Multiple observed counterparts to SGRBs
- Off-axis models show afterglows much fainter and longer timescales

[van Eerten and MacFadyen, 2011]

1.3 Radio

- Magnetically driven outflows prior to NS-NS merger or GW excitation of magnetized plasma post merger.
- Duration of a few minutes pre/post merger.
- Relativistic post-merger plasma could produce short, strong flare (<30 minutes).
- Mildly-/sub-relativistic material interacting with interstellar medium from SGRB could produce flares at timescales of weeks/years

1. Motivation

- EM counterparts of binary mergers
[SGRB Afterglows]

1.1 X-Rays

- Believed to originate from SGRBs outflow interaction with surrounding medium
- Off-axis models show timescales of 10 hours to a few days

1.2 Optical

- Also from interactions of relativistic outflows with surrounding medium
- Multiple observed counterparts to SGRBs
- Off-axis models show afterglows much fainter and longer timescales

[van Eerten and MacFadyen, 2011]

1.3 Radio

- Magnetically driven outflows prior to NS-NS merger or GW excitation of magnetized plasma post merger.
- Duration of a few minutes pre/post merger.
- Relativistic post-merger plasma could produce short, strong flare (<30 minutes).
- Mildly-/sub-relativistic material interacting with interstellar medium from SGRB could produce flares at timescales of weeks/years

2. Orphan afterglows



1. Motivation

- EM counterparts of binary mergers
[SGRB Afterglows]

1.1 X-Rays

- Believed to originate from SGRBs outflow interaction with surrounding medium
- Off-axis models show timescales of 10 hours to a few days

1.2 Optical

- Also from interactions of relativistic outflows with surrounding medium
- Multiple observed counterparts to SGRBs
- Off-axis models show afterglows much fainter and longer timescales

[van Eerten and MacFadyen, 2011]

(+)

- Time scales of the order of hours/days
- Not too collimated as SGRBs
- Several detections

1.3 Radio

- Magnetically driven outflows prior to NS-NS merger or GW excitation of magnetized plasma post merger.
- Duration of a few minutes pre/post merger.
- Relativistic post-merger plasma could produce short, strong flare (<30 minutes).
- Mildly-/sub-relativistic material interacting with interstellar medium from SGRB could produce flares at timescales of weeks/years

2. Orphan afterglows

1. Motivation

- EM counterparts of binary mergers
[SGRB Afterglows]

1.1 X-Rays

- Believed to originate from SGRBs outflow interaction with surrounding medium
- Off-axis models show timescales of 10 hours to a few days

1.2 Optical

- Also from interactions of relativistic outflows with surrounding medium
- Multiple observed counterparts to SGRBs
- Off-axis models show afterglows much fainter and longer timescales

[van Eerten and MacFadyen, 2011]

(+)

- Time scales of the order of hours/days
- Not too collimated as SGRBs
- Several detections

(-)

1.3 Radio

- Magnetically driven outflows prior to NS-NS merger or GW excitation of magnetized plasma post merger.
- Duration of a few minutes pre/post merger.
- Relativistic post-merger plasma could produce short, strong flare (<30 minutes).
- Mildly-/sub-relativistic material interacting with interstellar medium from SGRB could produce flares at timescales of weeks/years

2. Orphan afterglows

- Dispersion
- Low energy radiation
- Not necessarily coincident with merger
- Not always isotropic (radio signals can be isotropic but peak in months/years)

[Metzger and Berger, 2011]

1. Motivation

- EM counterparts of binary mergers

3. Kilonovae

Radioactive decay of heavy
r-process elements in ejecta

- * Of the order 10^{44} J energy release
- * $5 \times 10^4 \sim 1 \times 10^5$ times Sun luminosity
(~ 1000 brighter than a typical nova)

1. Motivation

- EM counterparts of binary mergers

3. Kilonovae

Radioactive decay of heavy
r-process elements in ejecta

- * Of the order 10^{44} J energy release
- * $5 \times 10^4 \sim 1 \times 10^5$ times Sun luminosity
(~1000 brighter than a typical nova)

(+)

- Lasts hours to ~ weeks
- Isotropic radiation
- Likely to localize a large sample of GW sources

1. Motivation

- EM counterparts of binary mergers

3. Kilonovae

Radioactive decay of heavy
r-process elements in ejecta

- * Of the order 10^{44} J energy release
- * $5 \times 10^4 \sim 1 \times 10^5$ times Sun luminosity
(~1000 brighter than a typical nova)

(+)

- Lasts hours to ~ weeks
- Isotropic radiation
- Likely to localize a large sample of GW sources

(-)

- High dispersion
- Relatively faint and red due to opacity from heavy elements
- Not necessarily coincident with merger

1. Motivation

- EM counterparts of binary mergers

3. Kilonovae

Radioactive decay of heavy r-process elements in ejecta

- * Of the order 10^{44} J energy release
- * $5 \times 10^4 \sim 1 \times 10^5$ times Sun luminosity (~ 1000 brighter than a typical nova)


(+)

- Lasts hours to \sim weeks
- Isotropic radiation
- Likely to localize a large sample of GW sources

(-)

- High dispersion
- Relatively faint and red due to opacity from heavy elements
- Not necessarily coincident with merger

One single detected event: GRB 130603B



[Tanvir et al. 2013, Berger et al. 2013]

1. Motivation

- EM counterparts of binary mergers

3. Kilonovae

Radioactive decay of heavy r-process elements in ejecta

- * Of the order 10^{44} J energy release
- * $5 \times 10^4 \sim 1 \times 10^5$ times Sun luminosity (~ 1000 brighter than a typical nova)


(+)

- Lasts hours to \sim weeks
- Isotropic radiation
- Likely to localize a large sample of GW sources

(-)

- High dispersion
- Relatively faint and red due to opacity from heavy elements
- Not necessarily coincident with merger

Overall picture



[Metzger and Berger, 2011]

Plan of the talk

1. Motivation

- The role of electromagnetic (EM) counterparts
- EM counterparts of binary systems mergers
- Short Gamma-ray Bursts afterglows

2. An alternative EM counterpart

3. Preamble: the single pulsar case

- Light curves
- Pulsar configuration
- Magnetosphere models

4. The binary neutron star case

5. Summary

2. An alternative EM counterpart

**Consider Gamma-Ray emission from binary systems
due to magnetosphere interaction prior to merger**

2. An alternative EM counterpart

**Consider Gamma-Ray emission from binary systems
due to magnetosphere interaction prior to merger**

A priori

(+)

- Almost coincidental with merger time
- Characterizes orbital behaviour

2. An alternative EM counterpart

**Consider Gamma-Ray emission from binary systems
due to magnetosphere interaction prior to merger**

A priori

(+)

- Almost coincidental with merger time
- Characterizes orbital behaviour

Key questions:


1. Is it powerful enough to be observed in Earth?

2. An alternative EM counterpart

Consider Gamma-Ray emission from binary systems
due to magnetosphere interaction prior to merger

A priori
(+)


- Almost coincidental with merger time
- Characterizes orbital behaviour



Key questions:

1. Is it powerful enough to be observed in Earth? Yes! → [Palenzuela et al, 2013]

$$L \simeq 10^{(0 \sim 2)} (B/10^{11}) \text{ Mpc}$$



2. An alternative EM counterpart

Consider Gamma-Ray emission from binary systems due to magnetosphere interaction prior to merger

A priori
(+)

- Almost coincidental with merger time
- Characterizes orbital behaviour



Key questions:

1. Is it powerful enough to be observed in Earth? Yes! → [Palenzuela et al, 2013]

$$L \simeq 10^{(0 \sim 2)} (B/10^{11}) \text{ Mpc}$$

2. Is it highly collimated radiation in general?

3. Characteristic light curves?



2. An alternative EM counterpart

Consider Gamma-Ray emission from binary systems due to magnetosphere interaction prior to merger

A priori
(+)

- Almost coincidental with merger time
- Characterizes orbital behaviour

Key questions:

1. Is it powerful enough to be observed in Earth? Yes! → [Palenzuela et al, 2013]

$$L \simeq 10^{(0 \sim 2)} (B/10^{11}) \text{ Mpc}$$



2. Is it highly collimated radiation in general?

3. Characteristic light curves?

Have to figure it out ...

Idea: Find all directions of gamma-ray emission, generate sky maps of radiation and light curves

Strategy: Extract magnetosphere EM fields from existing numerical simulations



Plan of the talk

1. Motivation

- The role of electromagnetic (EM) counterparts
- EM counterparts of binary systems mergers
- Short Gamma-ray Bursts afterglows

2. An alternative EM counterpart

3. Preamble: the single pulsar case

- Light curves
- Pulsar configuration
- Magnetosphere models


4. The binary neutron star case

5. Summary

3. Preamble: the single pulsar case

Pulsars are assumed to be rotating neutron stars.


Their strong magnetic fields power
highly collimated synchrotron radiation
in the radio/X-ray/gamma-ray band



3. Preamble: the single pulsar case

Pulsars are assumed to be rotating neutron stars.

Their strong magnetic fields power
highly collimated synchrotron radiation
in the radio/X-ray/gamma-ray band




3. Preamble: the single pulsar case

Pulsars are assumed to be rotating neutron stars.

Their strong magnetic fields power
highly collimated synchrotron radiation
in the radio/X-ray/gamma-ray band

~2,300 known pulsars
~230 millisecond pulsars

In average, ~1kPa away
Typical radius: 10~20km
Typical mass: $1.4 \sim 2M_{\odot}$
Typical magnetic field strength: $10^8 \sim 10^{11}\text{G}$



Credit: NASA/Hester et al

3. Preamble: the single pulsar case

Pulsars are assumed to be rotating neutron stars.

Their strong magnetic fields power
highly collimated synchrotron radiation
in the radio/X-ray/gamma-ray band


~2,300 known pulsars
~230 millisecond pulsars

In average, ~1kPa away
Typical radius: 10~20km
Typical mass: $1.4 \sim 2M_{\odot}$
Typical magnetic field strength: $10^8 \sim 10^{11}\text{G}$

Vela: ~11Hz

Crab: ~30Hz

PSR J1748-2446ad: 716Hz



Credit: NASA/Hester et al

3. Preamble: the single pulsar case

Pulsars are assumed to be rotating neutron stars.

Their strong magnetic fields power
highly collimated synchrotron radiation
in the radio/X-ray/gamma-ray band


~2,300 known pulsars
~230 millisecond pulsars

In average, ~1kPa away
Typical radius: 10~20km
Typical mass: $1.4 \sim 2M_{\odot}$
Typical magnetic field strength: $10^8 \sim 10^{11}\text{G}$


Vela: ~11Hz

Crab: ~30Hz

PSR J1748-2446ad: 716Hz



Credit: NASA/Hester et al



Credit: B. O' hare

3. Preamble: the single pulsar case

Pulsars are assumed to be rotating neutron stars.

Their strong magnetic fields power
highly collimated synchrotron radiation
in the radio/X-ray/gamma-ray band

~2,300 known pulsars
~230 millisecond pulsars


In average, ~1kPa away
Typical radius: 10~20km
Typical mass: $1.4 \sim 2M_{\odot}$
Typical magnetic field strength: $10^8 \sim 10^{11}\text{G}$

Vela: ~11Hz


Crab: ~30Hz

PSR J1748-2446ad: 716Hz

The problem ~30 years ago:
Reproduce observed light curves
of Crab-like pulsars



Credit: NASA/Hester et al




Credit: B. O'hare

3. Preamble: the single pulsar case

- Light curves

Some refined, light curves of observed pulsars:




Credit: Jim Napolitano

3. Preamble: the single pulsar case

- Light curves

Some refined, light curves of observed pulsars:




Credit: Jim Napolitano

To try to reproduce this light curves: assume your favourite magnetic field configuration, find all directions of emission, cut at certain observation angle.

3. Preamble: the single pulsar case

- Light curves

Some refined, light curves of observed pulsars:




Credit: Jim Napolitano

To try to reproduce this light curves: assume your favourite magnetic field configuration, find all directions of emission, cut at certain observation angle.

Key ingredient: Magnetosphere model. Will discuss some of them soon...

3. Preamble: the single pulsar case


- Pulsar configuration




Credit: Robert Mallozzi,
University of Alabama.

3. Preamble: the single pulsar case

- Pulsar configuration



Main emission process:
Synchrotron + Inverse Compton scattering



nrumiano

<http://lightandcolourinmoderntechnology.com/>

Credit: Robert Mallozzi,
University of Alabama.

3. Preamble: the single pulsar case


- Magnetosphere models

Popular/successful magnetosphere models

- Dipole in vacuum
- Force-free plasma $\longrightarrow \rho\mathbf{E} + \mathbf{J} \times \mathbf{B} = 0$
[Goldreich and Julian, 1969]

3. Preamble: the single pulsar case

- Magnetosphere models




Popular/successful magnetosphere models

- Dipole in vacuum
 - Force-free plasma $\rightarrow \rho\mathbf{E} + \mathbf{J} \times \mathbf{B} = 0$
[Goldreich and Julian, 1969]
- Different emission zones

[Dyks and Rudak, 2003]

3. Preamble: the single pulsar case

- Magnetosphere models



[Dyks and Rudak, 2003]

Popular/successful magnetosphere models

- Dipole in vacuum
- Force-free plasma $\rightarrow \rho\mathbf{E} + \mathbf{J} \times \mathbf{B} = 0$
[Goldreich and Julian, 1969]

Different emission zones


We adopt the *Two Polar Caustics Model*

Main assumptions:


- Vacuum magnetosphere
- Emission along Last Open Field Lines
- Emission up to the Light Cylinder

[Arons and Scharlemann 1979; Arons 1983;
Muslimov and Harding 2003, 2004; Dyks et al 2004]


3. Preamble: the single pulsar case




Last open field lines




3. Preamble: the single pulsar case




Last open field lines




Sky map





3. Preamble: the single pulsar case



Last open field lines



Sky map



Light curve example

Plan of the talk

1. Motivation

- The role of electromagnetic (EM) counterparts
- EM counterparts of binary systems mergers
- Short Gamma-ray Bursts afterglows

2. An alternative EM counterpart

3. Preamble: the single pulsar case

- Light curves
- Pulsar configuration
- Magnetosphere models

4. The binary neutron star case

5. Summary

4. Binary Neutron Star case

We extracting EM fields from
BNS simulations prior to merger

Data kindly provided by Carlos Palenzuela

Main ingredients:

- Plasma magnetosphere
- Force Free approximation
- Aligned magnetic dipoles

4. Binary Neutron Star case

We extracting EM fields from
BNS simulations prior to merger

Data kindly provided by Carlos Palenzuela

Main ingredients:

- Plasma magnetosphere
- Force Free approximation
- Aligned magnetic dipoles


We adopt the Separatrix Layer Model,
and extension of the Two Polar Caustic approach

[Bai and Spitkovsky 2010]


Assumptions:

- Emission along Open Field Lines
- Emission from strong current layers beyond the Light Cylinder

4. Binary Neutron Star case




$t \sim -4.6\text{ms}$ (around 1 orbit prior to merger)



Magnetic field lines

Cap regions

4. Binary Neutron Star case






Resulting sky map

$t \sim -4.6\text{ms}$
(around 1 orbit
prior to merger)


Notice not collimated
emission!

Some light curves





4. Binary Neutron Star case

Comparing the emission at different times before merger...



$t = -4.6\text{ms}$



$t = -5.4\text{ms}$

5. Summary

- Detection of EM counterparts will enable exciting advances in the field of gravitational wave astronomy

5. Summary

- Detection of EM counterparts will enable exciting advances in the field of gravitational wave astronomy
- We are exploring Gamma-ray emission due to magnetosphere interaction in BNS systems as an EM counterpart because:

- It is not highly collimated radiation in general
- It is almost coincidental with merger time
- Its characteristics are tied to the orbital behaviour
- It is powerful enough to be detected on Earth

5. Summary

- Detection of EM counterparts will enable exciting advances in the field of gravitational wave astronomy
- We are exploring Gamma-ray emission due to magnetosphere interaction in BNS systems as an EM counterpart because:

- It is not highly collimated radiation in general
- It is almost coincidental with merger time
- Its characteristics are tied to the orbital behaviour
- It is powerful enough to be detected on Earth



- Misaligned spins case.
 - Exploring EM counterparts in modified theories of gravity
- Report appearing soon... we hope.