

Corneal Biomechanical Vector Analysis

For Intracorneal Ring Segment Planning

Dr. Miguel Reis
Dr. Jordana Sandes

2026

ÍNDICE

(Clique com botão direito → Atualizar campo para gerar o índice)

Chapter 1 — The Keratoconic Cornea as a Mechanical Structure

****Corneal Biomechanical Vector Analysis for Intracorneal Ring Segment Planning**** *Part I — The Problem and Fundamentals*

****Key Point:**** Keratoconus is not merely a curved cornea — it is a disease of the stromal material. Two eyes with the same K-steep may have completely different biomechanical states. A nomogram that ignores this is treating the symptom, not the disease.

1.1 Introduction

Keratoconus is the most common corneal ectasia, with prevalence estimates ranging from 1 in 375 in population-based studies to as high as 1 in 84 when modern videokeratographic screening criteria are applied (Godefrooij et al., 2017; Torres Netto et al., 2018). Traditionally, the condition has been defined by its striking topographic features — elevated keratometric readings, inferior corneal steepening, irregular astigmatism, and progressive thinning — features that place it squarely in the domain of optical geometry. Indeed, the standard clinical workflow for keratoconus diagnosis centers on sagittal curvature maps, keratometric indices, and pachymetric profiles: measurements that describe the *shape* of the cornea but say little about its *mechanical state*. This geometric perspective has practical consequences for treatment planning. When a surgeon selects an intracorneal ring segment (ICRS) for a keratoconic eye, the decision is typically guided by nomograms that map one or two geometric parameters — keratometry, Q-value, manifest refraction — to a ring configuration. The implicit assumption is that corneal shape is a sufficient proxy for corneal mechanics. As we will demonstrate throughout this book, this assumption is fundamentally flawed. Over the past two decades, a structural and bibliographic paradigm shift has been gaining momentum in ophthalmology. The seminal work of Dupps and Roberts (2001, consolidated in their 2014 article "*Biomechanics of corneal ectasia and biomechanical treatments*"), together with the structural investigations of Meek and Knupp (2015), unequivocally established that keratoconus is, at its core, a **primarily biomechanical disease** — a disorder of stromal stress distribution, with focal degradation of the elastic material properties.

The classic geometric manifestations (elevated K, pachymetric thinning, apical protrusion) are not the pathology itself but rather the *secondary mechanical consequences (symptoms)* of a weakened structure yielding under normal intraocular pressure. Two corneas with identical K-steep values may harbor vastly different stress states, material stiffnesses, and progression propensities. Treating them identically and on purely geometric grounds, as a classical nomogram does, is to confuse the symptom (shape) with the disease (mechanical weakness).

This opening chapter establishes the biomechanical foundation upon which the entire AVBC framework rests. We begin with the fibrillar architecture of the corneal stroma — the load-bearing structure that determines mechanical behavior — and progress through constitutive modeling, the decompensation cycle that drives keratoconus progression, and the clinical evidence for a biomechanical, rather than purely geometric, understanding of the disease. The reader who grasps this foundation will understand why rational ICRS planning requires a biomechanical language, not a geometric look-up table.

1.2 Fibrillar Anatomy of the Corneal Stroma

****For the Clinician:**** This section explains *why* the cornea behaves as it does under pressure. The central concept is simple: the cornea is a composite material (like fiberglass), with collagen fibers (providing stiffness) embedded in a soft matrix (providing

flexibility). The orientation of these fibers determines where the cornea is strong and where it is vulnerable. The corneal stroma constitutes approximately 90% of the corneal thickness and is the primary determinant of its mechanical behavior. It is composed of 200 to 500 collagen lamellae, each 1–2 μm thick and 10–250 μm wide, arranged in a highly organized matrix that serves a dual purpose: optical transparency and mechanical load bearing (Komai & Ushiki, 1991; Meek & Knupp, 2015).

Collagen Fiber Organization

The predominant collagen types in the corneal stroma are Type I (approximately 70% by dry weight) and Type V (approximately 15%), with minor contributions from Types III, VI, and XII. Type I collagen provides tensile strength, while Type V regulates fibril diameter and spacing — a critical factor for transparency (Birk et al., 1990). The fibrils are remarkably uniform in diameter (approximately 31 nm in humans), and their regular spacing (approximately 56 nm center-to-center) produces destructive interference of scattered light in all directions except forward, resulting in the cornea's remarkable transparency (Maurice, 1957; Meek, 2009).

Wide-angle X-ray scattering (WAXS) studies, conducted most extensively by Meek, Boote, and collaborators at Cardiff University, have revealed that collagen fiber orientation is not random but follows a characteristic pattern that varies by region (Meek & Boote, 2004; Meek & Knupp, 2015):

Central cornea: Fibers are preferentially oriented along the nasal-temporal and superior-inferior meridians, creating an approximately orthogonal arrangement aligned with the cardinal directions. This creates an anisotropic mechanical response — the central cornea is stiffer along these preferred directions.

Limbus and peripheral cornea: Fibers adopt a predominantly circumferential (tangential) orientation, forming an annular reinforcement analogous to barrel hoops. This limbal ring anchors the cornea to the sclera and resists radial expansion under intraocular pressure (IOP).

Transition zone (paracentral): A gradual shift occurs from cardinal to circumferential orientation, with increased fiber dispersion.

Anterior versus Posterior Stroma

There is a critical distinction between the anterior and posterior stroma. The anterior third exhibits extensive lamellar *interweaving* — the lamellae branch, merge, and insert into Bowman's layer at oblique angles, creating a mechanically interconnected mesh (Komai & Ushiki, 1991; Winkler et al., 2013). The posterior two-thirds, in contrast, feature more parallel, flat lamellae with less interweaving. This architectural difference has direct mechanical consequences:

The anterior stroma is approximately **two to three times stiffer** than the posterior stroma when measured by nanoindentation (Thomasy et al., 2014) or Brillouin microscopy (Scarcelli et al., 2015).

The posterior stroma is more susceptible to shear failure and lamellar slippage.

In keratoconus, the posterior stroma fails first (Dupps & Roberts, 2014). The earliest histopathological changes — fiber disorganization, breaks in lamellar continuity — occur in the posterior layers, consistent with the observation that posterior elevation abnormalities are among the earliest tomographic signs of subclinical keratoconus.

The Proteoglycan Matrix

Between the collagen fibrils lies the interfibrillar matrix, composed primarily of proteoglycans — keratan sulfate proteoglycans (lumican, keratocan, mimecan) and dermatan sulfate proteoglycans (decorin, biglycan). These molecules regulate fibril spacing and hydration, contributing to both transparency and mechanical coupling between lamellae (Quantock et al., 2010). In keratoconus, proteoglycan expression is altered, and the relationship between matrix hydration and fibril spacing becomes disrupted — an alteration that precedes observable thinning (Meek et al., 2005).

For constitutive modeling purposes, the key conclusion is that the corneal stroma is a **fiber-reinforced composite** — an extracellular matrix reinforced by collagen fibers with direction-dependent stiffness. Any realistic mechanical model must capture this anisotropy.

1.3 The Cornea as an Anisotropic Hyperelastic Shell

Why Linear Elasticity Fails

The earliest biomechanical models of the cornea treated it as a linearly elastic, isotropic material — characterized by a single Young's modulus E and Poisson's ratio ν . This simplification is inadequate for several reasons:

Nonlinearity: Corneal tissue exhibits pronounced strain-stiffening. At low strains (physiological range), the tissue is relatively compliant; at higher strains, collagen fibers become taut and stiffness increases dramatically. This J-shaped stress-strain curve cannot be captured by a constant E .

Anisotropy: As described above, the mechanical response depends on fiber orientation. Stiffness measured along collagen fibers differs substantially from stiffness measured perpendicular to them.

Near-incompressibility: Hydrated corneal tissue has a water content of approximately 78%, making it nearly incompressible (Poisson's ratio approaching 0.5).

Viscoelasticity: The cornea exhibits time-dependent mechanical behavior — creep under sustained load and stress relaxation. However, for quasi-static analyses (such as IOP loading or ICRS implantation), hyperelastic models are generally sufficient.

The Holzapfel-Gasser-Ogden (HGO) Model

****For the Clinician:**** The HGO model is the mathematical equation we use to simulate the cornea on the computer. You don't need to memorize the equation — what matters are the ****5 parameters**** in the table below. Each describes a different aspect of the corneal material, and our simulations use these values throughout this book.

Among the constitutive models available for biological soft tissues, the Holzapfel-Gasser-Ogden (HGO) model has emerged as the standard for corneal biomechanics (Holzapfel et al., 2000; Pandolfi & Manganiello, 2006; Nguyen et al., 2018). The HGO model was originally developed for arterial wall mechanics but is well suited to any fiber-reinforced tissue where fibers are embedded in a softer ground substance.

The strain energy function is:

$$\Psi = c(\bar{I}_1 - 3) + (k_1/2k_2) \sum_i \{ \exp[k_2(\kappa(\bar{I}_1 - 3) + (1 - 3\kappa)(\bar{I}_{4i} - 1))^2] - 1 \}$$

where:

Parameter	Symbol	Value (FEBio)	Physical Meaning
Matrix shear modulus	c	0.05 MPa	Stiffness of the ground substance (proteoglycan matrix without fibers)
Fiber stiffness	k_1	0.22 MPa	Collagen fiber stiffness at small strain
Fiber nonlinearity	k_2	100 (dimensionless)	Rate of exponential stiffness increase; larger values $\rightarrow$ sharper J-curve

Fiber dispersion	α	0.09 (dimensionless)	Degree of fiber alignment: 0 = perfectly aligned; 1/3 = random (isotropic)
Bulk modulus	k	4.76 MPa	Resistance to volumetric deformation (near-incompressibility)

Table 1.1. HGO material parameters used in FEBio 4.12 simulations throughout this book. Values from Nguyen et al. (2018).

Clinical interpretation of each parameter:

c = 0.05 MPa represents the baseline stiffness of the cornea *without* the contribution of collagen fibers — only the proteoglycan matrix. It is relatively soft, comparable to a firm gel. In keratoconus, matrix degradation further reduces c, making the tissue more compliant.

k₁ = 0.22 MPa captures the stiffness contributed by collagen fibers. This is the parameter most directly affected by collagen cross-linking (CXL), which increases k₁ by an estimated 50–200% (Wollensak et al., 2003).

k₂ = 100 controls how rapidly the tissue becomes stiffer as strain increases. A value of 100 simulates a strongly nonlinear J-curve — the cornea is compliant at physiological strains but resists large deformations. This is a protective mechanism: the tissue accommodates small IOP fluctuations but resists rupture at high pressures.

α = 0.09 indicates that fibers are *mostly* aligned (α = 0 would mean perfectly aligned; α = 1/3 would mean perfectly random). A value of 0.09 is consistent with WAXS data showing preferential, but not exclusive, tangential orientation in the central cornea.

k = 4.76 MPa enforces near-incompressibility, appropriate for a hydrated biological tissue.

The Macaulay brackets $\langle \cdot \rangle$ in the equation ensure that fibers contribute to stiffness only when stretched, not when compressed — a physically realistic condition since collagen fibers buckle under compression and do not carry load.

The first invariant I_1 captures isotropic (matrix) deformation, while the pseudo-invariants I_4 capture the stretch specifically in the fiber direction. This separation is the mathematical embodiment of the stroma's composite nature: matrix plus fibers.

1.4 The Dupps Decomensation Cycle

****Key Point:**** The Dupps cycle is why keratoconus progresses. It is a vicious cycle: weakness → deformation → more stress → more weakness. It is also why CXL works (it breaks the cycle by reinforcing fibers) and why ICRS works (it redistributes stresses).

Perhaps the most important conceptual advance in understanding keratoconus came from Dupps and Roberts, who formalized the **biomechanical decompensation cycle** — a positive feedback cycle that explains why keratoconus is progressive and why isolated geometric descriptors are insufficient (Dupps & Roberts, 2001; Dupps & Roberts, 2014).

The Cycle

The cycle proceeds as follows:

Enzymatic weakening: Elevated matrix metalloproteinase (MMP) activity — particularly MMP-2 and MMP-9 — degrades collagen and proteoglycans in a focal region of the stroma (Seppälä et al., 2006). This may be initiated by genetic predisposition, environmental factors (eye rubbing, atopy), or a combination of both.

Increased local compliance: The degraded region becomes more compliant (lower c, lower k₁ in HGO terms). Under the same IOP, it deforms more than the surrounding tissue.

Stress redistribution: According to Laplace's relation for a pressurized shell ($\sigma = PR/2t$, where P is pressure, R is radius, and t is thickness), the increased curvature at the weakened zone increases local membrane stress. Thinning (reduced t) further elevates stress.

Further deformation: Higher stress in a weaker material produces more deformation — leading to further steepening and thinning.

More enzymatic activity: Mechanical strain stimulates MMP expression through mechanotransduction pathways in keratocytes. Elevated strain in the steepened zone further promotes enzymatic degradation.

Return to step 1: The cycle repeats, creating progressive ectasia.

This is a classic **positive feedback cycle** — a vicious circle that, once initiated, drives disease progression until enzymatic stimulus attenuates (which may occur naturally in the fifth decade of life) or an external intervention breaks the cycle (such as CXL, which increases k₁ and interrupts the cycle at step 3).

The Role of IOP

A critical insight from the decompensation cycle is that **IOP is the mechanical engine of keratoconus progression** — not because IOP is elevated (it is typically normal), but because even normal IOP (15 mmHg) generates significant stress in a weakened stroma. Our finite element simulations confirm this quantitatively: a baseline cornea under a 15 mmHg IOP load shows an apical displacement of **360.9 μ m** — a substantial deformation even in a biomechanically normal stroma. In keratoconus, where regional weakening reduces local stiffness by 30–60% (Scarcelli et al., 2015), the same IOP produces disproportionately larger local deformations.

This is why IOP reduction (though theoretically attractive) is not a practical treatment for keratoconus — IOP is not the problem; material weakness is. The therapeutic target should be the material properties (CXL) or the stress distribution (ICRS).

1.5 Keratoconus: From Geometry to Biomechanics

Geometric Descriptors and Their Limitations

The standard clinical parameters for keratoconus — K-steep, K-flat, Kmax, corneal astigmatism, Q-value, thinnest pachymetry — are geometric descriptors. They describe the current shape and thickness of the cornea but provide no information about:

The underlying **material stiffness** (which varies regionally and between patients)

The **stress distribution** (which depends simultaneously on geometry, material properties, and IOP)

The **propensity for progression** (which depends on the interaction between stress and material degradation)

Two corneas with identical K-steep values of 48.0 D may have fundamentally different biomechanical profiles. One may have uniform material properties with a relatively recent and still-progressing ectasia; the other may have regionally compensated stiffness (e.g., from prior CXL or natural cross-link accumulation) with a stable ectasia. A nomogram treats them identically. Mechanics does not.

Evidence from Brillouin Microscopy

The most direct evidence for regional stiffness variation in keratoconus comes from Brillouin microscopy, a technique that measures the local longitudinal modulus in living tissues without contact (Scarcelli et al., 2012). Studies by Scarcelli, Yun, and collaborators demonstrated that:

The Brillouin frequency shift (a proxy for tissue stiffness) is **reduced by 0.1–0.3 GHz** in the cone region compared to surrounding tissue (Scarcelli et al., 2015).

This reduction is present even in **subclinical keratoconus** — eyes with normal topography but suspicious posterior elevation — suggesting that material weakness precedes geometric alteration.

The spatial distribution of the Brillouin reduction correlates with the steepening zone but is not identical to it, reinforcing the distinction between geometry and mechanics.

The Ocular Response Analyzer and Corvis ST

In clinical practice, the Ocular Response Analyzer (ORA, Reichert) and Corvis ST (Oculus) provide indirect measurements of corneal biomechanics through air-puff tonometry. The key parameters — Corneal Hysteresis (CH), Corneal Resistance Factor (CRF), and the Corvis Biomechanical Index (CBI) — capture aspects of the viscoelastic response to deformation (Luce, 2005; Roberts & Dupps, 2014).

In keratoconus, CH and CRF are consistently reduced, and this reduction often **precedes detectable topographic changes** (Fontes et al., 2010). This temporal sequence — biomechanical alteration before geometric alteration — is strong evidence that the disease is fundamentally biomechanical and that geometric descriptors are *lagging indicators* of an ongoing mechanical process.

The Case for Biomechanical Assessment

If geometric descriptors are lagging indicators and biomechanical assessment can detect abnormalities earlier, it follows that any intervention planning that relies exclusively on geometry is inherently delayed — it responds to the *consequences* of the disease rather than to its *current state*. The clinical implication is profound: rational ICRS planning must incorporate biomechanical assessment, not depend exclusively on keratometric and pachymetric data.

1.6 Implications for ICRS Planning

If keratoconus is fundamentally a biomechanical disease, it follows logically that an effective intervention must address the biomechanics — not merely the geometry. The ICRS, as a structural implant inserted into the stroma, is inherently a biomechanical intervention. It modifies the stress distribution, alters the displacement field, and constrains stromal deformation at the implantation site. However, current planning approaches treat the procedure as if it were a geometric intervention — selecting ring parameters based solely on topographic measurements.

The disconnect is evident. Current nomograms prescribe ring thickness based on K-steep or Q-value, implicitly assuming that keratometry captures enough of the biomechanical state to predict the ring's effect. But our FEM simulations reveal that the ICRS effect depends on at least three independent mechanical variables — radial displacement (VR), tangential stress redistribution (VT), and asymmetric torque (V τ) — each modulated by different ring parameters. Thickness controls VR; arc length controls VT; ring asymmetry controls V τ . A nomogram that maps K-steep to thickness captures only one of these three mechanisms, and even that imperfectly.

The AVBC framework, introduced in the following chapters, proposes to bridge this gap. By decomposing the ICRS effect into three biomechanical vectors, each linked to a specific ring parameter, it provides a language for mechanistic planning. The surgeon who understands VR, VT, and V τ can reason about *why* a particular ring configuration produces a specific effect — rather than consulting a look-up table and hoping for the best outcome.

1.7 Summary

Keratoconus is a **biomechanical disease**, not merely a geometric deformity. The underlying process involves collagen and matrix degradation, stress redistribution, and a positive feedback cycle (the Dupps decompensation cycle).

The corneal stroma is an **anisotropic, hyperelastic, fiber-reinforced composite**. Its mechanical behavior is best captured by the HGO constitutive model, with parameters $c = 0.05$ MPa (matrix), $k_1 = 0.22$ MPa (fiber), $k_2 = 100$ (nonlinearity), and $\kappa = 0.09$ (dispersion).

Geometric descriptors (K-steep, Q-value, pachymetry) are lagging indicators that describe consequences, not causes.

Biomechanical assessment — via Brillouin microscopy, ORA/Corvis, or constitutive modeling — provides earlier and more informative characterization.

Even **normal IOP (15 mmHg)** produces significant corneal deformation ($uz = 360.9$ μ m in our FEM baseline), highlighting IOP as the mechanical engine of ectatic progression.

Rational ICRS planning demands understanding of the biomechanical mechanisms of the intervention — not merely the geometric outcome. This is the premise of the AVBC framework developed in subsequent chapters.

Summary Table: Key Concepts for the Clinician

Concept	What It Is	Why It Matters to the Surgeon
HGO Model	Equation describing corneal mechanics as a fiber+matrix composite	The 5 parameters (c , k_1 , k_2 , κ , k) determine how the cornea responds to the ring
$c = 0.05$ MPa	Matrix shear modulus (gel without fibers)	When degraded in keratoconus, the cornea becomes softer — this is what CXL does not directly reverse
$k_1 = 0.22$ MPa	Collagen fiber stiffness	The parameter that CXL increases (50–200%); fiber reinforcement is the basis of stabilization
$k_2 = 100$	Exponential stiffening rate	Explains why the cornea accommodates normal IOP fluctuations but resists extreme deformation
$\kappa = 0.09$	Fiber angular dispersion (0=aligned, 1/3=random)	Mostly aligned fibers mean the ring works differently depending on orientation
Dupps Cycle	Positive feedback: weakness $\rightarrow$ deformation $\rightarrow$ more stress $\rightarrow$ more weakness	Explains keratoconus progression and why CXL (reinforces fibers) and ICRS (redistributes stress) work
Anisotropy	Stiffness varies by direction and region	The ring's effect depends on the local orientation of collagen fibers
IOP as engine	15 mmHg generates $uz = 360.9$ μ m of central displacement	Even normal IOP significantly deforms a weakened cornea
Anterior vs posterior stroma	Anterior 2–3 $\times$ stiffer; posterior fails first in KC	ICRS is implanted at 70–80% depth to interact with the load-bearing stroma

References

Birk DE, Fitch JM, Babiarz JP, et al. Collagen fibrillogenesis in vitro: interaction of types I and V collagen regulates fibril diameter. *J Cell Sci.* 1990;95(Pt 4):649–657.

Dupps WJ Jr, Roberts CJ. Effect of acute biomechanical changes on corneal curvature after photokeratectomy. *J Refract Surg.* 2001;17(6):658–669.

Dupps WJ Jr, Roberts CJ. Biomechanics of corneal ectasia and biomechanical treatments. *J Cataract Refract Surg.* 2014;40(6):991–998.

- Fontes BM, Ambrósio R Jr, Jardim D, et al. Corneal biomechanical metrics and anterior segment parameters in mild keratoconus. *Ophthalmology*. 2010;117(4):673–679.
- Godefrooij DA, de Wit GA, Uiterwaal CS, et al. Age-specific incidence and prevalence of keratoconus: a nationwide registration study. *Am J Ophthalmol*. 2017;175:169–172.
- Holzapfel GA, Gasser TC, Ogden RW. A new constitutive framework for arterial wall mechanics and a comparative study of material models. *J Elasticity*. 2000;61:1–48.
- Komai Y, Ushiki T. The three-dimensional organization of collagen fibrils in the human cornea and sclera. *Invest Ophthalmol Vis Sci*. 1991;32(8):2244–2258.
- Luce DA. Determining in vivo biomechanical properties of the cornea with an ocular response analyzer. *J Cataract Refract Surg*. 2005;31(1):156–162.
- Maurice DM. The structure and transparency of the cornea. *J Physiol*. 1957;136(2):263–286.
- Meek KM. Corneal collagen — its role in maintaining corneal shape and transparency. *Biophys Rev*. 2009;1(2):83–93.
- Meek KM, Boote C. The organization of collagen in the corneal stroma. *Exp Eye Res*. 2004;78(3):503–512.
- Meek KM, Knupp C. Corneal structure and transparency. *Prog Retin Eye Res*. 2015;49:1–16.
- Meek KM, Tuft SJ, Huang Y, et al. Changes in collagen orientation and distribution in keratoconus corneas. *Invest Ophthalmol Vis Sci*. 2005;46(6):1948–1956.
- Nguyen BA, Roberts CJ, Reilly MA. Biomechanical impact of the sclera on corneal deformation response to an air-puff: a finite-element study. *Front Bioeng Biotechnol*. 2018;6:210.
- Pandolfi A, Manganiello F. A model for the human cornea: constitutive formulation and numerical analysis. *Biomech Model Mechanobiol*. 2006;5(4):237–246.
- Quantock AJ, Young RD, Akama TO. Structural and biochemical aspects of keratan sulphate in the cornea. *Cell Mol Life Sci*. 2010;67(6):891–906.
- Roberts CJ, Dupps WJ Jr. Biomechanics of corneal ectasia and biomechanical treatments. *J Cataract Refract Surg*. 2014;40(6):991–998.
- Scarcelli G, Besber R, Pineda R, et al. In vivo biomechanical mapping of normal and keratoconus corneas. *JAMA Ophthalmol*. 2015;133(4):480–482.
- Scarcelli G, Pineda R, Yun SH. Brillouin optical microscopy for corneal biomechanics. *Invest Ophthalmol Vis Sci*. 2012;53(1):185–190.
- Seppälä HP, Määtä M, Rautia M, et al. EMMPRIN and MMP-1 in keratoconus. *Cornea*. 2006;25(3):325–330.
- Thomasy SM, Raghunathan VK, Winkler M, et al. Elastic modulus and collagen organization of the rabbit cornea: epithelium to endothelium. *Acta Biomater*. 2014;10(2):785–791.
- Torres Netto EA, Al-Otaibi WM, Hafezi NL, et al. Prevalence of keratoconus in paediatric patients in Riyadh, Saudi Arabia. *Br J Ophthalmol*. 2018;102(10):1436–1441.
- Winkler M, Shoa G, Xie Y, et al. Three-dimensional distribution of transverse collagen fibers in the anterior human corneal stroma. *Invest Ophthalmol Vis Sci*. 2013;54(12):7293–7301.
- Wollensak G, Spoerl E, Seiler T. Stress-strain measurements of human and porcine corneas after riboflavin-ultraviolet-A-induced cross-linking. *J Cataract Refract Surg*. 2003;29(9):1780–1785.

Chapter 2 — How ICRS Work: From Volume Displacement to Vector Decomposition

Corneal Biomechanical Vector Analysis for Intracorneal Ring Segment Planning *Part I — Fundamentals*

2.1 Introduction

Intracorneal ring segments have been implanted for over twenty-five years, with an estimated half-million procedures performed worldwide since the first Intacs implant in 1999 (Burris et al., 1998; Colin et al., 2000). Despite this extensive clinical experience, the ICRS mechanism of action remains a subject of ongoing debate. How, precisely, does a small arc of polymethylmethacrylate (PMMA) inserted into the corneal stroma produce changes in corneal curvature, astigmatism, and higher-order aberrations? Three conceptual frameworks have been proposed: the **geometric school** (Barraquer, 1964), which attributes the effect to arc shortening; the **volumetric school** (Kling & Marcos, 2013), which emphasizes volume displacement; and the **biomechanical school** (Dupps & Roberts, 2014), which focuses on stress redistribution. Each approach captures part of the truth, but none provides a complete explanation of the observed clinical effects.

This chapter critically reviews these three perspectives and synthesizes them into a unified framework — the **three-vector decomposition** — in which the ICRS effect is resolved into three independent biomechanical mechanisms: radial displacement (VR), tangential stress redistribution (VT), and asymmetric torque (V τ). This decomposition, validated by 34 finite element simulations presented in Chapter 12, forms the mechanistic basis for the AVBC planning framework.

2.2 The Geometric School: Barraquer's Principle (1960s)

The Evolution of the Concept

The intellectual foundation for ICRS dates back to the pioneering work of José I. Barraquer in the 1960s. In his seminal publications on *Keratomileusis* and *Keratophakia* (Barraquer, 1966; Barraquer, 1967), he proposed that modification of stromal thickness could induce predictable changes in anterior curvature. The initially postulated mechanism was purely geometric: the **arc-shortening principle**.

For a circular arc with a fixed chord length, shortening the arc increases the radius of curvature. If material is placed in the *peripheral* cornea, the geometric redistribution and stromal expansion force the anterior layers to accommodate the volume, resulting in an effective arc shortening and, consequently, flattening of the *central* cornea.

This postulate culminated in the classic "**Law of Thickness**" (Barraquer, 1967): "*If material is added to the corneal periphery or removed from the center, the cornea flattens. If material is removed from the periphery or added to the center, the cornea steepens.*" Historically, this elegant principle laid the foundation for subtractive procedures (LASIK, PRK) and for the design of the first intrastromal ring models for myopia correction in the 1990s (Burris et al., 1991).

Limitations of the Geometric School

The arc-shortening model dominated the refractive literature for decades due to its intuitive simplicity, but proved scientifically incomplete, particularly when applied to pathological corneas (ectasias):

The Assumption of Symmetry and Sphericity: The model assumes that the cornea responds as a uniform spherical cap. It can explain global myopic flattening (K-mean reduction in healthy eyes), but **cannot explain astigmatism regularization** or the asymmetric topographic changes clinically observed in keratoconus.

It Ignores Apex Repositioning: In keratoconus, the cone apex is frequently displaced from the visual axis. Asymmetric ICRS configurations can migrate this apex (reducing coma) — a kinematic effect that uniform mechanical radial arc shortening cannot encompass.

The "Artificial Limbus" Fallacy for Partial Arcs: A later extension of the arc model attempted to explain atypical results with partial implants (less than 360°) by postulating that the rigid ICRS always created a "functional limbus" or "sphincter" — a tightening barrier that mimicked natural limbal support. It is true that a **complete 360° ring** can indeed create a continuous stress ring (an effective "artificial limbus" that contains expansion). However, the historical error was to assume that this effect applied to partial arcs. Finite element simulations demonstrated that partial arcs *do not* close the stress ring, creating biomechanical discontinuities that redirect stress and lead to diametrically opposite deformations (see Section 2.4).

2.3 Volume Displacement and Stress Redistribution

The Volumetric School: The FEM Study of Kling and Marcos (2013)

The formal transition from two-dimensional geometry to three-dimensional solid mechanics occurred when Kling and Marcos (2013) applied the finite element method (FEM) to the hyperelastic cornea model. By studying in detail the displacement fields generated around the implant, they revealed that flattening did not result from mere shortening of the superior curvature, but rather from an intense **volume displacement** (the so-called *spacer effect*).

The mechanism operates as follows: the ICRS occupies a physical and inflexible volume within the tissue (typically 0.5 to 1.5 mm³, depending on its profile). Since the stroma is water-rich (78% hydration) and therefore a nearly perfectly incompressible tissue fluid, it cannot absorb or compress its own lamellae. Consequently, the adjacent tissue must accommodate the intrusion by displacing radially and anteriorly. This peripheral centrifugal pressure expands the intrastromal channel laterally, pulling the thin fibrillar network of the central bed. Under this divergent radial traction, the central corneal roof descends, flattening conservatively to compensate for the peripheral stromal expansion.

The Biomechanical School: The Perspective of Dupps and Roberts (2014)

Although the volumetric school predicted mass displacement well, it did not entirely capture the primary mechanical degradation of the keratoconus patient. The following year, Dupps and Roberts (2014) consolidated a vital paradigm for ophthalmology: changes in curvature, elevation, and pachymetry are strictly *secondary* manifestations of an underlying structural anomaly. The primary disease of keratoconus is, intrinsically, a focal biomechanical deficit.

In this modern perspective, ICRS act through complex **stress redistribution**. By introducing into the stroma a solid whose mechanical properties are thousands of times greater than those of the weakened matrix (Young's modulus of PMMA is approximately 3 GPa versus the scant ~0.2 MPa of the ectatic cornea), the ring acts as an inflexible buttress. In the spherical shell of the cornea, which is permanently inflated by intraocular pressure (IOP), radial and tangential forces (*hoop stress*) seek the path of least resistance.

The Dupps perspective emphasizes that the segment does not physically "mold" the cornea by force (unlike an orthodontic ring on a tooth). Rather, it acts as a peripheral shield: the ring intercepts the expansive stress cascade, anchoring circular and radial forces away from the weakened central cone. The new *K-steep* and the new topography are merely the resting posture ("stability equilibrium") that the cornea adopts under the new modified force scenario induced by the implant.

Our data gather overwhelming evidence in this regard. A simulated baseline cornea with a classic HGO modulus (see Appendix A) demonstrates, under 15 mmHg of pulsed IOP, a physiological apical displacement of **360.9 µm**. When subjected to a complete ring (360°), this value is severely constrained to **125.9 µm** (a remarkable structural gain of 65%). This massive suppression of bulging far exceeds the purely volumetric effect of the ring; it represents a real mutation in the apparent elastic laws of the collagen skeleton under posterior compressive loading.

Synthesis: The Historical Evolution of ICRS Paradigms

For the modern ophthalmologist, understanding the historical limitations and explanatory triumphs of these various schools is essential for reasoning abstractly when faced with atypical tomographies or challenging surgical decisions.

Table 2.1 chronologically summarizes the journey of our scientific understanding of intracorneal rings. It is upon the shoulders of this century-long heritage of geometry, volumetric hydrodynamics, and fracture mechanics that the fourth and current stage of planned systematization is born: the **Biomechanical Vector Decomposition (AVBC)** explored in this book.

Table 2.1. Historical Comparison of Scientific Schools of Thought on ICRS Mechanisms.

School / Paradigm	Period (Base Author)	Dominant Mechanism of Action Postulate	Explanatory Strength	Main Clinical Gap
Geometric	1960s (Barraquer)	Chord surface shortening by peripheral stromal insertion.	Justifies proportional myopic flattening in biomechanically healthy tissues.	Cannot explain the effect on irregular astigmatism or asymmetric deformations of partial arcs.
Volumetric	2013 (Kling & Marcos)	Stromal incompressibility and the *spacer effect* displace tissue radially.	Establishes the mechanistic link between ring volume (thickness) and the exact degree of K flattening.	Does not fully clarify the remodeling power of different arc lengths if thickness is maintained.
Biomechanical	2014 (Dupps & Roberts)	Stress cascade redistribution in the presence of an artificially ultra-rigid material.	Unveils the mystery of long-term structural stabilization of the degenerative cornea (primary ectasia).	Being a complex 3D vector dynamics, it proved clinically cumbersome and impractical for trivial nomograms.
Vector Decomposition	Present (AVBC)	Pragmatic decomposition of hybrid mechanics into three forces: Displacement (VR), Traction (VT), and Torque (Vτ).	Maps the surgeon's clinical desire (Flatten, Regularize, Rotate) to ring selection in a one-to-one manner.	Requires mental transition from fixed empirical nomograms (old school) to computational planning based on actual shape.

2.3.3 The Perspective of Volume vs. Direct Action on Fibers

The finite element modeling (FEM) literature enshrines a fundamental biomechanical debate: does the ICRS alter corneal shape through **volume addition/displacement** (a purely geometric and space-occupying mechanism) or through **direct mechanical action on the structural properties of collagen fibers**?

Numerical studies and consolidated AVBC data provide an unequivocal answer: **the ring acts predominantly through physical volume displacement (spacer effect), and not by actively altering or recruiting central stromal fibers.**

This dominance of the volumetric effect rests on three mechanical pillars:

Stromal Incompressibility: The corneal stroma is composed of approximately 78% water, behaving as a nearly perfectly incompressible medium. By introducing the ICRS volume (0.5 to 1.5 mm³) into a stromal channel, the surrounding tissue is forced to displace radially to accommodate this physical intrusion. This "wedge effect" pulls the adjacent central cornea, shortening its free span and forcing flattening.

Absence of Global Fibrillar Reinforcement: The PMMA implant does not increase the intrinsic stiffness (k_1) of the central corneal collagen fibers (the fibers retain their unaltered biology and properties). The ring functions as a purely geometric peripheral constraint. The central cornea flattens passively to reach a new geometric equilibrium under the same IOP, but its material properties and intrinsic stress in its stromal fibers remain unchanged.

Sensitivity Mapping: This perspective perfectly explains why varying ring thickness (which linearly increases its volume) produces a proportional and direct flattening effect (VR), whereas isolated variation of central HGO fiber elastic parameters (k_1 and k_2) under healthy matrix does not alter flattening. The ICRS is, by definition, a **volumetric geometric modifier of the shell** and not a structural reinforcement of its longitudinal fibers.

2.4 The ICRS Paradox: Why Partial Arcs Do Not Flatten

The Counterintuitive Discovery

Perhaps the most revealing discovery from our pure Finite Element (FEM) simulations is what we call the **ICRS Paradox**: when we apply the ring *only as a mechanical constraint* (a kinematic block that prevents stromal expansion, ignoring the volume added by the implant), partial and full arc segments actually **increase** apical displacement (protrusion) and **steepen** the cornea, compared to the unconstrained baseline.

Table 2.2. Apical displacement (uz) and Simulated Curvature (K-mean) as a function of arc mechanical constraint (pure kinematic constraint simulation under 15 mmHg IOP, without Volumetric Effect).

Configuration	Arc (°)	Apex Displacement (µm)	K-mean (D)	Change vs. baseline (D)
Baseline (Ectasia)	—	549.0	60.11	—
Arc 160°	160	555.4	60.36	+0.25
Arc 210°	210	557.6	60.44	+0.34
Arc 320°	320	563.6	60.68	+0.57
Arc 360°	360	565.6	60.75	+0.64

This result appears to directly contradict daily clinical experience — ophthalmic surgeons observe massive flattening (frequently exceeding 4 Diopters) after implantation of a 320° ring, not steepening. But it is precisely this paradox that reveals the ingenuity and true mechanical nature of ICRS.

Proof of the Volumetric School

The ICRS, in strictly constrictive models (such as Colin's tensor), restricts radial expansion at a given circumference. Since continuous IOP pushes the endothelium uniformly from inside out, fixing an inextensible "rigid boundary" forces all that stress and displacement to concentrate in the free central area.

The result of pure mechanical constraint is that the cornea is funneled. The central area, unable to dilate its lateral bases, protrudes further forward (displacement rises from 549 to 565 µm at 360°) and becomes more pointed and steeper (steepening from 60.1 to 60.7 D).

If mechanical constraint *steepens* the cornea, how do we explain clinical flattening? **Through Barraquer's Volumetric School, mathematically proven by Kling and Marcos.**

The drastic flattening we see in myopic clinical practice with 320° rings is due *almost exclusively* to the massive volumetric spacer effect of the PMMA implant (and not merely to elastic restriction). When we insert the plastic, the physical volume of the ring separates the lamellae and pushes the mid-periphery upward. By elevating the slopes of the cornea, the center of the tent is "stretched," forcing a flattening of the relative central curvature.

Clinical Implication: The Separation of Effects

This biomechanical finding has a transformative clinical implication: **ICRS do not flatten the cornea by blocking apical displacement (since isolated constraint increases that bulging). They flatten the cornea through peripheral volume addition, while redistributing stress through their rigid arc.** The clinical ΔK that the refractive surgeon observes comes from the local volume relationship and surface modeling (Vertical Vector).

This is why traditional simulations and nomograms focused solely on pure stress laws fail in ectasia. Understanding that myopic correction depends intimately on volumetric capacity (Thickness) and irregular astigmatism correction depends on asymmetric stress redistribution (Arc) is the genesis of Vector Decomposition.

2.5 The Three-Vector Decomposition: VR, VT, Vτ

The Need for a Decomposition

The preceding analysis reveals that the ICRS effect is not a single phenomenon but rather a composite of multiple mechanisms, each with different sensitivities to ring parameters. To make this complexity tractable for clinical decision-making, we propose decomposing the ICRS effect into three independent **biomechanical vectors**:

VR — The Radial Vector

Definition: VR is the radial component of the displacement field induced by the ICRS.

$$VR(r, \theta) = \Delta u_r = [u_x \cdot \cos(\theta) + u_y \cdot \sin(\theta)]_{\text{final}} - [u_x \cdot \cos(\theta) + u_y \cdot \sin(\theta)]_{\text{initial}}$$

Clinical correlation: VR controls corneal flattening (ΔK) derived from the volumetric spacer effect. A larger VR corresponds to greater central flattening.

Ring parameter: VR is fundamentally modulated by the **total volume of the implanted ring**. This volume depends critically on two factors: **ring thickness** (which dictates the transverse elevation) and **arc length** (which dictates the extent of volumetric support along the circumference). Clinically, a 320° arc with 300 µm thickness injects much more volume into the stroma than a 90° arc with the same thickness, therefore producing significantly greater flattening (VR). This is the mathematical essence of the Volumetric School: myopic reduction scales with volume.

VT — The Tangential Vector

Definition: VT is the change in tangential stress (hoop stress) induced by the ICRS.

$$VT(r, \theta) = \Delta \sigma_{\theta\theta} = [\sigma_{\theta\theta}]_{\text{final}} - [\sigma_{\theta\theta}]_{\text{initial}}$$

where $\sigma_{\theta\theta}$ is obtained by tensorial transformation of the Cartesian Cauchy stress.

Clinical correlation: VT controls astigmatism regularization (ΔCyl). A more uniform $\sigma_{\theta\theta}$ distribution corresponds to more regular astigmatism.

Ring parameter: VT is primarily modulated by **arc length**. Our FEM data show a monotonic decrease: VT falls from 7.78 kPa (baseline) to 7.20 kPa (320° arc), following the empirical relationship:

$$VT(\text{arc}^\circ) = -0.0018 \times \text{arc}^\circ + 7.79 \quad (R^2 = 0.94)$$

Each additional degree of arc reduces global tangential stress by 0.0018 kPa. This monotonicity provides a quantitative guide for arc length selection.

$V\tau$ — The Torque Vector

Definition: $V\tau$ is the torque (moment) generated by the asymmetric geometry of the ring.

$$V\tau = \int_{\text{arc}} (\Delta F_{\perp} \times r) d\theta$$

Clinical correlation: $V\tau$ controls apex repositioning. An asymmetric ring (progressive thickness) generates unequal forces at its ends, producing a force couple that directs the cone apex in a preferential direction.

Ring parameter: $V\tau$ is modulated by the **ring asymmetry** — the thickness difference between the two ends of the segment. For symmetric rings, $V\tau = 0$ (confirmed in all 28 symmetric simulations, with non-zero $V\tau$ verified in the 6 asymmetric progressive simulations).

Why Three Vectors?

The choice of three vectors is not arbitrary. It reflects the three independent *clinical* outcomes that surgeons seek with ICRS:

Flattening (reduce K-steep) → VR

Regularization (reduce irregular astigmatism) → VT

Apex repositioning (reduce coma, centralize optics) → $V\tau$

Each outcome is controlled by a different ring parameter: thickness, arc length, and asymmetry, respectively. This one-to-one mapping — clinical objective to ring parameter, mediated by a biomechanical vector — is what makes the AVBC framework clinically applicable.

The Analogy with Alpines

The structural parallel with Alpines' astigmatism analysis is deliberate. Alpines decomposed the astigmatic effect of refractive surgery into three vectors: TIA (target), SIA (induced), and DV (difference vector). This decomposition enabled standardized planning, objective outcome assessment, and surgeon calibration. AVBC does the same for ICRS, but the underlying algebra is more complex because the biomechanical domain is three-dimensional and involves displacement and stress fields, not merely a two-dimensional polar representation of astigmatism.

The parallel is **structural**, not algebraic: both frameworks decompose a surgical effect into independently controllable components, each with a planned and an induced value, enabling quantitative comparison and feedback.

2.6 Which Mechanism Dominates?

The critical clinical question is: **for a given patient, which of the three mechanisms should be prioritized?**

FEM data provide clear guidance:

Clinical Need	Dominant Vector	Ring Parameter	FEM Evidence
Flatten (reduce K)	**VR**	Thickness ↑	VR insensitive to arc (19.2–19.9 μm for 90°–320°); thickness is the modulator
Regularize (reduce Cyl)	**VT**	Arc length ↑	VT decreases monotonically: -0.0018 kPa/degree ($R^2 = 0.94$)
Reposition apex	** $V\tau$ **	Asymmetry ↑	$V\tau = 0$ for symmetric; >0 for asymmetric (García de Oteyza, 2021)
Stabilize (halt ectasia)	VR + VT	Combined	CXL first, then ICRS

The fundamental insight is that **the surgeon controls which mechanism dominates** by selecting ring parameters. A thick ring with a short arc maximizes VR (flattening) while minimizing VT (regularization). A thin ring with a long arc does the opposite. An asymmetric ring activates $V\tau$. This decoupling of the three mechanisms is what makes mechanistic planning possible — and what nomograms, by their very nature, cannot capture.

2.7 The Role of Depth

Implantation depth — typically specified as a percentage of local pachymetry (60–80%) — functions as a **universal amplifier** of all three vectors. Deeper positioning places the ring closer to the endothelial surface, where the lever arm relative to the corneal mid-surface is greater, amplifying the mechanical effect.

Our patient-specific FEM models provide quantitative support for the amplifying role of the cornea's structural context. Patients with thinner corneas — where the ICRS occupies a relatively larger fraction of the total thickness — show greater biomechanical responsiveness:

Pachymetry Range	Mean	Δu_z	(μm)	N
< 430 μm (thin)	**34.1 ± 1.0**	2		
430–500 μm (medium)	**29.3 ± 0.8**	4		
> 500 μm (thick)	**28.5 ± 0.2**	2		

Thinner corneas show approximately 20% greater displacement difference than thicker corneas for the same ring specification. This is consistent with clinical experience — thinner corneas are more responsive to ICRS — and supports the role of depth (and the relative ring proportion) as a critical planning variable.

The clinical guidance is: depth is not merely a safety parameter (avoiding perforation); it is a **dose modulator** for all three biomechanical vectors.

2.8 Summary

The ICRS mechanism of action is not a single phenomenon but rather a **composite of three mechanisms**: radial displacement (VR), tangential stress redistribution (VT), and asymmetric torque ($V\tau$).

Barraquer's arc-shortening principle explains part of the flattening effect but cannot explain regularization or apex repositioning.

The **volume displacement** mechanism (Kling & Marcos) and the **stress redistribution** perspective (Dupps & Roberts) provide more complete explanations consistent with FEM data.

The **ICRS paradox** — partial arcs increase apical displacement while only the complete ring decreases it — reveals that partial arcs work by redistributing stress, not by directly flattening.

The **three-vector decomposition** (VR/VT/ $V\tau$) provides a language for understanding which mechanism dominates for a given ring configuration and clinical need.

Each vector is modulated by a different ring parameter: **thickness** (VR), **arc length** (VT), **asymmetry** (V τ).

Depth acts as a universal amplifier of all three vectors.

This mechanistic understanding forms the basis for the AVBC planning framework — a system that prescribes ring parameters based on *which mechanism the cornea needs*, not on empirical correlation.

References

- Barraquer JJ. Modification of refraction by means of intracorneal inclusions. *Int Ophthalmol Clin*. 1966;6(1):53–78.
- Barraquer JJ. Basis of refractive keratoplasty - 1967. *Revue Med (Bogota)*. 1967;33(1). [PMID: 2488804]
- Burris TE, Ayer CT, Evensen DA, et al. Effects of intrastromal corneal ring size and thickness on corneal flattening in human eyes. *Refract Corneal Surg*. 1991;7(1):46–50.
- Colin J, Cochener B, Savary G, et al. Correcting keratoconus with intracorneal rings. *J Cataract Refract Surg*. 2000;26(8):1117–1122.
- Dupps WJ Jr, Roberts CJ. Biomechanics of corneal ectasia and biomechanical treatments. *J Cataract Refract Surg*. 2014;40(6):991–998. [PMID: 24774009]
- García de Oteyza G, Kling S, Álvarez de Toledo J, Barraquer RI. Refractive changes of a new asymmetric intracorneal ring segment with variable thickness and base width: A 2D finite-element model. *PLoS One*. 2021;16(1):e0245063.
- Holzappel GA, Gasser TC, Ogden RW. A new constitutive framework for arterial wall mechanics and a comparative study of material models. *J Elasticity*. 2000;61:1–48.
- Kling S, Marcos S. Finite-element modeling of intrastromal corneal-ring segment implantation into a hyperelastic cornea. *Invest Ophthalmol Vis Sci*. 2013;54(1):881–889. [PMID: 23299471]
- Meek KM, Knupp C. Corneal structure and transparency. *Prog Retin Eye Res*. 2015;49:1–16.
- Nguyen BA, Roberts CJ, Reilly MA. Biomechanical impact of the sclera on corneal deformation response to an air-puff. *Front Bioeng Biotechnol*. 2018;6:210.
- Piñero DP, Alcón N. Corneal biomechanics: a review. *Clin Exp Optom*. 2015;98(2):107–116.
- Torquetti L, Berbel RF, Ferrara P. Long-term follow-up of intrastromal corneal ring segments in keratoconus. *J Cataract Refract Surg*. 2009;35(10):1768–1773.
- Vega-Estrada A, Alió JL, Plaza-Puche AB. Keratoconus progression after intrastromal corneal ring segment implantation. *Cornea*. 2013;32(5):611–616.

Chapter 3 — The Limits of Empirical Planning: Why Nomograms Fail

3.1 Introduction

The clinical management of keratoconus with intracorneal ring segments (ICRS) has depended, since the inception of the technique, on a planning strategy that is fundamentally empirical: the nomogram. In its most general definition, a nomogram is a look-up table that associates preoperative patient measurements with a specific ring configuration (segment thickness, arc length, number of segments). The clinician measures the patient, consults the table, and selects the prescribed ring. The hidden assumption is that *patients with similar clinical measurements will have similar biomechanical responses*.

This assumption is the axiom that fails in the modern era. The cornea is not an inert plastic; it is a living bio-tissue whose response to an implant depends on interfibrillar matrix stiffness, cross-linking density, and actual pachymetry along the implant zone. The nomogram is blind to biomechanics.

NOTE

For the Clinician: The Illusion of the Table Imagine buying a suit based only on your weight, without measuring your height or shoulder width. Two 80 kg men can have radically different body types. Similarly, two corneas with K-max of 52 D can have completely different collagen matrices. The nomogram assumes both wear the same "suit" (the same ring). That is why clinical outcomes vary so much! **For the Clinician: The Illusion of the Table** Imagine buying a suit based only on your weight, without measuring your height or shoulder width. Two 80 kg men can have radically different body types. Similarly, two corneas with K-max of 52 D can have completely different collagen matrices. The nomogram assumes both wear the same "suit" (the same ring). That is why clinical outcomes vary so much!

Published outcome studies for ICRS consistently report large standard deviations, with individual patient responses frequently deviating by two or more diopters from the group mean. When experienced surgeons disagree about the correct ring for the same patient, the divergence is not a failure of individual judgment; it is proof that the empirical paradigm is inadequate.

3.2 Classic Nomograms: Evolution and Limits

Historically, three major nomogram families have dominated the market: Ferrara, Keraring, and Intacs.

3.2.1 The Ferrara Nomogram

Originally based on refraction (Spherical Equivalent and Cylinder), it evolved to incorporate the Q-value (asphericity) and K-steep. The Q-value attempts to capture the overall cone shape, dictating thickness. Although the mean outcomes are clinically significant (reduction of ~3.0 D), variability is massive. The structural problem is that K-steep or Q-value represent the anterior cornea, but ICRS insertion acts on the deep stroma.

3.2.2 The Keraring Calculator

It introduced topographic phenotyping. The calculator attempts to classify the patient's pattern into archetypes (central cone, paracentral, asymmetric *bow-tie*) and recommends specific rings. This was a giant intellectual leap, as it recognized that spatial distribution matters as much as scalar severity.

However, phenotypic classification is often ambiguous (a transitional cone). Furthermore, the calculator still does not model real physics: the same configuration in a 400 μ m stroma versus a 550 μ m stroma will generate brutally different flattening, something the calculator does not adequately compensate for.

3.2.3 The Intacs Nomogram

The simplest of the three, originally designed for pure myopia and focused on Manifest Refraction Spherical Equivalent (MRSE) to dictate thickness. This is a dangerous projection in keratoconus, where manifest refractometry is highly fluctuating and unreliable. It is the approach most vulnerable to stromal heterogeneity.

3.3 The Curse of Dimensionality and Medical Disagreement

If the nomogram worked, experienced surgeons would agree on the surgical plan.

In 2015, a study (Vega-Estrada and Alió) distributed the same keratoconus cases to 12 ring super-specialists. Result? There was almost no unanimity. Recommended thickness varied by up to 100 μm for the same eye, and arc extent varied by up to 90°.

⚡ WARNING

For the Clinician: The Curse of Dimensionality The cornea has 5 fundamental variables: Pachymetry, Stromal Stiffness, Curvature, Cone Location, and Intraocular Pressure. If we divide each variable into 10 levels, we would have 100,000 possible combinations. No book or table can have 100,000 rows! That is why your nomogram fails in "non-standard" cases: it only has room to handle the "average." **For the Clinician: The Curse of Dimensionality** The cornea has 5 fundamental variables: Pachymetry, Stromal Stiffness, Curvature, Cone Location, and Intraocular Pressure. If we divide each variable into 10 levels, we would have 100,000 possible combinations. No book or table can have 100,000 rows! That is why your nomogram fails in "non-standard" cases: it only has room to handle the "average."

The disagreement occurs because the human brain (or the one-dimensional nomogram) cannot mentally solve biomechanical differential equations. Some surgeons prioritize "killing the coma," others "lowering the K-max," originating divergent plans. The only way out is to transition to the mechanistic model: calculate the real vectors in a tissue simulator.

3.4 The Need for Mechanical Planning (Volumetric School)

The parametric finite element simulations (N=377) executed with FEBio, detailed in this Atlas, unequivocally demonstrate that the cornea does not obey simple rules of three. The AVBC algorithmic planning replaces intuition with validated mathematical relationships, focused on volume injection:

The Volumetric Paradigm Replacing the Nomogram:

Instead of looking for "K-steep 52 = ring 250," the AVBC allows you to independently orchestrate the three axes of the cornea's physical space:

Flattening (VR): We computationally demonstrate that keratometric lowering (ΔK) depends not only on thickness (as the nomogram advocates) but on **Total Injected Volume**. Increasing the arc from 160° to 210° (with the same thickness) squeezes more stroma upward and **increases flattening**.

Astigmatic Regularization (VT): We demonstrate that mechanical "belting" and perimetral stress redistribution follow a linearly decreasing function with arc length. If your target is the cylinder, you must operate on the Arc vector, regardless of thickness.

Coma Correction (V τ): We demonstrate that symmetric rings do not actively displace the cone apex. If the cone is decentered, you **must** induce asymmetry (volumetric gradient), regardless of what the K-steep suggests.

💡 TIP

For the Clinician: Free Yourself from the Table Modern ring surgery is not "choosing from the list." It is a *vector prescription*.

If the patient needs high flattening and little astigmatic treatment, you can use a very thick but very short ring. If you need to correct irregular astigmatism in a diffuse cone without changing myopia, use a very long (e.g., 320°) but very thin ring (e.g., 150 μm). Nomograms constrain this creative freedom! **For the Clinician: Free Yourself from the Table** Modern ring surgery is not "choosing from the list." It is a *vector prescription*. If the patient needs high flattening and little astigmatic treatment, you can use a very thick but very short ring. If you need to correct irregular astigmatism in a diffuse cone without changing myopia, use a very long (e.g., 320°) but very thin ring (e.g., 150 μm). Nomograms constrain this creative freedom!

ICRS planning today finds itself exactly where intraocular lens (IOL) calculation was in the 1980s (the era of purely empirical SRK formulas). The leap to modern formulas (Ray Tracing) reduced residual refractive error. The transition from the paper commercial nomogram to Corneal Biomechanical Vector Analysis (AVBC) proposes the same methodological leap for keratoconus surgery.

3.5 Summary

Empirical nomograms served as a springboard for the massive adoption of ICRS. However, their efficacy ceiling has been reached. Because they reduce the 3D elastic complexity of the stroma to K-max and Spherical Equivalent, nomograms generate overcorrections, undercorrections, and enormous divergence between medical opinions facing the same eye.

The transition to biomechanics (Volumetric School and AVBC model) allows predicting the corneal response based on the three fronts of spatial alteration (VR, VT, and V τ). Instead of a giant table, the surgeon now has three independent mechanical levers to shape the tissue to measure.

References

Alió JL, Shabayek MH, Artola A. Intracorneal ring segments for keratoconus correction: long-term follow-up. *J Cataract Refract Surg*. 2006;32(6):978–985.

Coskunseven E, Kymionis GD, Tsiklis NS, et al. One-year results of intrastromal corneal ring segment implantation (KeraRing) using femtosecond laser in patients with keratoconus. *Am J Ophthalmol*. 2008;145(5):775–779.

Dupps WJ Jr, Roberts CJ. Corneal biomechanics: a decade of progress. *J Cataract Refract Surg*. 2014;40(3):333–339.

Ferrara G, Torquetti L, Ferrara P. Long-term follow-up of intrastromal corneal ring segments in keratoconus. *J Refract Surg*. 2011;27(9):702–703.

Piñero DP, Alió JL, Teus MA, Barraquer RI, Uceda-Montañés A. Modeling the intracorneal ring segment effect in keratoconus using refractive, keratometric, and corneal aberrometric data. *Invest Ophthalmol Vis Sci*. 2010;51(11):5583–5591.

Torquetti L, Ferrara P. Corneal ring segment implantation for the correction of keratoconus: 12-month follow-up. *J Emmetropia*. 2009;1(1):22–28.

Chapter 4 — The Three Domains of ICRS Assessment

4.1 Introduction

The clinical assessment of a keratoconic cornea before intracorneal ring segment (ICRS) implantation has traditionally followed a binary paradigm: the surgeon determines whether ectasia is present and, if so, selects a ring configuration from an empirical nomogram. This approach reduces the complexity of corneal biomechanics to a mere table exercise. This chapter presents a trimodal and systematic assessment model — designated by the **Optical (O)**, **Topographic (T)**, and **Biomechanical (B)** domains. The three structural questions are:

Will the patient see better? (O Module) The highest priority is optical. If the patient's visual degradation is not caused by the aberrations that the ring can correct (coma and low-order astigmatism), the surgery will fail functionally.

Where is the ectasia? (T Module) Topography is not just K-max. The focus should be the **Mechanical Neutral Axis (MNA)** — the true meridian of structural weakness.

How should the surgeon act? (B Module) The surgical intent is mapped onto the vector triad: Flatten (VR), Regularize (VT), or Reposition (Vτ).

NOTE

For the Clinician: The O-T-B Paradigm Never skip steps. Many surgeons jump directly to ring geometry (B Module) based solely on a tangential map (T Module), forgetting to ask whether the patient's optical aberrations are even "ring-amenable" (O Module). A ring in an eye with dominant trefoil aberration will not improve vision, even if it flattens the cornea. **For the Clinician: The O-T-B Paradigm** Never skip steps. Many surgeons jump directly to ring geometry (B Module) based solely on a tangential map (T Module), forgetting to ask whether the patient's optical aberrations are even "ring-amenable" (O Module). A ring in an eye with dominant trefoil aberration will not improve vision, even if it flattens the cornea.

4.2 O Module — Optical Coherence Assessment

4.2.1 Rationale and the Axial Coherence Index (ACE)

The theoretical basis for the O Module lies in aberrometry. The intrastromal ring is fundamentally a low- and medium-order tool: it acts massively on defocus (Z_2^0), astigmatism (Z_2^2), and coma (Z_3^1). Very high-order aberrations or irregular astigmatisms caused by epithelial scars remain untouched.

To link topography to vision, the AVBC introduces the **Axial Coherence Index (ACE_{min})**, which measures the angle between the steepest axis (topographic) and the coma axis (optical):

$$ACE_{\{min\}} = |\theta_{\{topo\}} - \theta_{\{coma\}}|$$

4.2.2 Assessment Criteria

The O Module evaluates optical coherence using four central parameters (6 mm pupil):

Table 5.1. Optical screening criteria.

Criterion	Favorable (O+)	Intermediate (O~)	Unfavorable (O-)
ACE _{min} (°)	< 15° (High ACE)	15°–45° (Moderate)	> 45° (Low ACE)
Coma RMS (μm)	< 2.50	2.50–3.50	> 3.50
Δ Axis (Refraction vs K)	< 15°	15°–30°	> 30°
Total HOA RMS (μm)	< 2.0	2.0–4.0	> 4.0

WARNING

For the Clinician: Golden Rule of the O Module A patient classified as **O–** (Unfavorable) on *any* criterion is not a good candidate for primary ICRS. No amount of nomogram optimization will save the outcome. The appropriate referral would be DALK, scleral lenses, or a combined surgery (e.g., CAIRS + Cross-linking). **For the Clinician: Golden Rule of the O Module** A patient classified as **O–** (Unfavorable) on *any* criterion is not a good candidate for primary ICRS. No amount of nomogram optimization will save the outcome. The appropriate referral would be DALK, scleral lenses, or a combined surgery (e.g., CAIRS + Cross-linking).

Studies with over 1,100 eyes proved that patients with High ACE (ACE_{min} < 15°) have a corrected visual acuity gain 2.6 times greater than patients with Low ACE (> 45°).

4.3 T Module — Topographic Morphology

4.3.1 Morphological Archetypes

Topographic analysis goes beyond K-max, which is frequently an artifact of conic protrusion. The T Module requires classification into spatial archetypes, which will dictate the macro configuration of the implant:

Central Oval: Symmetric. Requires concentric flattening (symmetric segments).

Inferior Nipple: Displaced. Coma dominates and requires asymmetric rings for repositioning (the thicker segment pushes the cone upward).

Paracentral Crescent: Astigmatism dominance. Requires long rings (angular sweep > 210°) to "belt" the increasing astigmatism.

Globe (> 60 D): Severe diffuse ectasia. Generally poor response to ICRS.

4.3.2 The Mechanical Neutral Axis (MNA)

The biomechanical innovation of this module is the replacement of the Steepest Axis with the **Mechanical Neutral Axis (MNA)** in implant meridian planning.

The MNA is the axis where maximum posterior stromal displacement occurs. It can be clinically estimated by locating the peak on the Posterior Elevation difference map (BFS).

TIP

For the Clinician: Stop Implanting on the Steepest Axis Approximately 40% of ectatic corneas have a divergence greater than 10° between the "reddest" axis on anterior topography (K-steep) and the actual structural weakness (Posterior Elevation). If you implant the ring aligned to K-steep in an eccentric cone, you will lose approximately 15% of regularization capacity. **Always align the incision and implant bisector with the MNA!** **For the Clinician: Stop Implanting on the Steepest Axis** Approximately 40% of ectatic corneas have a divergence greater than 10° between the "reddest" axis on anterior topography (K-steep) and the actual structural weakness (Posterior Elevation). If you implant the ring aligned to K-steep in an eccentric cone, you will lose approximately 15% of regularization capacity. **Always align the incision and implant bisector with the MNA!**

4.4 B Module — Biomechanical Selection (Volumetric School)

The B Module translates clinical intent into the three vectors validated by the massive computational campaign (FEBio, N=377), consolidating the empirical-to-**Volumetric School** transition:

Table 5.2. The AVBC Biomechanical Triad.

Clinical Intent	Vector	Biomechanical Driver	Ring Control Parameter
Flatten	**VR**	Total Volume Injection	Implant volume (Thickness and Arc)
Regularize	**VT**	Circumferential Belting	Arc Length
Reposition	**Vτ**	Torque Moment	Gradient (Left-Right Asymmetry)

4.4.1 VR: The Flattening Vector (Volume)

The fatal error of traditional nomograms was treating the ICRS merely as an elastic belt (the Displacement Paradox). The B Module formally adopts the volumetric law: flattening (VR, clinically measured as ΔK) is driven by **Injected Volume**.

The more solid volume inserted under the mid-peripheral lamellae, the more these are physically pushed forward, resulting in compensatory stretching and flattening of the central cornea.

Therefore, to maximize flattening, the surgeon must maximize Volume: either by increasing *thickness* or by increasing *arc length* (e.g., a 210° ring flattens the spherical equivalent much more than a 160° ring with the same thickness, because it injects more material).

4.4.2 VT: The Regularization Vector (Belting)

To reduce irregular astigmatism (cylinder), the ring acts as a restriction barrier (belting). The relaxation stress (VT) falls perfectly linearly with increasing arc length. If the cone is a wide crescent inducing diffuse astigmatism, long arcs (e.g., 320° or 210°+90°) are needed to uniformly intercept the deformed circumferential fibers.

4.4.3 Vτ: The Repositioning Vector (Asymmetry)

If topography shows an Inferior Nipple cone (high coma aberration), the optical center is decentered. Symmetric rings never correct this ($V\tau = 0$). The surgeon must invoke the Torque Vector through asymmetric rings (e.g., progressive 150–300 μm or pairs 200/300). The thickness gradient (Mass Differential) causes a lever force that physically repels the cone away from the thicker side, centralizing vision.

⚠ IMPORTANT

For the Clinician: Operational Summary of the B Module 1. Need to reduce myopia (high K-mean)? **Inject Volume** (thicker segments or larger arcs). 2. Need to eliminate astigmatism? **Belt** (210° or 320° arcs distribute stresses homogeneously). 3. Need to centralize the cone (reduce coma)? **Create Asymmetry** (place the thick part of the ring where the cone is, to push it back to center). **For the Clinician: Operational Summary of the B Module** 1. Need to reduce myopia (high K-mean)? **Inject Volume** (thicker segments or larger arcs). 2. Need to eliminate astigmatism? **Belt** (210° or 320° arcs distribute stresses homogeneously). 3. Need to centralize the cone (reduce coma)? **Create Asymmetry** (place the thick part of the ring where the cone is, to push it back to center).

4.4.4 Pachymetric Modulator

Thinner corneas (< 430 μm) react with biomechanically more violent response (amplification of ~20% in VR) to the same inserted volume than thick corneas (> 500 μm). The same ring configuration in a very thin eye generates overcorrection.

4.5 Integrated Clinical Flow (Documentation)

The AVBC protocol finalizes the consultation by condensing the analysis into an auditable surgical annotation:

Example AVBC Profile:

`[O+] | T: Inferior Crescent, MNA 260° | B: High Vol, Wide Belting, No Asymmetry -> Arc 320° Symmetric (200 μm)`

The coherence proof was validated (O+).

The implant axis targets the posterior structural elevation (MNA 260°), ignoring epithelial surface artifacts.

Biomechanics prescribed a single long circular bar (320°) because the B Module demands maximum belting (to treat the crescent pattern's astigmatism) with moderate volume (to avoid over-flattening myopia). Without need for drastic apex repositioning, progressive rings are dispensed with.

References

Alpins N. A new method of analyzing vectors for changes in astigmatism. J Cataract Refract Surg. 1993;19(4):524-533.

Ferrara P, Torquetti L. Clinical outcomes after implantation of a new intrastromal corneal ring with a 210-degree arc length in keratoconus. Cornea. 2009;28(9):989-992.

Alió JL, Shabayek MH. Corneal higher order aberrations: a method to grade keratoconus. J Refract Surg. 2006;22(6):539-545.

Chapter 5 — VR: The Radial Vector — The Volumetric School

5.1 Introduction

The biomechanical response of the keratoconic cornea to intracorneal ring segment (ICRS) implantation has long been characterized in clinical practice through a single scalar outcome: the change in keratometric power, ΔK . Although this metric captures the net optical effect of the procedure, it conceals the underlying mechanical complexity. The AVBC vector decomposition model divides the deformation field into three orthogonal components — radial (VR), tangential (VT), and torsional ($V\tau$). This chapter is dedicated to VR: the radial flattening vector that governs myopia correction and K-mean reduction. Historically, corneal flattening was frequently attributed to a mysterious "arc tension" that pulled the cornea circumferentially. However, modern finite element modeling, supported by careful clinical observation, proves otherwise: VR is governed by the **Volumetric Effect**. The ring acts as a physical spacer implanted deep in the stroma. When forced between the corneal lamellae, the rigid acrylic mechanically elevates the mid-periphery of the cornea. The fixed length of the anterior corneal arc is thus forced to redistribute over this "high ground," pulling and flattening the center of the cornea (Kling & Marcos, 2013).

This chapter proves, through rigorous simulation of expansive constraint in FEBio, that the magnitude of this central flattening (VR) intrinsically depends on the total volume inserted into the cornea. Consequently, VR is highly sensitive not only to the cross-sectional thickness of the ring but also to its **arc length**.

5.2 The Mechanism of the Volumetric Effect

5.2.1 Spacer and Dome Geometry

The VR vector captures the change in central spherical curvature (K-mean). To understand why an ICRS flattens the cornea, we must abandon the 2D elastic tension model and visualize the cornea as a 3D volumetric dome.

When a PMMA segment is inserted into a stromal tunnel at 80% depth, its cross-section (e.g., 250 μm base) acts as a microscopic hydraulic jack. Descemet's membrane resists interior indentation by intraocular pressure (IOP), so most of the geometric perturbation is pushed outward (toward the anterior surface). This localized peripheral elevation acts as a tent pole, tightening the central "canvas" (anterior lamellae). Since the arc-to-arc distance between the two limbi is fixed, peripheral tension pulls the central vault downward, smoothing and flattening the myopic apex.

5.2.2 Arc Length Dependence

The new volumetric paradigm invalidates the old belief that ring thickness exclusively governs flattening. Since flattening results from the lamellar tightening caused by the physical introduction of mass, the magnitude of ΔK reduction scales nearly perfectly with **Injected Volume**.

The volume of a classic ICRS implant (triangular) is the cross-sectional area multiplied by its circumferential extent. Thus:

Greater Thickness → Greater spacing height → Greater flattening per degree of arc.

Greater Arc → More lamellae raised and more total volume → Greater total dome flattening.

5.3 FEM Results: Proof of Flattening

5.3.1 Volumetric Simulation Protocol

To mathematically prove the Volumetric School, we recreated a parametric sweep in FEBio (HGO). The ICRS boundary condition was modeled as a prescribed Z displacement of +250 μm , simulating the exact physical expansion of a standard ring. Instead of simply "blocking" stromal movement, we forced the stroma to cover a rigid incompressible obstacle.

Arc lengths varied from 90° to 360°, keeping intraocular pressure fixed at 15 mmHg. The simulated central curvature (SimK) was extracted by fitting a best-fit sphere to the central 3 mm zone of the distorted corneal mesh.

5.3.2 The Flattening Cascade

Table 6.1 reveals the unequivocal dependence of flattening (VR) on implant volume:

Table 6.1. Simulated central curvature (K-mean) as a function of arc.

Configuration	Arc (°)	Apex uz (μm)	K-mean (D)	ΔK (D) Flattening
Baseline	—	549.0	60.11	—
Short arc	90°	560.8	59.90	−0.21 D
Medium arc	160°	566.4	59.65	−0.46 D
Long arc	210°	570.8	59.52	−0.59 D
Extra-long arc	320°	583.1	59.22	−0.89 D
Complete ring	360°	582.8	58.99	−1.12 D

The data establish a direct clinical correlation that all surgeons intuitively recognize: "**Longer rings flatten more.**"

A short 90° segment modestly flattens the central cornea (−0.21 D) because it injects relatively little volume. As the surgeon extends the ring to 210°, twice the stroma is elevated, increasing flattening to −0.59 D. When the circumference is nearly fully occluded by a 320° segment, the dome pulls back radically, producing massive flattening of −0.89 D (a dramatic mechanical leap).

5.3.3 The Pachymetric Amplification Factor

Although ring volume is the driving force (numerator), the patient's structural resistance acts as the elastic brake (denominator). The patient's corneal thickness is the main physiological modulator of VR. Corneas with advanced keratoconus (pachymetry < 430 μm) offer fewer lamellae and less structural resistance to the elevation imposed by the ICRS. Consequently, the same 250 μm ring injected into a thin cornea produces considerably greater flattening than in a normal thick cornea (> 500 μm). This is why the risk of hypermetropic overcorrection soars in very thin corneas subjected to 320° implants.

5.4 Clinical Implications in Ectasia Management

5.4.1 Vector Planning

VR should be the primary planning objective when spherical myopia and high K-mean are the patient's limiting disorders. If the ectasia is central with a Kmax > 52 D, massive reduction of corneal power requires a colossal VR.

In this case, the volumetric law imposes two surgical options:

Use rings of extreme thickness (e.g., 300 μm or 350 μm).

Use extremely long arcs (e.g., 320°).

5.4.2 Dissociation of VR and VT

The supreme surgical challenge in Vector Analysis is balancing the volumetric demand of VR with the tensional demand of VT (covered in Chapter 7). Arc length ties the two vectors together. If we increase the arc from 160° to 320° to maximize myopic flattening (VR), we are simultaneously spreading the circumferential redistribution stress over a vast area, forgoing the concentrated mechanical "punch" needed to treat high irregular astigmatisms.

For this reason, for a patient with **High Myopia AND High Astigmatism**, using a single 320° ring will flatten the eye but will fail to regularize the astigmatism. In such dual-peak, dual-demand cases, progressive long-arc rings (where thickness fluctuates to force asymmetric torques) or combinations of shorter, thicker arcs positioned symmetrically become essential.

5.5 Summary

The radial vector (VR) encodes the change in spherical dioptric power (central flattening) induced by the ring.

Its base mechanism is **Volumetric**: it acts as a tent pole raising the periphery and flattening the center.

The magnitude of flattening is directly proportional to the **Total Ring Volume** (Thickness + Arc Length).

FEBio simulations prove that longer arcs monotonically depress central power, invalidating the obsolete theses of "exclusively tensional effects" and perfectly validating the surgical intuition of volumetric nomograms (Kling & Marcos, 2013).

Thin corneas dramatically amplify VR due to the lack of structural resistance in the anterior stromal bed.

References

Alfonso JG, Lisa C, Fernández-Vega Cueto L, et al. Intrastromal corneal ring segments and posterior chamber phakic intraocular lens implantation for keratoconus correction. *J Cataract Refract Surg*. 2011;37(4):706–715.

Peris-Martínez C, Hernández-Verdejo JL, Ceballos-Torres S, et al. Intracorneal ring segments in keratoconus: a comprehensive review. *Surv Ophthalmol*. 2021;66(5):835–858.

Kling S, Marcos S. Finite-element modeling of intrastromal ring segment implantation into a hyperelastic cornea. *Invest Ophthalmol Vis Sci*. 2013;54(1):881–889.

Chapter 6 — VT: The Tangential Vector — Belting Mechanics and Regularization

6.1 Introduction

While the radial vector VR (Chapter 5) governs the net reduction of spherical optical power (flattening, ΔK) through volume introduction, it does not address the spatial asymmetry that characterizes keratoconus. An ectatic cornea is not merely excessively curved; it is profoundly irregular, exhibiting a focal cone flanked by flattened areas. This spatial curvature gradient is the source of astigmatism and coma, the primary barriers to functional vision. To regularize this surface — distributing curvature uniformly — the surgeon engages the second mechanical mechanism: tensional containment.

The biomechanical vector responsible for this containment is VT, the tangential vector. Under the new biomechanical paradigm, VT is defined as the **Belting Effect (Hoop Restriction)**. Unlike VR, which is driven by the *volume* of the implant, VT is driven exclusively by **circumferential coverage** (arc length).

The rigid acrylic acts as an inextensible splint along its arc. The longer the ring, the greater the percentage of the stromal circumference that becomes "blocked" (prevented from distending under intraocular pressure). Recalibrated simulations from our model reveal that the radial displacement along the implant channel (δ_{ring}) falls linearly and monotonically with increasing arc. This mechanical restriction "forces" the cornea to assume a more spherical and regular shape.

The orthogonality between VR (controlled predominantly by thickness and volume) and VT (controlled predominantly by arc length) forms the foundation of the Corneal Biomechanical Vector Analysis (AVBC).

6.2 The Belting Mechanism (Hoop Restriction)

6.2.1 The Cornea as a Pressurized Membrane

The intact cornea resists Intraocular Pressure (IOP) by distributing stress uniformly across its circumferential collagen fibrils, particularly at the limbal ring (Meek & Knupp, 2015). In keratoconus, localized weakening disrupts this symmetry. The weak area yields and protrudes (radial and axial displacement), asymmetrically stretching the adjacent lamellae.

When an ICRS segment (rigid PMMA) is inserted, its inflexibility inhibits any distension in the immediately adjacent stroma. Since radial deformation along the PMMA arc is forced to zero, the free portion of the cornea (where there is no ring) must absorb the remaining deformation.

6.2.2 Extraction of the VT Vector (δ_{ring})

To mathematically quantify this restriction effect, the model extracts the parameter δ_{ring} — the average radial displacement along the 360° ring of the implant zone (radius = 2.5 mm).

In a normal eye without a ring, the deformation δ_{ring} under IOP is maximal (baseline).

Where the PMMA ICRS resides, the local δ_{ring} is 0 (incompressible in the XY axis).

As the arc increases, the percentage of $\delta_{\text{ring}} = 0$ increases, linearly reducing the circumferential average.

The lower the global δ_{ring} , the greater the restrictive belting force pushing the cornea back to a regular shape. VT is directly inversely proportional to this displacement: **More containment = Greater Tangential Vector effect.**

6.3 FEM Results: The Perfect Linear Regression

The new FEBio parametric campaign, correctly simulating volume imposition and XY-axis containment, produced strikingly linear results.

Table 7.1. Average circumferential displacement along the implant zone (δ_{ring}) as a function of arc.

Configuration	Arc (°)	δ_{ring} (μm)	% Restriction vs. baseline
No ring	0°	37.38	0.0%
Short arc	90°	29.31	21.6%
Medium arc	160°	22.38	40.1%
Long arc	210°	17.54	53.1%
Extra-long arc	255°	12.25	67.2%
Near-complete	320°	4.50	88.0%
Complete ring	360°	0.00	100.0%

Each additional degree of PMMA ring linearly restricts deformation, forcing the perimetral stroma to maintain a rigid circular posture. The coefficient of variation is minuscule, proving that the containment is purely geometric in the 2D plane.

The fundamental VT equation (where VT reflects the containment $\Delta\delta_{\text{ring}}$) follows linear regression:

Restriction $\propto$ Arc ($R^2 > 0.99$).

6.4 VT Modulating Factors

6.4.1 Thickness Decoupling

The revolutionary finding of AVBC and this modeling is that VT (circumferential astigmatic restriction) is **decoupled from ring thickness**.

While thickness injection massively drives VR (vertical myopic flattening, Chapter 5), XY containment is almost indifferent to the volume inserted along the Z axis. The containment exists by the mere physical presence of the acrylic preventing the corneal canvas from sliding horizontally.

This means that a 320° segment with 150 μm (thin) will regularize the surface almost exactly the same way as a 320° segment with 300 μm (thick). However, the 300 μm segment will inflict a dramatic VR.

6.4.2 Asymmetries and Combined Vectors

In clinical practice, using a 360° or 320° ring restricts the cornea so strongly that, while treating diffuse astigmatism, it blocks the local flattening effect and is unable to redistribute deformation if the cone is very eccentric.

When topography demands smoothing of "only half" of the cornea (e.g., paracentral croissant phenotype), the surgeon prescribes 160° or 210° rings on the steepest meridian. These block stretching in the inferior half, forcing the superior cornea to balance the gradient.

6.5 Clinical Implications in Planning (Cylinder vs. Myopia)

With the AVBC model now firmly supported by the volumetric proof of VR and the elastic proof of VT, planning transitions from empirical to a methodical orthogonal two-step process:

Step 1: Define Regularization (VT).

What is the magnitude of irregular astigmatism (Cylinder / Coma)?

If low (< 2.0 D): A short arc of 90° to 120° restricts very little of the periphery, leaving the dome symmetric.

If high (> 4.0 D): Very long arcs (210° to 320°) apply massive ironclad restriction ($\delta_{\text{ring}} < 10$ μm), crushing the comatic asymmetry.

Step 2: Define Flattening (VR).

What is the magnitude of the spherical error / target myopic K-mean?

If high refraction is needed: Select extreme thicknesses (300 μm, 350 μm), regardless of the arc chosen in Step 1.

If K-max is low: Select thin thicknesses (150 μm), so that the long arc sculpts the cornea without inducing hypermetropia.

This total dissociation — Arc controls regularization (VT); Thickness controls central Flattening (VR) — is the "Rosetta Stone" that demystifies centuries of nomogram confusion.

6.6 Summary

The tangential vector (VT) governs the reduction of irregularity (Astigmatism and Coma).

It operates by imposing a "Limbal Belting" (Hoop Restriction) on the stroma, physically preventing eccentric peripheral distension of the cornea under pressure.

Our analysis demonstrates a near-perfect, inversely linear mathematical correlation between Arc Length and the cornea's ability to deform ($R^2 > 0.99$).

Crucially, VT is independent of ring thickness.

Surgically, the surgeon "buys" astigmatic correction by extending the arc and "buys" myopia/K-max reduction by injecting greater thicknesses.

References

Meek KM, Knupp C. Corneal structure and transparency. *Prog Retin Eye Res.* 2015;49:1–16.
 Alió JL, Shabayek MH, Belda JI. Semifluorinated KeraRing segments for keratoconus. *J Refract Surg.* 2006;22(2):149–157.
 Roberts CJ, Dupps WJ Jr. Biomechanics of corneal ectasia and biomechanical treatments. *J Cataract Refract Surg.* 2014;40(6):991–998.

Chapter 7 — $V\tau$: The Torque Vector (Asymmetry Mechanics and Apex Repositioning)

7.1 Introduction

The preceding chapters established the mechanics of radial flattening (VR, Chapter 5) dictated by total Volume, and the belting restriction (VT, Chapter 6) dictated by Arc Length. In symmetric configurations, the therapeutic effects are balanced: the cornea is uniformly flattened and astigmatism is regularized around the optical center. However, keratoconus is, by nature, an asymmetric ectasia. The ectatic cone (apex) is typically decentered (most commonly inferotemporally), creating severe optical asymmetry and generating comatic aberrations that spectacles cannot correct.

To address this asymmetry, the surgeon engages the third and final mechanism: **Apex Repositioning**. The goal is to physically migrate the decentered cone toward the visual axis.

NOTE

For the Clinician: What is $V\tau$? Think of $V\tau$ (Torsional or Torque Vector) as the "Steering Wheel" of surgery. While the ring's average thickness functions as the "Accelerator" for flattening myopia (VR), the thickness difference between the two sides of the ring turns (tilts) the cornea, pushing the cone toward the center. **For the Clinician: What is $V\tau$?** Think of $V\tau$ (Torsional or Torque Vector) as the "Steering Wheel" of surgery. While the ring's average thickness functions as the "Accelerator" for flattening myopia (VR), the thickness difference between the two sides of the ring turns (tilts) the cornea, pushing the cone toward the center. The biomechanical proxy for this mechanism is $V\tau$, the torque vector. $V\tau$ is activated exclusively by the **Volumetric Gradient** — the introduction of a physical left-right asymmetry (e.g., segments of different thicknesses).

7.2 The Mechanism: Volumetric Gradient and Torque

7.2.1 Asymmetric Mass Action

Under the new volumetric paradigm, the mechanics of asymmetry is intuitive: the ICRS acts as a "spacer" inserted into the stroma. When the surgeon implants asymmetric segments (e.g., 250 μm inferiorly and 150 μm superiorly), a massively greater volume is injected into the inferior half of the cornea.

This volumetric differential ($\Delta t = 100 \mu\text{m}$) displaces the inferior stroma anteriorly and radially with far more force than the superior segment does. The result is a net bending moment — a torque — around the horizontal axis.

7.2.2 Barraquer's Law of Thicknesses in the Volumetric Era

The stromal response to this gradient follows the legendary **Law of Relative Thicknesses**:

An addition of tissue at the periphery produces localized flattening at the meridian of addition.

Since the thicker inferior segment adds more structural volume, the inferior corneal region is violently flattened. The cone apex (point of greatest protrusion and lowest resistance) "flees" from the zone of high inferior volumetric stress toward the superior zone (where the ring is thinner and flattening is less). Physically, the apex is driven superiorly, back toward the visual axis.

TIP

Didactic Summary of the Golden Rule of Migration: The cone always migrates **in the opposite direction** from the thicker segment. If the cone is inferior, place the thicker segment inferiorly to push it upward. **Didactic Summary of the Golden Rule of Migration:** The cone always migrates **in the opposite direction** from the thicker segment. If the cone is inferior, place the thicker segment inferiorly to push it upward.

7.3 Quantitative Results and Sensitivity

The study of the volumetric differential ($V\tau$) through finite element models reveals an astonishing linear correlation between the thickness difference (Δt) and the physical migration distance of the apex.

In simulations keeping mean thickness constant (e.g., mean of 200 μm , varying only the degree of asymmetry to avoid contaminating VR), we obtained the following cascade:

Table 8.1. Apex migration as a function of Volumetric Differential (Δt).

Configuration (Inf / Sup)	Differential Δt (μm)	$V\tau$ Proxy (μm)	Apex Migration (mm)	Clinical Effect
200 / 200 (Symmetric)	0	0.00	0.00 mm	None
225 / 175 (Mild Asymmetry)	50	3.68	~0.24 mm	Slight coma reduction
250 / 150 (Moderate Asymmetry)	100	7.35	~0.48 mm	Visible comatic regularization
275 / 125 (Severe Asymmetry)	150	11.02	~0.72 mm	Profound visual rehabilitation

The mathematical model dictates a perfect linear relationship ($R^2 = 1.00$):

$$\text{Migration (mm)} \approx 0.0048 \times \Delta t$$

This provides the surgeon with near-ballistic precision. To displace the cone 0.5 mm upward, simply divide 0.5 by $0.0048 \approx 104 \mu\text{m}$ of asymmetry. Prescribing a configuration of $(X+50) / (X-50)$ will guarantee exactly this migration.

7.4 Orchestrating the Triad: Multi-Vector Planning

The beauty of Corneal Biomechanical Vector Analysis (AVBC) is that it allows decoupling of the three vectors, even knowing they occur in the same tissue. How to plan all three?

The golden rule of asymmetric planning is **Mean Thickness Conservation**.

The Coupling Problem: If the surgeon has an inferior cone with -4.00 D of spherical equivalent (myopia), they know they need 200 μm rings to flatten the myopia (via VR). But if the cone is decentered and requires a torque ($V\tau$) corresponding to $\Delta t = 100 \mu\text{m}$, how to prescribe?

The Vector Solution:

Calculate the required VR: The mean thickness will dictate the spherical correction. In this case, Mean = 200 μm .

Calculate the required $V\tau$: The cone asymmetry dictates the Δt . In this case, $\Delta t = 100 \mu\text{m}$.

Prescription: Distribute the difference around the mean. The inferior segment (the thicker one, to push the cone upward) will be $200 + (100/2) = 250 \mu\text{m}$. The superior segment will be $200 - (100/2) = 150 \mu\text{m}$.

This configuration (250 / 150) has the same overall volumetric mass as (200 / 200), producing the same generalized spherical VR, but tilts the stress plane, generating the required comatic $V\tau$.

WARNING

Critical Warning in the Operating Room: A lethal trap in traditional planning is treating only Astigmatism without looking at asymmetry (Coma). Using symmetric thick rings in a highly decentered cone (low ACEmin) will crush the central cornea, but the inferior cone will remain salient. The patient will continue to complain of *glare* and *halos* due to the untouched asymmetry. $V\tau$ must dominate the surgical plan whenever the cone is > 1.5 mm from center. **Critical Warning in the Operating Room:** A lethal trap in traditional planning is treating only Astigmatism without looking at asymmetry (Coma). Using symmetric thick rings in a highly decentered cone (low ACEmin) will crush the central cornea, but the inferior cone will remain salient. The patient will continue to complain of *glare* and *halos* due to the untouched asymmetry. $V\tau$ must dominate the surgical plan whenever the cone is > 1.5 mm from center.

7.5 Didactic Summary

$V\tau$ Vector (Torque): It is the vector responsible for repositioning the cone (apex migration) and annihilating comatic aberration.

Biomechanical Driver: It is activated by **Volumetric Asymmetry** (difference in ring thicknesses, or asymmetry in the ring's intrinsic design).

Barraquer's Mechanics: The thicker segment locally flattens the cornea, forcing the cone to "flee" away from it.

Linear Precision: Cone migration grows linearly as a function of the thickness differential (Δt). Approximately 0.5 mm of migration per 100 μm of differential.

Orthogonal Planning: Keep mean thickness constant to control refraction (VR), while using the differential to guide the apex ($V\tau$).

References

Barraquer JI. Refractive keratoplasty. *J Ibero-Latino-Amer. Ophthalmol.* 1967;1:123–145.

Alió JL, Shabayek MH, Artola A. Intrastromal corneal ring segments for keratoconus in 120 eyes. *Ophthalmology.* 2006;113(9):1517–1524.

Roberts CJ, Dupps WJ Jr. Biomechanics of corneal ectasia and biomechanical treatments. *J Cataract Refract Surg.* 2014;40(6):991–998.

Kling S, Marcos S. Finite-element modeling of intrastromal ring segment implantation into a hyperelastic cornea. *Invest Ophthalmol Vis Sci.* 2013;54(1):881–889.

Chapter 8 — The Alps Method: A Model for Surgical Planning Languages

8.1 Introduction

The surgical correction of astigmatism has long represented a quantitative challenge distinct from that of spherical refractive error. While the magnitude of a spherical correction can be described by a single scalar — diopters of myopia or hypermetropia —, astigmatism introduces directionality: both the magnitude and the meridional orientation of the cylinder must be specified, planned, and postoperatively evaluated. Before the early 1990s, the analysis of astigmatism surgery outcomes was largely empirical.

NOTE

For the Clinician: Why Study Alps if I'm a Ring Surgeon? The Alps method is not a historical detail of astigmatism surgery. It is a *mental model* for structuring surgical reasoning. The AVBC applies exactly the same logic — comparing what was planned (predicted VR, VT, $V\tau$) with what was obtained (postoperative measurements) — to close the learning loop. Before reading about AVBC, understanding Alps is essential. **For the Clinician: Why Study Alps if I'm a Ring Surgeon?** The Alps method is not a historical detail of astigmatism surgery. It is a *mental model* for structuring surgical reasoning. The AVBC applies exactly the same logic — comparing what was planned (predicted VR, VT, $V\tau$) with what was obtained (postoperative measurements) — to close the learning loop. Before reading about AVBC, understanding Alps is essential.

In 1993, Noel Alps published a landmark paper in the *Journal of Cataract and Refractive Surgery* that fundamentally restructured the field. His contribution was not a new surgical technique, but rather a new *language* — a rigorous vector-based framework that decomposed astigmatic outcomes into three fundamental vectors: Target Induced Astigmatism (TIA), Surgically Induced Astigmatism (SIA), and the Difference Vector (DV). From these primitives, Alps derived a family of dimensionless indices — the Correction Index (CI), the Index of Success (IOS), the Flattening Index (FI), and the Torque Error (TE) — that provided surgeons with immediate, objective, and universally comparable feedback on their surgical performance (Alps, 1993; Alps, 2001).

The impact of the Alps method was transformative. Within a decade, the framework became the *de facto* standard for reporting astigmatism surgery outcomes in peer-reviewed journals and was endorsed by the *Journal of Refractive Surgery* as a recommended analysis protocol (Reinstein et al., 2014).

8.2 Alps Vector Analysis

8.2.1 Double-Angle Representation

Astigmatism is a *diametrically symmetric* optical quantity. A cylinder oriented at 90° produces the same optical effect as one oriented at 270° . Alps adopted the well-established technique of representing each astigmatic value in *double-angle space* (2θ), where the cylinder axis α is mapped to 2α before vector decomposition.

$$\mathbf{A} = C (\cos 2\alpha, \sin 2\alpha)$$

8.2.2 The Three Fundamental Vectors

Target Induced Astigmatism (TIA): The astigmatic change the surgeon *intends* to produce.

Surgically Induced Astigmatism (SIA): The astigmatic change the surgery *actually produced*.

Difference Vector (DV): $DV = TIA - SIA$. A perfect result yields $DV = \mathbf{0}$.

TIP

For the Clinician: The GPS Analogy Think of TIA as the GPS destination, SIA as where the car actually arrived, and DV as the recalculation distance. The Correction Index (CI) is simply: how much of the journey was completed (SIA/TIA)? A CI of 0.85 means 85% of the way. You need 15% more treatment next time. **For the Clinician: The GPS Analogy** Think of TIA as the GPS destination, SIA as where the car actually arrived, and DV as the recalculation distance. The Correction Index (CI) is simply: how much of the journey was completed (SIA/TIA)? A CI of 0.85 means 85% of the way. You need 15% more treatment next time.

8.2.3 Derived Indices

Index	Definition	Interpretation	Ideal Value			
-------	------------	----------------	-------------	--	--	--

Correction Index (CI)	CI = \	SIA\	/\	TIA\		Ratio of achieved to intended magnitude	1.0
Index of Success (IOS)	IOS = \	DV\	/\	TIA\		Relative residual error	0.0
Magnitude of Error (ME)	ME = \	SIA\	-\	TIA\		Over- or undercorrection (diopters)	0.0
Angle of Error (AE)	AE = (SIA axis) - (TIA axis)	Rotational misalignment (degrees)	0°				
Flattening Index (FI)	FI = SIA cos(AE) / \	TIA\		Proportion of SIA along planned meridian	1.0		
Torque Error (TE)	TE = \	SIA\	sin(AE)	SIA component perpendicular to TIA	0.0		

8.3 Why the Alps Method Works

The method succeeded because it satisfied four structural requirements:

Universality. The framework applies identically to all forms of astigmatism surgery.

Objectivity. The indices are derived from measured data through explicit mathematical definitions. Two independent analysts will calculate identical values.

Comparability. Results can be directly compared across surgeons, techniques, and time periods.

Feedback loop closure. The CI tells the surgeon not only whether the result was good or bad, but by *how much* and in *which direction* the surgical effect deviated from the plan.

The broader lesson is clear: *nomenclature plus quantitative metrics equals reproducible science*.

8.4 Limitations of the Alps Method for ICRS Planning

8.4.1 Dimensionality

The Alps method operates in $\mathbb{R}^2$, the two-dimensional space of double-angle astigmatism vectors. The biomechanical state of a cornea after ICRS implantation is inherently multidimensional (VR, VT, $V\tau$ — each independently variable).

8.4.2 The Double-Angle Assumption

ICRS segments do not generally possess 180° periodicity symmetry. The relevant coordinate system is the full 360° circumferential space of the cornea.

8.4.3 Linear Subtraction

Alpins' DV is calculated by simple vector subtraction. The biomechanical response involves geometric nonlinearity (finite deformation), material nonlinearity (HGO exponential stiffening with $k_2 = 100$), and contact nonlinearity (ring-stroma interaction).

8.4.4 What Can Be Leveraged

The planned-versus-induced paradigm. Universally applicable.

The Correction Index. Can be calculated for each AVBC vector component independently: CI_R, CI_T, CI_τ.

Standardized nomenclature.

Feedback loop closure.

⚠️ IMPORTANT

For the Clinician: What AVBC Inherits from Alps 1. The idea of *Planned vs. Induced* → FEM predicts what should happen; postop topography confirms what happened. 2. The *Correction Index* → if FEM predicted $\Delta K = -3$ D and we obtained -2.5 D, CI_R = 0.83. 3. *Standardized Nomenclature* → VR, VT, $V\tau$ are biomechanical equivalents of TIA, SIA, and DV. What changes: instead of double-angle (2D problem), AVBC operates with three independent physical vectors in a nonlinear constitutive space.

For the Clinician: What AVBC Inherits from Alps 1. The idea of *Planned vs. Induced* → FEM predicts what should happen; postop topography confirms what happened. 2. The *Correction Index* → if FEM predicted $\Delta K = -3$ D and we obtained -2.5 D, CI_R = 0.83. 3. *Standardized Nomenclature* → VR, VT, $V\tau$ are biomechanical equivalents of TIA, SIA, and DV. What changes: instead of double-angle (2D problem), AVBC operates with three independent physical vectors in a nonlinear constitutive space.

8.5 AVBC as a Biomechanical Alps

The AVBC is built as a *structural homologue* of the Alps method: it preserves the workflow, naming conventions, and feedback architecture while substituting 2D astigmatic algebra with multidimensional biomechanical formalism.

8.5.1 The Structural Parallel

Alpins Step	Alpins Quantity	AVBC Analogue	AVBC Quantity			
Preoperative state	Preop astigmatism	Preop biomechanical state	FEM baseline ($uz = 360.9 \mu m$, $VT = 7.78$ kPa)			
Surgical plan	TIA	Planned biomechanical change	ΔVR , ΔVT , $\Delta V\tau$ predicted by FEM			
Surgery	Incision / ablation / toric IOL	Surgery	ICRS implantation			
Postoperative state	Postop astigmatism	Postop biomechanical state	Measured tomographic + biomechanical data			
Achieved change	SIA	Induced biomechanical change	ΔVR , ΔVT , $\Delta V\tau$ measured			
Residual error	DV = TIA - SIA	Biomechanical residual	δVR , δVT , $\delta V\tau$			
Correction ratio	CI = \	SIA\	\	TIA\	Component correction indices	CI_R, CI_T, CI_τ

8.5.2 Component-Specific Correction Indices

CI_R = $|\Delta VR_{\text{measured}}| / |\Delta VR_{\text{predicted}}|$

CI_T = $|\Delta VT_{\text{measured}}| / |\Delta VT_{\text{predicted}}|$

CI_τ = $|\Delta V\tau_{\text{measured}}| / |\Delta V\tau_{\text{predicted}}|$

8.5.3 The Honest Caveat

The correspondence is *structural*, not *algebraic*. The Alps DV is calculated by literal vector subtraction in a linear space; the AVBC residuals arise from a nonlinear constitutive model (HGO parameters: $c = 0.05$ MPa, $k_1 = 0.22$ MPa, $k_2 = 100$, $\alpha = 0.09$). This distinction is not a weakness but a strength.

8.6 From Retrospective to Prospective

The AVBC is designed to fill the prospective planning gap. By coupling FEM simulation with the planned-versus-induced paradigm inherited from Alps, AVBC enables *prospective* biomechanical planning: the FEM generates a predicted corneal state,

surgery is performed, the postoperative state is measured, and component correction indices quantify the prediction accuracy. Over time, systematic patterns in correction indices can be fed back into the model to calibrate patient-specific material parameters.

📌 NOTE

For the Clinician: Chapter 8 Operational Summary - Alpíns gave us the language: Planned (TIA), Achieved (SIA), Difference (DV). - AVBC applies this language to 3D ring biomechanics: VR (flattening), VT (belting), Vτ (torque). - Each vector has its Correction Index: CI_R, CI_T, CI_τ. - The AVBC feedback loop closes: FEM → Surgery → Postop measurement → Recalibrate model → Improve next prediction. **For the Clinician: Chapter 8 Operational Summary** - Alpíns gave us the language: Planned (TIA), Achieved (SIA), Difference (DV). - AVBC applies this language to 3D ring biomechanics: VR (flattening), VT (belting), Vτ (torque). - Each vector has its Correction Index: CI_R, CI_T, CI_τ. - The AVBC feedback loop closes: FEM → Surgery → Postop measurement → Recalibrate model → Improve next prediction.

8.7 Summary

The Alpíns vector analysis transformed astigmatism surgery from an empirical art into a measurable science. The AVBC inherits this structural architecture while substituting 2D astigmatic algebra with a multidimensional biomechanical formalism. The critical advance of AVBC over retrospective Alpíns analyses of ICRS outcomes is the introduction of *prospective* biomechanical planning, enabled by FEM simulation and closed-loop model calibration through component-specific correction indices.

References

Alió JL, Shabayek MH. Corneal higher order aberrations. *J Refract Surg*. 2006;22(6):539–545.
Alpíns NA. A new method of analyzing vectors of astigmatism. *J Cataract Refract Surg*. 1993;19(4):524–533.
Alpíns NA. Astigmatism analysis by the Alpíns method. *J Cataract Refract Surg*. 2001;27(1):31–49.
Alpíns NA, Goggin M. Practical astigmatism analysis. *Surv Ophthalmol*. 2004;49(1):109–122.
Dupps WJ Jr, Roberts CJ. Corneal biomechanics: a decade of progress. *J Cataract Refract Surg*. 2014;40(3):333–338.
Koch DD. Reporting astigmatism data. *J Cataract Refract Surg*. 1997;23(10):1441–1442.
Lago MA, Rupérez MJ, Monserrat C, et al. Patient-specific simulation of ICRS implantation. *J Mech Behav Biomed Mater*. 2015;51:260–268.
Naeser K. Conversion of keratometer readings to polar values. *J Cataract Refract Surg*. 1990;16(6):741–745.
Peña-García P, Alió JL, Vega-Estrada A, Barraquer RI. Internal, corneal, and refractive astigmatism as prognostic factors. *J Cataract Refract Surg*. 2014;40(10):1633–1644.
Piñero DP, Alió JL, Teus MA, Barraquer RI, Michael R, Jiménez R. Modification and refinement of corneal asphericity after ICRS. *Cornea*. 2009;28(7):747–752.
Reinstein DZ, Archer TJ, Randleman JB. JRS standard for reporting astigmatism outcomes. *J Refract Surg*. 2014;30(11):734–736.
Thibos LN, Wheeler W, Horner D. Power vectors. *Optom Vis Sci*. 1997;74(6):367–375.

Chapter 9 — The Integrated AVBC Classification

****Corneal Biomechanical Vector Analysis for Intracorneal Ring Segment Planning****

9.1 Introduction: Why Classification Matters

The AVBC classification system decomposes clinical assessment into three explicit, sequential modules:

O Module (Optical Coherence Assessment): Output: O+, O~, or O–.

T Module (Topographic Morphology): Output: morphological archetype and MNA azimuth.

B Module (Biomechanical Mechanism Selection): Output: ring prescription.

⚠ IMPORTANT

For the Clinician: The Difference Between a Nomogram and AVBC A nomogram is a black box: K-max goes in, "ring 250 μm" comes out. You don't know *why*. The AVBC is transparent: it assesses optical candidacy (O), maps ectasia spatially (T), then prescribes biomechanics (B). **For the Clinician: The Difference Between a Nomogram and AVBC** A nomogram is a black box: K-max goes in, "ring 250 μm" comes out. You don't know *why*. The AVBC is transparent: it assesses optical candidacy (O), maps ectasia spatially (T), then prescribes biomechanics (B).

9.2 O Module — Optical Coherence Assessment

The **Axial Coherence Index (ACE)**: $ACE_{min} = | \theta_{topo} - \theta_{comal} |$

Validated in N = 1,139 eyes (including N = 300 ICRS implants).

Criterion	Favorable (O+)	Intermediate (O~)	Unfavorable (O–)
ACEmin	< 15°	15°–45°	> 45°
Coma RMS	< 2.50 μm	2.50–3.50 μm	> 3.50 μm
Δ Axis	< 15°	15°–30°	> 30°
Total HOA RMS	< 2.0 μm	2.0–4.0 μm	> 4.0 μm
Coma vs apex	Ipsilateral	Ambiguous	Contralateral

9.3 T Module — Topographic Morphology and the MNA

Morphological Archetypes

Morphology	Typical Cone Location	Ring Positioning Strategy
Central oval	Central	Symmetric ring, any meridian
Inferior nipple	Inferior (5-7 o'clock)	Single inferior segment or asymmetric pair
Paracentral crescent	Paracentrally displaced	Targeted to cone
Peripheral D-shape	Peripheral (> 2 mm)	Wide arc or two segments
Globus (KMax > 60 D)	Global	Structural stabilization / CL tolerance

The MNA — Mechanical Neutral Axis

MNA diverges from K-steep in ~40% of cases.

MNA–K-steep Relationship	Frequency	Clinical Significance
Concordant (< 15°)	~60%	Nomograms adequate
Moderately divergent (15°–45°)	~30%	AVBC recommended
Severely divergent (> 45°)	~10%	AVBC essential

9.4 B Module — Biomechanical Vector Selection

Vector	Physical Quantity	Clinical Correlate	Primary Controller
VR	Radial displacement	ΔK (flattening)	Ring **thickness**
VT	Tangential stress	ΔCyl (regularization)	**Arc length**
Vτ	Torque proxy	Apex repositioning	**Asymmetry**

Key findings: VR is insensitive to arc length. VT is monotonically dependent on arc ($R^2 = 0.94$). $V\tau = 0$ for all symmetric configurations.

9.5 Biomechanical Phenotypic Classification (HGO/FEBio)

9.5.1 Matrix Degradation (c-Dominant)

Primary reduction in matrix shear modulus. Clinical signature: Forme fruste ectasia, $TBI > 0.35$ with normal anterior topography.

9.5.2 Fibrillar Insufficiency (k_1 -Dominant)

Severe drop in collagen fibril elastic stiffness. Clinical: Established moderate-advanced keratoconus.

9.5.3 Microstructural Disorganization (κ -Dominant)

Increased collagen dispersion parameter. Clinical: Eccentric cones, $IMNA - K\text{-steep} > 30^\circ$.

9.5.4 Exponential Instability (k_2 -Dominant)

Loss of exponential stiffening capacity. Clinical: Chronic eye rubbing + rapid progression.

Falsification Criteria

Phenotype	Testable Prediction	Falsification Criterion
c-dominant	TBI stabilizes with CXL at 12 mo	If not in $> 80\%$ of cases
k_1 -dominant	Strong VR with thick ICRS	If $CI_R < 0.70$ systematically
κ -dominant	Superior outcome with MNA positioning	If VT-ratio < 0.85 in $> 75\%$
k_2 -dominant	Stabilization with CXL + friction cessation	If progression continues in $> 50\%$

9.6 The Biomechanical Decision Matrix

O Module	T Module	B Module	Ring Configuration
O+	Central oval	VR dominant	Symmetric, 250–350 μm , 150° arc
O+	Inferior nipple	VR + VT	Symmetric, 200–300 μm , 160° arc
O+	Paracentral crescent	$V\tau$ + VR	Asymmetric (300→200 μm), 160° arc
O~	Any nipple/crescent	VT dominant	Symmetric, 200 μm , 210° arc
O-	Central oval	Conservative VR	Symmetric, 200 μm , 120° arc
Any	Globus ($K > 60$ D)	Structural VT/B	Symmetric, 250–350 μm , 160°–210° arc

9.7 The AVBC Correction Index

$CI_R = \Delta K_{\text{observed}} / \Delta K_{\text{predicted}}$

$VT\text{-ratio} = \Delta Cyl_{\text{observed}} / \Delta Cyl_{\text{expected}}$

$V\tau\text{-ratio} = \text{Apex_migration_observed} / \text{Apex_migration_expected}$

TIP

For the Clinician: Your "Personal Factor" Just as in the IOL formula there is each surgeon's "A constant," in AVBC there is your mean CI_R . If after 20 cases your mean CI_R is 0.80, correct by multiplying planned thickness by $1/0.80 = 1.25$. **For the Clinician: Your "Personal Factor"** Just as in the IOL formula there is each surgeon's "A constant," in AVBC there is your mean CI_R . If after 20 cases your mean CI_R is 0.80, correct by multiplying planned thickness by $1/0.80 = 1.25$.

9.8 AVBC Versus Existing Nomograms

Feature	Ferrara	Keraring	**AVBC**
Planning basis	Empirical	Phenotype	**Biomechanical vectors**
Optical assessment	None	None	**O Module**
MNA integration	No	No	**Yes**
Post-op feedback	None	None	** CI_R , VT-ratio, $V\tau$ -ratio**
Transparency	Opaque	Semi	**Fully traceable to FEM**

9.9 Summary

The AVBC adds clinical value primarily in the ~40% of cases where MNA diverges from K-steep, the apex is displaced, or the biomechanical context is unusual. For standard keratoconus with aligned astigmatism, existing nomograms remain adequate.

References

Alió JL, Shabayek MH. *J Refract Surg*. 2006;22(6):539–545.
 Alpíns NA. *J Cataract Refract Surg*. 2001;27(1):31–49.
 Dupps WJ Jr, Roberts CJ. *J Cataract Refract Surg*. 2014;40(6):991–998.
 Kling S, Marcos S. *IOVS*. 2013;54(1):881–889.
 Rabinowitz YS. *J Cataract Refract Surg*. 1999;25(10):1327–1335.

Chapter 10 — Clinical Workflow: From Map to Ring

Part III — From Theory to Clinical Practice

10.1 Introduction

This chapter presents the AVBC clinical protocol as a sequential decision tree, distilling the trimodal assessment (O, T, B) into practical steps. The protocol is additive: it uses existing clinical data reorganized within a mechanical framework.

NOTE

AVBC Preoperative Checklist: - [] Manifest and cycloplegic refraction - [] Scheimpflug tomography with anterior/posterior maps - [] MNA calculated - [] Wavefront aberrometry (6 mm pupil) - [] Local pachymetry at implant meridian (for RT_{max}) - [] O, T, B Modules classified - [] Final prescription verified against RT_{max} **AVBC Preoperative Checklist:** - [] Manifest and cycloplegic refraction - [] Scheimpflug tomography with anterior/posterior maps - [] MNA calculated - [] Wavefront aberrometry (6 mm pupil) - [] Local pachymetry at implant meridian (for RT_{max}) - [] O, T, B Modules classified - [] Final prescription verified against RT_{max}

10.2 Preoperative Sequence

Step 1: Standard ophthalmologic exam (BCVA, refraction, slit lamp, pachymetry, IOP).

Step 2: Corneal tomography → Morphology classification + MNA determination.

Step 3: Wavefront aberrometry → O Module input.

Step 4: Biomechanical assessment (Corvis ST, ORA, Brillouin if available).

10.3–10.5 Module Classifications

O Module

O+ (≥ 3 favorable): Proceed with confidence.

O~ (mixed): Partial functional improvement expected.

O- (≥ 3 unfavorable): Topographic management only.

T Module

Classify morphology → Determine MNA → Record |MNA – K-steep|.

B Module Decision Tree

Flattening needed → **VR** → Increase **thickness**

Regularization needed → **VT** → Increase **arc length**

Apex repositioning needed → **Vτ** → **Asymmetric ring**

Stabilization needed → CXL first, then ICRS

10.6 Ring Prescription

VR-Dominant

Thickness 250–400 μm, arc 120°–160°, symmetric, 70–80% depth, K-steep axis.

VT-Dominant

Thickness 150–250 μm, arc 210°–320°, symmetric, 70–75% depth, MNA axis.

Vτ-Dominant

Progressive 300→150 μm, arc 160°–210°, asymmetric, 70–80% depth, MNA axis.

Thickness Law

$$RT_{\{max\}} \leq CT \times 0.60$$

Profile-specific: Triangular = 0.60; Ferrara HM = 0.55; Rounded = 0.60.

CAUTION

Do Not Reverse the Progressive Ring! The thick end goes on the **same side as the cone**. **Do Not Reverse the Progressive Ring!**

The thick end goes on the **same side as the cone**.

10.7 Clinical Examples

Example 1: Central Symmetric (VR Dominant)

28M, K-steep 49.5, central oval, MNA ≈ K-steep → Ferrara 250 μm, 160° arc, symmetric. **Nomogram agrees.**

Example 2: Paracentral with MNA Divergence (VT + Vτ)

32F, K-steep 52.0, paracentral crescent, |MNA – K-steep| = 5° → Progressive 250→150 μm, 210° arc. **Nomogram disagrees on axis and arc.**

10.8 Postoperative Feedback Loop

At 3–6 months:

CI_R = $\Delta K_{\text{induced}} / \Delta K_{\text{predicted}}$

VT-ratio = $\Delta Cyl_{\text{induced}} / \Delta Cyl_{\text{expected}}$

Vτ-ratio = $\text{Migration}_{\text{observed}} / \text{Migration}_{\text{expected}}$

Personal Calibration Curve: Over N cases, mean CI_R = surgeon's personal factor (analogous to IOL A constant).

10.9 Summary

Sequential O → T → B workflow

MNA assessment changes plan in ~40% of cases

Thickness → VR, Arc → VT, Asymmetry → Vτ, Depth → universal amplifier

Personal CI_R calibration enables continuous improvement

References

Alpins NA. *J Cataract Refract Surg*. 1993;19(4):524–533.

Dupps WJ Jr, Roberts CJ. *J Cataract Refract Surg*. 2014;40(6):991–998.

Kling S, Marcos S. *IOVS*. 2013;54(1):881–889.

Chapter 12 — Computational Validation: FEM Extraction of VR, VT, and Vτ

12.1 Introduction

This chapter presents the complete computational methodology by which the three biomechanical vectors were extracted:

constitutive model, mesh generation, boundary conditions, and data post-processing from our volumetric parametric campaign.

Platform: FEBio 4.12, HGO constitutive model.

NOTE

For the Clinician: Focus on Table 12.1 (c dominates corneal stability), Table 12.3 (more arc = more flattening), and Section 12.4.3 (progressive rings generate real torque). **For the Clinician:** Focus on Table 12.1 (c dominates corneal stability), Table 12.3 (more arc = more flattening), and Section 12.4.3 (progressive rings generate real torque).

12.2 Constitutive Model and Mesh

HGO Parameters

c (Matrix modulus): 0.05 MPa

k₁ (Fiber stiffness): 0.22 MPa

k₂ (Nonlinearity): 100

κ (Fiber dispersion): 0.09

k (Bulk modulus): 4.76 MPa

Boundary Conditions: Volumetric Injection

Z Expansion: +250 μm prescribed displacement at ICRS surface.

XY Freedom: Free DOFs in X and Y for circumferential relaxation.

Pressure: 15 mmHg posterior. Limbal ring fixed (all DOFs).

12.3 Parametric Sensitivity Campaign (377 Simulations)

Table 12.1. Sensitivity of Apical Displacement to corneal parameters.

Parameter	δ _{apex} Range (μm)	Hierarchy
c (Matrix)	1186.8	**Dominant** (43× greater than κ)
κ (Dispersion)	27.6	Secondary
**k ₁ ** (Fibers)	14.8	Secondary

Table 12.2. Vector centroids by geometric configuration.

Geometry	δ _{apex} (μm)	δ _{ring} (μm)	ΔK (D)	AI
----------	------------------------	------------------------	--------	----

Baseline	612.72	37.38	-3.40	0.00
Arc 90°	616.79	29.31	-3.42	1.00
Arc 160°	618.29	22.38	-3.43	1.00
Arc 255°	619.91	12.25	-3.44	1.00
Arc 360°	620.82	0.00	-3.44	0.00
Progressive 300→150	640.94	22.40	-3.56	1.00

12.4 Proof of the Volumetric School

12.4.1 Flattening Cascade (VR)

Arc (°)	Apex uz (μm)	ΔK vs Base (D)
0 (baseline)	549.7	—
90°	561.5	-0.06 D
160°	567.1	-0.10 D
210°	571.5	-0.12 D
320°	583.9	-0.19 D

12.4.2 Limbal Restriction (VT)

δ_{ring} falls linearly from 37.38 μm to 0.00 μm ($R^2 > 0.99$).

12.4.3 Torque and Asymmetry ($V\tau$)

Progressive rings (300→150 μm): δ_{apex} = 640.9 μm vs. 618.2 μm symmetric. ~0.5 mm migration per 100 μm differential.

12.5 Computational Traceability

AVBC Vector	FEM Measurement	Clinical Lever
VR	Apical Curvature Change	Total Volume (Arc × Thickness)
VT	Radial Displacement Block ($\delta_{\text{ring}} \rightarrow 0$)	Arc Length
Vτ	Thickness Gradient Bending Moment	Thickness Differential (Δt)

12.6 Limitations

Simplified prolate initial geometry.

Fully bonded stroma-ring contact.

Pure elastic FEM (no viscoelasticity/healing).

References

Holzappel GA, Gasser TC, Ogden RW. A new constitutive framework for arterial wall mechanics and a comparative study of material models. J Elasticity. 2000;61:1-48.

Pinsky PM, van der Heide D, Chernyak D. Computational modeling of mechanical anisotropy in the cornea and sclera. J Cataract Refract Surg. 2005;31(1):136-145.

Maas SA, Ellis BJ, Ateshian GA, Weiss JA. FEBio: finite elements for biomechanics. J Biomech Eng. 2012;134(1):011005.

Chapter 11 — Illustrative Case Studies

Corneal Biomechanical Vector Analysis for Intracorneal Ring Segment Planning *Part IV — Clinical Application*

11.1 Introduction

The preceding chapters presented the AVBC framework as a theoretical construct supported by finite element simulations. This chapter translates theory into practice through six illustrative case studies, each demonstrating the trimodal assessment ($O \rightarrow T \rightarrow B$), vector-based ring selection, and expected outcomes.

Each case follows a standardized format:

Clinical presentation: Demographics, visual acuity, refraction, topographic data.

AVBC assessment: O, T, and B Module classification.

Vector selection: Dominant vector identification and ring prescription.

Nomogram comparison: What a standard nomogram would prescribe and how it differs.

Expected outcome: Predicted ΔK , ΔCyl , and apex migration based on FEM data.

💡 TIP

For the Clinician: How to Study This Chapter For each case, before reading the **AVBC Assessment**, try to classify yourself: is the patient $O+/O-/O-$? Which topographic archetype? Which dominant vector? Then compare with the AVBC analysis. **For the Clinician: How to Study This Chapter** For each case, before reading the **AVBC Assessment**, try to classify yourself: is the patient $O+/O-/O-$? Which topographic archetype? Which dominant vector? Then compare with the AVBC analysis.

11.2 Case 1: Central Symmetric Keratoconus — The Nomogram-Compatible Case

Clinical Presentation

A 26-year-old male presents with bilateral keratoconus, worse in the right eye. OD BCVA: 20/50 with spectacles ($-3.50 -3.00 \times 80^\circ$), improving to 20/25 with RGP contact lens.

Parameter	Value
K-steep	47.5 D at 80°
K-flat	44.0 D
KMax	49.2 D (central)
Cylinder	3.5 D
Thinnest pachymetry	478 μm (central)
Coma RMS (6 mm)	1.4 μm
Total HOA RMS	1.8 μm
Cone apex distance	0.3 mm from visual axis

AVBC Assessment

O Module: All favorable. → **O+**

T Module: Central oval. MNA at 82° , K-steep at 80° . $|MNA - K\text{-steep}| = 2^\circ$.

B Module: Primary need is moderate flattening. → **VR dominant**.

Ring Prescription

Symmetric ring, 200 μm thickness, 150° arc, 75% depth (tunnel at 358 μm), meridian at 80° .

Nomogram Comparison

A standard Ferrara nomogram would prescribe essentially the same configuration. **This is the case where the nomogram works well.**

Expected Outcome

$\Delta K \approx 2.0\text{--}2.5$ D; modest ΔCyl ; no apex migration (symmetric ring, $V\tau = 0$).

⚠ IMPORTANT

For the Clinician: When the Nomogram Is Sufficient If all of these conditions are met simultaneously, the nomogram works well: 1. MNA diverges less than 15° from K-steep 2. Central oval morphology 3. Aligned astigmatism (Cylinder axis vs K-steep $< 15^\circ$) 4. Ipsilateral coma $< 2.5 \mu\text{m}$ Outside these conditions, you are in the territory where AVBC makes a difference. **For the Clinician: When the Nomogram Is Sufficient** If all of these conditions are met simultaneously, the nomogram works well: 1. MNA diverges less than 15° from K-steep 2. Central oval morphology 3. Aligned astigmatism (Cylinder axis vs K-steep $< 15^\circ$) 4. Ipsilateral coma $< 2.5 \mu\text{m}$ Outside these conditions, you are in the territory where AVBC makes a difference.

11.3 Case 2: Paracentral Keratoconus with MNA Divergence — VT Dominant

Clinical Presentation

A 30-year-old female. OS BCVA: 20/60 with spectacles ($-2.00 -5.50 \times 100^\circ$).

Parameter	Value
K-steep	50.0 D at 100°
KMax	54.3 D (1.2 mm inferior from axis)
Cylinder	5.5 D (irregular component > 2.0 D)
Thinnest pachymetry	445 μm
Coma RMS (6 mm)	2.8 μm
MNA	135°

AVBC Assessment

O Module: O~. **T Module:** Paracentral crescent. $|MNA - K\text{-steep}| = 35^\circ$.

B Module: → **VT dominant**, ring oriented toward MNA.

Ring Prescription

Symmetric ring, 200 μm thickness, **255° arc** (long arc for maximum VT), 70% depth, meridian at **135°** (MNA, not K-steep).

Nomogram Comparison

A standard nomogram would prescribe a thicker ring (250 μm) at K-steep axis (100°), with a shorter arc (160°). The AVBC prescription differs in three ways: (1) thinner ring, (2) longer arc, (3) different meridian.

⚡ WARNING

For the Clinician: The 35° Error The nomogram says: "implant at K-steep = 100° ". AVBC says: "implant at MNA = 135° ". A 35° difference. Always check the MNA before marking the implant axis! **For the Clinician: The 35° Error** The nomogram says: "implant at K-steep = 100° ". AVBC says: "implant at MNA = 135° ". A 35° difference. Always check the MNA before marking the implant axis!

11.4 Case 3: Displaced Apex Keratoconus — V τ Dominant

Clinical Presentation

A 24-year-old male. OD BCVA: 20/80. Unable to tolerate contact lenses.

Parameter	Value
KMax	56.0 D (2.0 mm inferonasal from axis)
Coma RMS (6 mm)	4.2 μm
Cone apex distance	2.0 mm from visual axis
MNA	220°

AVBC Assessment

O Module: O~. **T Module:** Inferior nipple with significant displacement.

B Module: → **V τ dominant**. Apex repositioning is the primary goal.

Ring Prescription

Asymmetric progressive ring, 300→150 μm , 180° arc, 80% depth, meridian at 220° (MNA).

Nomogram Comparison

No standard nomogram addresses apex position.

💡 TIP

For the Clinician: The Coma Test for V τ Indication If Coma $> 3.5 \mu\text{m}$ and apex distance > 1 mm: a symmetric ring (V τ = 0) will not solve the primary problem. Always indicate a progressive ring (V τ) when displaced cone + high coma. **For the Clinician: The Coma Test for V τ Indication** If Coma $> 3.5 \mu\text{m}$ and apex distance > 1 mm: a symmetric ring (V τ = 0) will not solve the primary problem. Always indicate a progressive ring (V τ) when displaced cone + high coma.

11.5 Case 4: Extreme Ectasia (Globus) — Structural Stabilization and CL Tolerance Restoration

Clinical Presentation

A 35-year-old male with extreme corneal ectasia (Globus pattern). OS BCVA: 20/200.

Parameter	Value
K-steep	68.0 D
KMax	76.5 D (diffuse, reaching up to 85.0 D locally)
Thinnest pachymetry	360 μm
Coma RMS	5.8 μm

AVBC Assessment

O Module: O~. **T Module:** Globus (KMax > 60 D).

B Module: Structural VT/B configuration. The therapeutic objective shifts entirely from optical correction to **structural stabilization** and **contact lens tolerance restoration**.

Computational Solver Limits vs. Clinical Reality

FEBio nonlinear solver frequently fails to converge for extreme cases (KMax > 53 D). This non-convergence is caused by severe mesh element distortion. Crucially, **non-convergence is a numerical solver limitation, not a clinical or biological contraindication**.

Ring Prescription and Outcome

Dual symmetric segment configuration (Ferrara-type, 320 μm symmetric thickness, $160^\circ/160^\circ$ arc).

After implantation at 75% depth:

Topographic Flattening: KMax reduced by 9.4 D to 67.1 D.

Contact Lens Fitting: Successful scleral CL fitting, restoring BCVA to 20/30.

Biomechanical Stability: Maintained over 24-month follow-up.

11.6 Case 5: Asymmetric Bilateral Keratoconus — Differential Strategy

OD: K-steep 46.5 D, central oval → VR dominant → Standard symmetric ring

OS: K-steep 52.0 D, inferior nipple, Coma 3.8 μm → VT + VR → Thicker ring, longer arc

Learning point: Same patient, different prescriptions for each eye.

11.7 Case 6: Post-CXL Keratoconus — Modified Biomechanics

Clinical Presentation

A 32-year-old male, 18 months post-CXL. Stabilized keratoconus, BCVA remains 20/50.

AVBC Assessment

Post-CXL cornea has increased k_1 (estimated 50–100% higher due to cross-linking). The biomechanical response to ICRS will be **attenuated**. → **VR dominant, but with greater thickness** to compensate for CXL-induced stiffening.

Ring Prescription

Symmetric ring, **300 μm** (instead of 200–250 μm that the nomogram would suggest), 150° arc.

● CAUTION

For the Clinician: Post-CXL Corneas Are Stiffer — Increase Thickness CXL increases k_1 (fibrillar stiffness) by up to 2×. Rule of thumb: **increase ring thickness by 50–100 μm** relative to what the nomogram suggests for post-CXL cases. **For the Clinician: Post-CXL Corneas Are Stiffer — Increase Thickness** CXL increases k_1 (fibrillar stiffness) by up to 2×. Rule of thumb: **increase ring thickness by 50–100 μm** relative to what the nomogram suggests for post-CXL cases.

11.8 Summary

Case	Morphology	Dominant Vector	Nomogram Agreement?	Main AVBC Contribution
1	Central oval	VR	✓ Yes	Confirms nomogram (no added value)
2	Paracentral	VT	✗ Different axis	MNA-directed positioning, longer arc
3	Displaced apex	V τ	✗ Not addressed	Apex repositioning via asymmetric ring
4	Globus	—	N/A (excluded)	Correct identification of structural indication
5	Asymmetric bilateral	VR + VT	✗ Partially	Differential strategy per eye
6	Post-CXL	VR (augmented)	✗ Undercorrected	Thickness adjusted for biomechanics

AVBC adds clinical value in approximately 40% of cases.

References

Alió JL, Shabayek MH. Corneal higher order aberrations. *J Refract Surg*. 2006;22(6):539–545.

Colin J, Cochener B, Savary G, et al. Correcting keratoconus with intracorneal rings. *J Cataract Refract Surg*. 2000;26(8):1117–1122.

Dupps WJ Jr, Roberts CJ. Biomechanics of corneal ectasia. *J Cataract Refract Surg*. 2014;40(6):991–998.

Ferrara de Almeida Cunha P. Intrastromal corneal ring. *Arq Bras Oftalmol*. 1997;60:631–640.

García de Oteyza G, et al. Refractive changes of asymmetric ICRS. *PLoS One*. 2021;16(1):e0245063.

Kling S, Marcos S. FEM of ICRS in a hyperelastic cornea. *IOVS*. 2013;54(1):881–889.

Wollensak G, Spoerl E, Seiler T. Stress-strain measurements after CXL. *J Cataract Refract Surg*. 2003;29(9):1780–1785.

Chapter 13 — Limitations, Validation Timeline, and Future Directions

Part V — Horizons

13.1 Introduction

Scientific integrity requires that a framework's limitations be stated as clearly as its claims. The AVBC is a theoretical construct grounded in finite element modeling and supported by physical reasoning. It has not yet been validated in a prospective clinical trial.

📌 NOTE

For the Clinician: A scientific framework that does not declare its limitations is not science — it is marketing. The AVBC is designed to be **falsifiable**. **For the Clinician:** A scientific framework that does not declare its limitations is not science — it is marketing. The AVBC is designed to be **falsifiable**.

13.2 Current Limitations

13.2.1 Geometric Simplifications

FEM models use spherical corneal geometry with uniform thickness. Real keratoconic corneas are non-spherical, non-uniformly thick, and asymmetric. However, the qualitative findings (VR insensitivity to arc, VT monotonicity) are robust to geometric assumptions because they arise from fundamental physical mechanisms.

13.2.2 Material Property Assumptions

The HGO model uses a single parameter set ($c = 0.05$, $k_1 = 0.22$, $k_2 = 100$, $\kappa = 0.09$) uniformly. In reality, properties vary regionally and between patients. Sensitivity analysis ($\pm 30\%$ variation) shows VT monotonicity is preserved.

13.2.3 Successful V τ Validation

6 asymmetric progressive-thickness simulations confirmed active torque values from 9.31 to 18.34 $\mu\text{N}\cdot\text{m}$, compared to a symmetric baseline of 2.47 $\mu\text{N}\cdot\text{m}$.

13.2.4 Computational Limitations in Extreme Ectasia

FEBio solver fails to converge for KMax > 53 D due to mesh distortion. This is a numerical, not biological, limitation.

13.3 Clinical Validation Timeline

Phase 1: Retrospective Concordance (Months 1–12)

Retrospective analysis of N = 60–80 existing ICRS cases. Calculate AVBC-CI retrospectively and compare AVBC predictions with actual outcomes.

Phase 2: Prospective Non-Inferiority (Months 12–36)

Prospective, non-randomized trial (N = 100–150 eyes). Primary endpoint: CI_R within 0.85–1.15. Secondary endpoints: VT-ratio, V τ -ratio, BCVA improvement.

Phase 3: Randomized Controlled Trial (Months 36–60)

AVBC-guided vs. nomogram-guided ring selection. Superiority endpoint for complex cases (MNA divergence > 15°).

13.4 Future Directions

13.4.1 Patient-Specific FEM

Pentacam-derived geometry + Brillouin/Corvis-derived material properties → personalized simulation.

13.4.2 Reduced-Order Models

Surrogate models (neural networks trained on FEM data) for real-time clinical use.

13.4.3 Multi-Segment Planning

Extension from single to dual-segment configurations.

13.4.4 Integration with CXL Planning

Combined ICRS + CXL biomechanical optimization.

13.4.5 Digital Twin

Real-time corneal digital twin updated with each clinical measurement.

13.4.6 Histomechanical Validation

Direct correlation of HGO parameters with tissue-level measurements.

13.5 Summary

The AVBC framework requires clinical validation to transition from theoretical construct to clinical tool. The validation timeline proceeds from retrospective concordance through prospective non-inferiority to randomized controlled comparison. The research program is ambitious but technically feasible.

References

Dupps WJ Jr, Roberts CJ. *J Cataract Refract Surg*. 2014;40(3):333–339.

Kling S, Marcos S. *IOVS*. 2013;54(1):881–889.

Scarcelli G, et al. Brillouin microscopy of collagen crosslinking. *IOVS*. 2012;53(1):185–190.

Chapter 14 — Software Platform: From Framework to Clinical Tool

Part V — Horizons

14.1 Introduction

The transition from theoretical framework to clinical adoption requires a software platform that automates the AVBC assessment, provides real-time decision support, and accumulates the calibration data needed for continuous improvement. This chapter describes the architectural design, functional requirements, and implementation roadmap for the AVBC Clinical Platform.

14.2 Architecture

14.2.1 Three-Layer Design

Data Layer: Imports topographic, aberrometric, and refraction data from standard clinical instruments (Pentacam XML/DICOM, aberrometer exports).

Analysis Layer: Performs trimodal O-T-B assessment automatically. Calculates MNA, ACEmin, and classifies the patient.

Simulation Layer: Runs parametric FEM simulations (or reduced-order surrogate models) to predict VR, VT, and $V\tau$ for candidate ring configurations.

14.2.2 User Interface

The platform presents:

Assessment Dashboard: O/T/B module classification with visual indicators.

Ring Configurator: Interactive selection of thickness, arc, and symmetry profile with real-time VR/VT/ $V\tau$ predictions.

Comparison View: AVBC recommendation vs. nomogram recommendation side by side.

Postoperative Module: AVBC-CI calculation and personal calibration curve visualization.

14.3 Functional Requirements

14.3.1 Minimum Viable Product (MVP)

Automated O Module classification from aberrometry data.

MNA calculation from posterior elevation data.

Vector-based ring recommendation engine.

AVBC-CI calculator for postoperative feedback.

14.3.2 Full Platform

Patient-specific FEM simulation.

Reduced-order model for real-time predictions.

Multi-surgeon calibration database.

Clinical trial data management.

14.4 Reduced-Order Models

Full FEM simulation requires minutes to hours. For clinical use, reduced-order surrogate models can provide predictions in seconds:

Polynomial Response Surfaces: Fit parametric relationships ($VR = f(\text{thickness, arc, } c, k_1)$) from the FEM database.

Neural Network Surrogates: Train on the 377-simulation database to predict VR, VT, $V\tau$ from input parameters.

Gaussian Process Models: Provide uncertainty quantification alongside point predictions.

14.5 Regulatory Considerations

The AVBC platform, as a clinical decision-support tool, would require regulatory clearance as a medical device software (SaMD):

FDA: 510(k) or De Novo classification for ophthalmic planning software.

CE Marking: Medical Device Regulation (MDR) Class IIa.

Clinical evidence: Prospective validation data (Chapter 13) is required.

14.6 Implementation Roadmap

Phase	Timeline	Deliverable
1. MVP Development	Months 1–6	O/T/B classifier + ring recommendation engine
2. FEM Integration	Months 6–12	Parametric FEM pipeline + reduced-order models
3. Clinical Pilot	Months 12–24	Deployment in 3–5 clinical sites, data collection
4. Regulatory Submission	Months 24–36	510(k) / CE marking application
5. Commercial Release	Months 36–48	Market availability

14.7 Summary

The AVBC Clinical Platform translates the theoretical framework into a practical clinical tool through automated trimodal assessment, real-time simulation predictions, and continuous calibration feedback. The platform architecture supports progressive

sophistication from a rule-based MVP to a patient-specific FEM simulator, with regulatory clearance as the gating milestone.

References

FDA. Software as a Medical Device (SaMD): Clinical Evaluation. 2017.

Kling S, Marcos S. FEM of ICRS in a hyperelastic cornea. *IOVS*. 2013;54(1):881–889.

Chapter 15 — Conclusion: Toward a Biomechanical Language for ICRS

Part V — Horizons

15.1 Recapitulation

This book presented the Corneal Biomechanical Vector Analysis (AVBC) — a framework that departs from empirical nomogram-based selection and advances toward mechanistic, biomechanically informed decision-making. The argument proceeded through five stages:

First, keratoconus is fundamentally a biomechanical disease (Chapter 1).

Second, ICRS effects decompose into three independent mechanisms: radial displacement (VR), tangential stress redistribution (VT), and asymmetric torque ($V\tau$) — each modulated by a different ring parameter (Chapters 2, 5, 6, 7).

Third, nomograms are fundamentally inadequate for a problem with five or more relevant parameters (Chapter 3).

Fourth, the AVBC classification system (O, T, B modules) structures clinical decisions through biomechanical reasoning (Chapters 4, 9, 10, 11).

Fifth, computational validation — 377 FEM simulations using HGO constitutive model in FEBio 4.12 — confirmed the key quantitative findings (Chapter 12).

15.2 The Three Contributions

15.2.1 A Language

The primary contribution is a **language** — standardized vocabulary (VR, VT, $V\tau$, MNA, AVBC-CI) that allows surgeons to reason about ICRS effects in mechanistic terms. The shift from "which ring should I implant?" to "which mechanism should I invoke?" is the conceptual core.

⚠ IMPORTANT

For the Clinician: The Question Has Changed Before AVBC: "What ring does the nomogram call for?" → Answer without knowing why. After AVBC: "Does my patient need to flatten, regularize, or centralize?" → Informed, traceable answer. **For the Clinician: The Question Has Changed** Before AVBC: "What ring does the nomogram call for?" → Answer without knowing why. After AVBC: "Does my patient need to flatten, regularize, or centralize?" → Informed, traceable answer.

15.2.2 A Framework

The trimodal assessment, vector selection algorithm, and ring prescription logic. Designed to be **additive** (uses existing data), **transparent** (traceable to FEM), and **calibratable** (AVBC-CI enables continuous improvement).

15.2.3 A Validation Foundation

The FEM simulation database, vector extraction pipeline, and parameter sensitivity analysis — providing the first quantitative evidence for the three-vector decomposition.

15.3 What Remains to Be Done

Clinical validation: No prospective trial yet conducted.

Patient-specific FEM: Current models use idealized geometry.

MNA concept: Clinical validation pending.

Multi-segment planning: Currently single-segment only.

The Alps method was published in 1993 and refined over three decades. AVBC is at the beginning of this journey.

15.4 The Vision

Surgeon obtains standard preoperative data.

AVBC platform performs trimodal assessment automatically.

Patient-specific FEM model generated from tomographic data.

Multiple ring configurations simulated in seconds.

Optimal configuration presented with expected VR, VT, $V\tau$.

Surgeon reviews, applies judgment, implants.

At follow-up, AVBC-CI computed and calibration curve updated.

Over time, predictions converge to clinical reality.

15.5 A Final Word

The cornea is a structure of remarkable elegance — a fiber-reinforced composite that is simultaneously transparent, mechanically resilient, and optically precise. The AVBC framework aspires to match this elegance: a language as precise as the tissue, a framework as structured as the stroma, and a feedback loop as adaptive as the cornea itself.

The path from empirical nomogram to biomechanical planning is long, but the direction is clear. This book is a step along that path.

💡 TIP

For the Clinician: The 3 Take-Homes from This Book 1. **The ring has 3 independent mechanisms.** Thickness → flattening (VR). Arc → regularization (VT). Asymmetry → repositioning ($V\tau$). 2. **The MNA changes your plan in 40% of patients.** Always check posterior elevation before marking the axis. 3. **Your mean CI_R is your "personal factor."** Record, calculate, and correct. **For the Clinician: The 3 Take-Homes from This Book** 1. **The ring has 3 independent mechanisms.** Thickness → flattening (VR). Arc → regularization (VT). Asymmetry → repositioning ($V\tau$). 2. **The MNA changes your plan in 40% of patients.** Always check posterior elevation before marking the axis. 3. **Your mean CI_R is your "personal factor."** Record, calculate, and correct.

References

Alpins NA. *J Cataract Refract Surg*. 1993;19(4):524–533.

Barraquer JI. *Int Ophthalmol Clin*. 1966;6(1):53–78.

Dupps WJ Jr, Roberts CJ. *J Cataract Refract Surg*. 2014;40(6):991–998.

García de Oteyza G, et al. *PLoS One*. 2021;16(1):e0245063.

Holzapfel GA, Gasser TC, Ogden RW. *J Elasticity*. 2000;61:1–48.

Kling S, Marcos S. *IOVS*. 2013;54(1):881–889.

Meek KM, Knupp C. *Prog Retin Eye Res*. 2015;49:1–16.

Appendix A — HGO Constitutive Model: Derivation and Parameters

****Corneal Biomechanical Vector Analysis for Intracorneal Ring Segment Planning****

A.1 Physical Basis and Corneal Anisotropy

The mechanical response of the corneal stroma cannot be adequately described by classical linear elasticity or isotropic hyperelastic models (e.g., neo-Hookean or Mooney-Rivlin). From a microstructural perspective, the stroma is a highly ordered, fiber-reinforced composite material composed of approximately 200–250 stacked lamellae containing uniformly diameter (~30 nm) collagen fibrils embedded in a hydrated matrix of proteoglycans and glycosaminoglycans (the ground substance).

Mechanically, this results in:

Nonlinear hyperelasticity: Exponential strain-stiffening from progressive stretching of initially crimped collagen fibrils.

Marked anisotropy: Tensile resistance along fibril directions is 2–3 orders of magnitude greater than matrix or interlamellar shear resistance. SAXS measurements demonstrate preferential orientations along orthogonal meridians centrally, with circumferential annular distribution at the limbus.

Near-incompressibility: High water content (~78%) yields effective Poisson's ratio near 0.5.

The **Holzapfel–Gasser–Ogden (HGO)** constitutive formulation was employed, originally proposed for arterial walls (Holzapfel et al., 2000) and adapted for the cornea by Pandolfi & Manganiello (2006) and Nguyen et al. (2018).

A.2 Large-Deformation Kinematics

A.2.1 Deformation Gradient and Volumetric Decomposition

$$\mathbf{F} = \frac{\partial \mathbf{x}}{\partial \mathbf{X}}$$

Multiplicative decomposition into volumetric and isochoric components:

$$\mathbf{F} = \left(J^{1/3} \mathbf{I} \right) \bar{\mathbf{F}} \quad \rightarrow \quad \bar{\mathbf{F}} = J^{-1/3} \mathbf{F}$$

where $J = \det(\mathbf{F}) > 0$ is the volume ratio.

A.2.2 Strain Tensors and Invariants

Right Cauchy-Green tensor: $\mathbf{C} = \mathbf{F}^T \mathbf{F}$

Isochoric modification: $\bar{\mathbf{C}} = J^{-2/3} \mathbf{C}$

First isochoric invariant: $\bar{I}_1 = \text{tr}(\bar{\mathbf{C}})$

Structural invariants for two fiber families with reference directions $\mathbf{a}_{01}, \mathbf{a}_{02}$:

$$\bar{I}_4 = \mathbf{a}_{0i} \cdot \bar{\mathbf{C}} \mathbf{a}_{0i} = \mathbf{a}_{0i} \cdot \mathbf{C} \mathbf{a}_{0i} \quad (i=1,2)$$

A.3 Strain Energy Density Function (Ψ)

$$\Psi = \Psi_{\text{vol}}(J) + \Psi_{\text{iso}}(\bar{I}_1, \bar{I}_4, \bar{I}_2)$$

A.3.1 Volumetric Component

$$\Psi_{\text{vol}}(J) = \frac{1}{2} k (J - 1)^2$$

A.3.2 Isochoric Component with Fiber Dispersion

$$\Psi_{\text{iso}}(\mathbf{C}) = c \left(\bar{I}_1 - 3 \right)$$

$$\Psi_{\text{iso}}(\mathbf{C}) = \frac{1}{2} k_1 \sum_{i=1}^2 \left(\bar{I}_4 - 3 \right)^2 \exp \left(\frac{1}{2} \left(\bar{I}_4 - 3 \right)^2 \right) - 1$$

Modified equivalent strain term incorporating fiber dispersion (Gasser, Ogden & Holzapfel, 2006):

$$\bar{I}_4 = \kappa \left(\bar{I}_1 - 3 \right) + (1 - 3\kappa) \left(\bar{I}_4 - 3 \right)$$

Dispersion parameter $\kappa \in [0, 1/3]$:

$\kappa = 0$: Perfectly aligned fibers (pure transverse anisotropy).

$\kappa = 1/3$: Fully isotropic dispersion.

$\kappa = 0.09$ (adopted): Strong preferential alignment with significant physical dispersion.

A.4 Associated Stress Tensors

A.4.1 Second Piola-Kirchhoff Stress ($\mathbf{S}$)

$$\mathbf{S} = 2 \frac{\partial \Psi}{\partial \mathbf{C}} = \frac{\partial \Psi_{\text{vol}}}{\partial \mathbf{C}} + \frac{\partial \Psi_{\text{iso}}}{\partial \mathbf{C}}$$

Volumetric: $\mathbf{S}_{\text{vol}} = k J (J - 1) \mathbf{C}^{-1}$

Isochoric:

$$\mathbf{S}_{\text{iso}} = J^{-2/3} \left[\bar{\mathbf{C}} - \frac{1}{3} \bar{I}_1 \bar{\mathbf{C}} \right] + \frac{1}{J} \left(\bar{\mathbf{C}} - \frac{1}{3} \bar{I}_1 \bar{\mathbf{C}} \right) \mathbf{C}^{-1}$$

where:

$$\frac{\partial \Psi_{\text{iso}}}{\partial \bar{I}_1} = c + \frac{1}{2} \sum_{i=1}^2 \kappa \left(\bar{I}_4 - 3 \right) \exp \left(\frac{1}{2} \left(\bar{I}_4 - 3 \right)^2 \right)$$

$$\frac{\partial \Psi_{\text{iso}}}{\partial \bar{I}_4} = \frac{1}{2} (1 - 3\kappa) \exp \left(\frac{1}{2} \left(\bar{I}_4 - 3 \right)^2 \right)$$

A.4.2 Cauchy Stress Tensor ($\boldsymbol{\sigma}$)

$$\boldsymbol{\sigma} = J^{-1} \mathbf{F} \mathbf{S} \mathbf{F}^T$$

Volumetric: $\boldsymbol{\sigma}_{\text{vol}} = k (J - 1) \mathbf{I} = -p \mathbf{I}$

Isochoric: $\boldsymbol{\sigma}_{\text{iso}} = J^{-1} \left(\bar{\boldsymbol{\sigma}} - \frac{1}{3} \bar{I}_1 \bar{\boldsymbol{\sigma}} \right)$

A.5 Physical Meaning and Parameter Calibration

Parameter	Symbol	Canonical Value	Physical Meaning	Clinical Role
Matrix shear modulus	c	0.05 MPa	Ground substance (GAGs, proteoglycans) elasticity	**Dominant** parameter; degradation initiates ectasia
Collagen fibril stiffness	k ₁	0.22 MPa	Intrinsic elastic stiffness of collagen fibrils	Primary load-bearing barrier; CXL increases 3–5×
Fiber nonlinearity	k ₂	100	Rate of exponential stiffening	Prevents catastrophic deformation under IOP spikes
Fiber dispersion	κ	0.09	Angular dispersion around preferred direction	κ → 1/3 in advanced cones → loss of anisotropy
Bulk modulus	k	4.76 MPa	Resistance to volumetric compression	Enforces near-incompressibility (ν ≈ 0.49)

References

Gasser TC, Ogden RW, Holzapfel GA. *J Royal Soc Interface*. 2006;3(6):15–35.

Holzapfel GA, Gasser TC, Ogden RW. *J Elasticity*. 2000;61(1–3):1–48.

Nguyen BA, Roberts CJ, Reilly MA. *Front Bioeng Biotechnol*. 2018;6:210.

Pandolfi A, Manganiello F. *Biomech Model Mechanobiol*. 2006;5(4):237–246.

Meek KM, Knupp C. *Prog Retin Eye Res*. 2015;49:1–16.

Appendix B — FEBio Scripts and Extraction Pipeline Documentation

B.1 Computational Pipeline Overview

The AVBC computational pipeline consists of four interlinked Python scripts that automate generation, execution, extraction, and analysis of finite element simulations. All scripts are located in the `simulations/` directory.

`simulations/`

```
├── generate_febio_model.py      # .feb model generator
├── run_parametric_campaign.py   # Parametric campaign orchestrator
├── extract_parametric_vectors.py # Vector metric extraction
├── analyze_phenotype_discrimination.py # Phenotypic discrimination analysis
├── calc_curvature.py           # Curvature calculation from displacements
├── models/                     # Generated .feb files
├── results/                    # Simulation outputs
└── analysis/                   # Discrimination reports
```

Sequential workflow:

Generate → `generate_febio_model.py` creates `.feb` files with configurable HGO parameters and ICRS geometry

Execute → `run_parametric_campaign.py` orchestrates parallel execution on FEBio 4.12

Extract → `extract_parametric_vectors.py` reads output files and calculates VR, VT, $V\tau$

Analyze → `analyze_phenotype_discrimination.py` performs sensitivity and discrimination analysis

B.2 Model Generator: `generate_febio_model.py`

Mesh Specification

Parameter	Value	Description
Radius of curvature (R)	7.8 mm	Mean corneal radius
Angular extent (θ_{max})	46.5°	Apex to limbus
Radial rings	20	Apex (ring 0) to limbus (ring 20)
Circumferential sectors	24	15° each
Node layers	2	Posterior ($z=0$) and anterior ($z=1$)
Total nodes	962	$20 \times 24 \times 2 + 2$ (apical nodes)
Pentahedral elements (apex)	24	Penta6 to avoid singularity
Hexahedral elements (body)	456	Linear Hex8
Total elements	**480**	
Stromal thickness (t_0)	500 μm	Uniform (canonical model)

ICRS Positioning

Ring 14 ($r \approx 4.2$ mm from apex), corresponding to the 5–6 mm optical zone.

Boundary Conditions

Follower pressure: 15 mmHg (0.002 MPa) on posterior face.

Limbal fixation: All DOFs fixed at ring 20.

ICRS prescribed displacement: +250 μm anteriorly (z), free in x and y .

Usage

```
python generate_febio_model.py \
--c 0.05 --k1 0.22 --k2 100 --kappa 0.09 \
--arc 160 --thickness 250 \
--output model_arc160_t250.feb
```

Flag	Parameter	Default	Unit
<code>--c</code>	Matrix stiffness	0.05	MPa
<code>--k1</code>	Fibrillar stiffness	0.22	MPa
<code>--k2</code>	Nonlinearity	100	—
<code>--kappa</code>	Dispersion	0.09	—
<code>--arc</code>	Arc length	160	degrees
<code>--thickness</code>	ICRS thickness	250	μm
<code>--progressive</code>	Progressive profile	—	linear/parabolic
<code>--output</code>	Output file	model.feb	—

B.3 Parametric Campaign: `run_parametric_campaign.py`

Experimental Design (OAT Factorial)

Material variations (20 configurations):

c : 0.01 / 0.02 / 0.03 / 0.05 / 0.07 / 0.10 MPa (6 values)

k_1 : 0.05 / 0.10 / 0.22 / 0.30 / 0.50 MPa (5 values)

k_2 : 10 / 50 / 100 / 200 / 500 (5 values)

κ : 0.00 / 0.05 / 0.09 / 0.15 / 0.333 (5 values, including isotropic)

Geometric variations (34 configurations):

Arcs: 90° / 120° / 160° / 210° / 255° / 290° / 320° / 360°

Asymmetric: 150→300 / 200→300 / 300→150 / 300→200 μm , linear and parabolic

Pachymetry: 400 / 500 / 600 μm

Total: 377 simulations converged from 680 attempted (convergence rate: 55.4%).

```
python run_parametric_campaign.py --jobs 4      # Full campaign
python run_parametric_campaign.py --subset geometry --jobs 2 # Geometry only
python run_parametric_campaign.py --dry-run      # Generate .feb without executing
```

B.4 Vector Extraction: `extract_parametric_vectors.py`

Extracted Metrics

Metric	Symbol	Description	AVBC Vector
Apical displacement	δ_{apex}	uz displacement at node 1 (apex)	VR
Radial displacement	δ_{ring}	Mean radial displacement at ICRS ring	VT
Keratometric change	ΔK	Refractive power change (D)	VR
Circumferential strain	ϵ_{circ}	Mean tangential strain at ring	VT

Maximum stress	$\sigma_{\max}$	Maximum von Mises stress (MPa)	—
Asymmetry index	AI	Displacement field asymmetry	$V\tau$
Flattest meridian	θ_{flat}	Direction of maximum flattening (°)	—
Steepest meridian	θ_{steep}	Direction of maximum curvature (°)	—

Curvature Calculation (ΔK)

$$\Delta K = \frac{n_{\text{ref}} - 1}{R_{\text{post}}} - \frac{n_{\text{ref}} - 1}{R_{\text{pre}}}$$

where $n_{\text{ref}} = 1.3375$ and R is fitted to a local sphere over the central 3 mm.

B.5 Discrimination Analysis: `analyze\_phenotype\_discrimination.py`

Canonical Results

Parameter	CV of δ_{apex}	Variance Explained
c (matrix)	**0.671**	**98.9%**
κ (dispersion)	0.011	1.1%
k_1 (fibers)	0.000	< 0.02%
k_2 (stiffening)	0.000	0.0%

B.6 Software Requirements

Component	Version	Source
FEBio	4.12	febio.org
Python	≥ 3.8	—
NumPy	≥ 1.20	`pip install numpy`
Pandas	≥ 1.3	`pip install pandas`
SciPy	≥ 1.7	`pip install scipy`

Full Pipeline Execution

```
python run_parametric_campaign.py --dry-run --jobs 4 # Generate models
python run_parametric_campaign.py --jobs 4 # Execute campaign
python extract_parametric_vectors.py --campaign results/campaign_summary.json
python analyze_phenotype_discrimination.py
```

B.7 Reproducibility

All input files (`.feb`), binary outputs (`.xplt`), solver logs, and data extractions are versioned. Scripts are deterministic: given identical input parameters, the same `.feb` file is generated byte-for-byte. Random seed is fixed at `seed=42`.

Appendix C — AVBC Terminology Glossary

Biomechanical Terms

Flattening (*Aplanamento*)

Reduction of central corneal curvature, measured as decrease in K-steep or K-mean. In the AVBC framework, flattening is mediated by the VR vector and controlled by total implant volume (thickness $\times$ arc).

Anisotropy

Material property where mechanical behavior varies with direction. The corneal stroma is anisotropic: stiffer along nasal-temporal and superior-inferior meridians (cardinal fibers) than in oblique directions.

AVBC — Corneal Biomechanical Vector Analysis

Surgical planning framework for ICRS that decomposes the biomechanical ring effect into three independent vectors (VR, VT, $V\tau$), replacing empirical nomogram-based selection.

Hoop restriction (*Cintagem*)

Circumferential restriction effect exerted by the ICRS on the peripheral stroma. Analogous to a barrel hoop: prevents radial expansion under IOP. Governs the VT vector and is controlled by arc length.

CXL — Corneal Collagen Crosslinking

Procedure that reinforces interfibrillar collagen cross-links using riboflavin and UVA radiation. Increases the HGO model parameter k_1 (fibrillar stiffness).

Apical displacement (δ_{apex})

Vertical displacement of the corneal apex under intraocular pressure. Primary deformation metric in FEM simulations. Canonical baseline value: 360.9 μm (without ring).

Mechanical Neutral Axis (MNA) (*Eixo Neutro Mecânico, ENM*)

Axis dividing the cornea into two halves with approximately equal biomechanical properties, determined by posterior elevation and minimum thickness. Diverges from the steepest curvature axis (K-steep) in up to 40% of cases. ICRS should be positioned relative to the MNA, not K-steep.

Volumetric School (*Escola Volumétrica*)

Paradigm explaining the ICRS effect through volume displacement: the ring occupies physical volume in the incompressible stroma, forcing adjacent tissue to displace radially and anteriorly. Foundation: Kling & Marcos (IOVS, 2013).

FEM — Finite Element Method

Computational technique for solving partial differential equations in complex geometries. In the AVBC context, used to simulate corneal response to ICRS insertion.

FEBio

Open-source finite element software (University of Utah) specialized in biomechanics. Version used: FEBio 4.12. Natively supports HGO constitutive model.

HGO — Holzapfel–Gasser–Ogden Model

Anisotropic hyperelastic constitutive model for fiber-reinforced materials. Decomposes strain energy into isotropic matrix contribution (c) and anisotropic fiber contributions (k_1 , k_2 , κ). Reference: Holzapfel, Gasser & Ogden (*J Elasticity*, 2000).

Hoop stress — Tangential stress

Circumferential stress in a pressurized shell (like the cornea under IOP). Governs the tendency for radial expansion. ICRS interrupts this stress by creating a mechanical barrier (see Hoop restriction).

ICRS — Intrastromal Corneal Ring Segment

PMMA implant inserted into the corneal stroma to modify curvature, astigmatism, and aberrations. Commercial types: Intacs, Ferrara Ring, Keraring.

Incompressibility

Material property of resistance to volume change. The corneal stroma (78% water) is nearly incompressible: when the ICRS occupies volume, adjacent tissue must displace, not compress.

HGO Constitutive Parameters

Parameter	Symbol	Canonical Value	Clinical Meaning
Matrix stiffness	c	0.05 MPa	Proteoglycan degradation → keratoconus. **Dominant parameter.**
Fibrillar stiffness	k_1	0.22 MPa	Collagen weakening; CXL increases k_1 .
Exponential stiffening	k_2	100	Nonlinear fiber response to large deformations.
Fiber dispersion	α	0.09	Lamellar disorganization; $\alpha \rightarrow 1/3$ = total isotropy.
Bulk modulus	k	4.76 MPa	Enforces near-incompressibility.

AVBC Vectors

VR — Radial Displacement Vector

Quantifies the flattening effect. Controlled by ring thickness (injected volume). Insensitive to arc length in partial arcs. FEM range: 19.2–19.9 μm (partial arcs), 125.9 μm (full 360° ring).

VT — Tangential Stress Vector

Quantifies astigmatism redistribution through circumferential belting. Controlled by arc length. Empirical equation: $VT = -0.0018 \times \text{arc}^\circ + 7.79$ ($R^2 = 0.94$). FEM range: 7.20–7.78 kPa.

V τ — Asymmetric Torque Vector

Quantifies apex repositioning. Controlled by thickness differential in progressive rings. Numerical zero for symmetric rings. FEM range: 9.31 $\mu\text{N}\cdot\text{m}$ (linear progressive) to 18.34 $\mu\text{N}\cdot\text{m}$ (parabolic progressive).

Trimodal Classification (O/T/B Modules)

O Module — Optical Coherence Assessment

Classifies visual recovery potential: O+ (BCVA $\geq 20/60$, coma $< 2.5 \mu\text{m}$) or O– (BCVA $< 20/60$ or coma $> 3.5 \mu\text{m}$).

T Module — Topographic Morphology

Classifies the spatial cone configuration: Central, Crescent (paracentral), Peripheral (decentered), or Global (diffuse).

B Module — Biomechanical Selection

Translates O+T classification into ring prescription using the three vectors. Determines the dominant vector and corresponding parameters.

AVBC Correction Indices

CI_R — Radial Correction Index

$CI_R = \Delta K_{\text{observed}} / \Delta K_{\text{predicted}}$. Target: 1.0 ± 0.15 .

CI_T — Tangential Correction Index

$CI_T = \Delta Cyl_{\text{observed}} / \Delta Cyl_{\text{predicted}}$. Target: 1.0 ± 0.15 .

ACE — Axial Coherence Index

ACE_{min} = minimum angle between steepest curvature axis and MNA. Screening threshold: ACE_{min} $< 28^\circ$ predicts ≥ 3 lines of gain (AUC 0.82).

Acronyms

Acronym	Meaning
AVBC	Corneal Biomechanical Vector Analysis
AE	Angle of Error (Alpins)
BCVA	Best Corrected Visual Acuity
CI	Correction Index
CXL	Corneal Collagen Crosslinking
DV	Difference Vector (Alpins)
FEM	Finite Element Method
HGO	Holzapfel–Gasser–Ogden
HOA	Higher-Order Aberrations
ICRS	Intrastromal Corneal Ring Segment
IOP	Intraocular Pressure
MAE	Mean Absolute Error
MNA	Mechanical Neutral Axis
ORA	Ocular Response Analyzer
PMMA	Polymethylmethacrylate
SIA	Surgically Induced Astigmatism (Alpins)
TIA	Target Induced Astigmatism (Alpins)
UCVA	Uncorrected Visual Acuity
VR	Radial Displacement Vector
VT	Tangential Stress Vector
V τ	Asymmetric Torque Vector
WAXS	Wide-Angle X-ray Scattering