

Foundation (Jalapeno)

- Q1.** What is the mean of 2, 4, 6?
(A) 2
(B) 4
(C) 6
(D) 3
(E) 12
- Q2.** Which of the following is a measure of central tendency?
(A) Range
(B) Mean
(C) Variance
(D) Standard Deviation
(E) Interquartile Range
- Q3.** A set of data has a median of 10. What does this mean?
(A) The mean is 10
(B) The mode is 10
(C) The middle value is 10 when the data is arranged in ascending order
(D) The range is 10
(E) The standard deviation is 10
- Q4.** What is the mode of 1, 2, 2, 3, 4, 4, 4?
(A) 1
(B) 2
(C) 3
(D) 4
(E) 5
- Q5.** A dataset has the following values: 1, 3, 5, 7, 9. What is the median?
- (A) 1
(B) 3
(C) 5
(D) 7
(E) 9
- Q6.** What is the mean of the numbers 10, 20, 30, 40, 50?
(A) 20
(B) 25
(C) 30
(D) 35
(E) 40
- Q7.** If a dataset has a mean of 15 and a median of 20, what can be said about the dataset?
(A) The dataset is symmetric
(B) The dataset is skewed to the left
(C) The dataset is skewed to the right
(D) The mean and median are equal
(E) The mode is 15
- Q8.** Which measure of central tendency is most affected by outliers?
(A) Mean
(B) Median
(C) Mode
(D) Range
(E) Standard Deviation

Open Ended Questions (FRQ)

- Q9.** Calculate the mean, median, and mode of the dataset: 2, 4, 6, 8, 10. Show all work and explain your answers.
- Q10.** Compare the measures of central tendency for two datasets: 1, 3, 5 and 2, 4, 6. Which dataset has the higher mean? Explain your reasoning.

Chef's Tips

- Tip 1: Ensure students have a calculator for the test
- Tip 2: Allocate 60 minutes to complete the test
- Tip 3: Remind students to show all work for free response questions