

Foundation (Jalapeno)

Q1. Solve for x : $2x + 5 = 11$

- (A) $x = 2$
- (B) $x = 3$
- (C) $x = 4$
- (D) $x = 5$
- (E) $x = 6$

Q2. Solve for y : $y - 3 = 7$

- (A) $y = 4$
- (B) $y = 5$
- (C) $y = 10$
- (D) $y = 11$
- (E) $y = 12$

Q3. Solve for z : $z + 2 = 9$

- (A) $z = 5$
- (B) $z = 7$
- (C) $z = 9$
- (D) $z = 11$
- (E) $z = 13$

Q4. Solve for x : $x/4 = 9$

- (A) $x = 36$
- (B) $x = 40$
- (C) $x = 45$
- (D) $x = 50$
- (E) $x = 55$

Q5. Solve for y : $3y = 24$

- (A) $y = 6$
- (B) $y = 7$
- (C) $y = 8$
- (D) $y = 9$
- (E) $y = 10$

Q6. Solve for x : $x - 2 = 7$

- (A) $x = 5$
- (B) $x = 9$
- (C) $x = 10$
- (D) $x = 11$
- (E) $x = 12$

Q7. Solve for z : $z/3 = 4$

- (A) $z = 10$
- (B) $z = 12$
- (C) $z = 15$
- (D) $z = 16$
- (E) $z = 20$

Q8. Solve for x : $5x = 35$

- (A) $x = 5$
- (B) $x = 6$
- (C) $x = 7$
- (D) $x = 8$
- (E) $x = 9$

Open Ended Questions (FRQ)

Q9. Solve for x : $2x + 5 = 11$ and $x - 2 = 7$. Show your work.

Q10. Rearrange the formula $C = 2\pi r$ to isolate r . Then solve for r when $C = 12\pi$.

Chef's Tips

- Provide a formula sheet with common algebra formulas.
- Allow students to use calculators for checking.
- Review literal equations and variable isolation before the test.