

Historic effective population
size

Effective Population size


- Effective population size:
 - the size of a population that shows the same amount of dispersion of allele frequencies under drift as an ideal population
- Effective population size is important for the conservation of endangered species
 - indicates the effective number of breeders in a population
 - Often lower than census population size $N_e = 0.2 * N_c$
 - If N_e is low there is a high chance of inbreeding, reducing N_e further (-> extinction vortex)
- Genetics allow us to estimate N_e with a single sample for current population sizes (NeEstimator)
- Genomics allow us to estimate N_e with a single sample for historic population sizes (<- but how good are they)

Methods

- MSMC, PMSC
 - Need whole genome sequences for each individual
- Stairways2, Epos, Snep (using the site frequency spectrum)
 - Using SNPs only
 - Based on the Coalescent approach
- Gone, LinkNe
 - Using linkage disequilibrium
 - Using SNPs and a linkage map (needs a good reference genome)

Coalescent (Kingsman 1980)

- IBD (identical by descent) [not isolation by distance!!!]
- Inheritance of a chromosome from a great parent
- For the two cousins the overlapping segment is said to have coalesced in their great-grandfather. (IBD)




(from Pritchards new book)

[<https://web.stanford.edu/group/pritchardlab/HGbook/>]

Coalescent (Kingsman 1980)

- IBD (identical by descent) [not isolation by distance!!!]
- Inheritance of a chromosome from a great parent
- For the two cousins the overlapping segment is said to have coalesced in their great-grandfather. (IBD)




(from Pritchards new book)

[<https://web.stanford.edu/group/pritchardlab/HGbook/>]

Coalescent (Kingsman 1980)

Backwards in time


- Two copies of a locus in the present generation are marked by red balls.
- These descend from a common ancestor (i.e. they coalesce) two generations ago. In coalescent models it is most natural to measure time backward from the present.




(from Pritchards new book)

[<https://web.stanford.edu/group/pritchardlab/HGbook/>]


Coalescent (Kingsman 1980)



WF genealogy for a small population.
six chromosomes sampled at the present day, in red.



Red circles and arrows indicate the ancestors of the sampled chromosomes



The coalescent genealogy abstracts away all irrelevant details of the WF process, showing only the ancestral relationships of the 6 samples and the coalescent times.

(from Pritchards new book)

[<https://web.stanford.edu/group/pritchardlab/HGbook/>]

Coalescent (Kingsman 1980)

Lots of calculation can be done (probability to coalesce in t generation) $(1 - \frac{1}{2N})^t$.

Most recent common ancestor is: $4N_e$

We now add mutations in the mix

Coalescent (Kingsman 1980)

Lots of calculation can be done (probability to coalesce in t generation) $(1 - \frac{1}{2N})^t$.

Most recent common ancestor is: $4Ne$

We now add mutations in the mix SFS





Figure 1

The locations of mutations on the coalescent genealogy (*left*) give rise to patterns of genetic variation data (*top right*). The SFS depicts the mutational patterns seen in the genetic variation data. Abbreviations: SFS, site frequency spectrum; SNPs, single nucleotide polymorphisms.


Site frequency spectrum

- Population history influences the shape of genealogies and the SFS.
- The yellow shaded areas on the left denote the history of each population.
- These demographic histories give rise to the genealogies shown within each model. Blue stars denote mutations that occur on the genealogies.
- The histograms denote the SFS for each model that is generated from the mutational pattern that occurred on the genealogy.







Site frequency spectrum

- A more realistic SFS
- ($N_e=100$, constant last 400 years)



Site frequency spectrum

- SFS for different population trajectories



Epos and Stairways





- Aim to find the historic population trajectories by fitting the SFS
- Basically all possible SFS are created and compared to the actual one
- Methods that simulate certain histories (Fastsimcoal, dadi) [testing certain hypothesis]
- Methods that do not need a proposal history: Epos/Stairways[/Gone]

Input for Epos/Stairways2

- SNPs (genlight object)
- L, mu and SFS (minbinsize)
- gl.sfs creates a SFS based on dartR/genlight object
- `> gl.sfs(gl, minbinsize = 1, folded=TRUE, singlepop=TRUE)`
- Exercise 1: Parameter in gl.sfs
- Exercise 2: Compare sfs for different data sets
(glcon, glinc, gldec, glbottle)



Input for Epos/Stairways2

- `> gl.sfs(gl, minbinsize = X, folded=TRUE, singlepop=TRUE)`



Input for Epos/Stairways2

- `> gl.sfs(gl, minbinsize = 1, folded=TRUE/FALSE)`



Input for Epos/Stairways2

- L: the length of all combined sequences (genome length)
- mu: mutation rate


$$x = \mu \times L \times 2 \times N_e$$

Principle: x: number of mutations in a generation

So the “trajectory” is calibrated for this product





- Exercise 3: Run gl.epos for different settings of L and mu

Epos




Epos

- Exercise 4: Run Epos for all four data sets







Stairways2

- Stairways2 (has the same input), but takes >50 times longer or so you can repeat the same exercise for all four scenarios.



Stairways [all four scenarios]





Isobel Walcott (Honours) Demographic Inference in a Conservation Context

- Standard cases – decline, expansion, stable
 - 586 scenario combinations, 30 replicates
 - 6 levels of loci subsampling for each
 - 3 methods tested
 - 316,440 runs (using NCI infrastructure)
- Bottleneck cases – based on known decline-recovery cases
 - Saltwater crocodiles, southern right whales, fur seals, Fleay's barred frog




BEER ACCOUNT

- Bernd Gruber
- 062 924
- 10173614

- If you only drink soft drinks it is \$10
- \$20 for alcoholic drinks
- Talk to Jason in case you want something special

- Board games at 7:00

Stairway2 and Epos Example runs



- 200 individuals
- 20k loci
- Decline: Population crash from 500 to 250 over 30 years
- Expansion: Population increase from 250 to 500 over 30 years
- Stable: Population of 500

Crocodiles

Population history

- Up until the mid-20th century, saltwater crocodiles were abundant across northern Australia.
- unregulated hunting and habitat destruction led to a dramatic decline in their numbers (~1950 – 1970)
- The saltwater crocodile was fully protected under Australian law by 1971.
- conservation efforts included habitat protection, regulated farming, and sustainable use initiatives, which have been pivotal in the species' recovery (>100.000)
- length of genome: $l=2.123e9$ (or
- length of $\mu = 7.9e-9$ (slowest found in reptiles)


DOI: 10.1002/jwmg.22525

RESEARCH ARTICLE



Natal origin and dispersal of problem saltwater crocodiles in the Darwin Harbor, Australia


Yusuke Fukuda^{1,2} | Craig Moritz² | Nancy N. FitzSimmons³ |
Namchul Jang⁴ | Grahame Webb⁵ | Garry Lindner⁶ |
Hamish Campbell⁷ | Keith Christian⁷ | Steven Leeder⁸ |
Sam Banks⁷



```
> nLoc(crocs)
[1] 1602
> nInd(crocs)
[1] 497
> gl.sfs(crocs, singlepop = TRUE)
```


Crocodiles

- sfs





Crocodiles

- trajectory



Foxes


- 160 foxes, 21259 loci
- SFS



Foxes

- 160 foxes, 21259 loci

- SFS




LD methods


- Linkage disequilibrium
- Hayes 2003 (CSH phased) and LD (for phased and unphased data)
- LD (CSH) along; $LD = (1 + 4N_e \times c)$
- LD changes over distances along the chromosomes and number of recombination are determined by effective population size. LD over distance
- Recombination rate c is unknown (using distance between as a proxy)

Linkage (Dis)equilibrium

Linked genes on a pair of homologous chromosomes:




Replication takes place at the beginning of meiosis:



The homologous chromosomes undergo synapsis and crossover occurs between adjacent chromatids:



The chromatids separate:



Now four different kinds of gametes form

Ne and LD

- ***Ne***: the number of breeding individuals in an idealized population that would show the same amount of dispersion of allele frequencies under random genetic drift
- $E(\hat{r}^2)$: LD [Linkage disequilibrium]
- S : sample size (number of individuals)
- c : recombination rate (for all pairs of SNPs)

$$E(\hat{r}_{\Delta}^2) \approx \frac{(1-c)^2 + c^2}{2N_e c(2-c)} + \frac{1}{S},$$

Ne and LD

- **Ne**: the number of breeding individuals in an idealized population that would show the same amount of dispersion of allele frequencies under random genetic drift
- $E(\hat{r}^2)$: LD [Linkage disequilibrium]
- S : sample size (number of individuals)
- c : recombination rate (for all pairs of SNPs)

$$E(\hat{r}_{\Delta}^2) \approx \frac{(1-c)^2 + c^2}{2N_e c(2-c)} + \frac{1}{S},$$

If $c=0.5$:

$$E(\hat{r}_{\Delta}^2) \approx \frac{1}{3N_e} + \frac{1}{S}$$

LD

- LD between pairs of SNP at binned distances provide information on population size at times in the past
- Originally only for expanding and declining population
- New methods eg GONE using phased, unphased genotypes)
 - Input SNPs + linkage map (position of SNPs on the chromosomes)


Recent Demographic History Inferred by High-Resolution Analysis of Linkage Disequilibrium

Enrique Santiago,^{*1} Irene Novo,² Antonio F. Pardiñas,³ María Saura,⁴ Jinliang Wang,⁵ and Armando Caballero²

Gone

- Comparison to other methods
- Number of SNPs: 255,000 or 450,000
- Number of samples: 20 (4 MSMC)

Gone —
MSMC —
Relate —
LinkNe —



Gone

- Very data hungry
- $N_e=1000$ and 100 individuals sampled.
- Implemented to work with dartR, but you need a linkage map




FIG. 2. Estimates of temporal N_e by GONE under different simulated demographic scenarios from present (generation 0) to 220 generations in the past. The true population size is the green shadowed area and n is the sample size of individuals for analysis. For all panels, the black lines refer to an analysis where all recombination bins from $c = 0.001$ up to $c = 0.5$ are considered (option $hc = 0.5$), whereas the red lines refer to analyses with rate bins from $c = 0.001$ up to only 0.05 ($hc = 0.05$). (A) and (B) Detection of bottlenecks occurring at different times. (C) Scenario with overlapping generations with three cohorts per generation and mixed-cohort sampling. (D) A population $N_e = 1,000$ was divided into two populations $N_e = 1,000$ each, which were isolated for 100 generations and then mixed 50 generations ago into a single population with $N_e = 1,000$. (E) and (F) Metapopulation composed of two subpopulations $N_e = 1,000$ each with 2% and 0.2% of migration, respectively, between them. (G) Estimations under different base-calling error rates. From top to bottom, 10%, 1%, 0.1% and 0%, the latter two being indistinguishable. (H) A hundred individuals were sampled from the population over a period of 100 consecutive generations at a rate of one sampled individual per generation. (I) and (J) Eight small samples ($n = 10$ each) were taken from the same population at the same time.


DnaDot - Fixing Ecology and Evolution's Blind Spot, Population size N_c

Ecological Indicators 2024

WB Sherwin EERC UNSW-Sydney

W.Sherwin@unsw.edu.au

N_c from: Mark Recapture **MR**;
Close Kin Mark Recapture **CKMR**



N_c is **CRUCIAL** – abundance needed for:

- $\pm 10\%$ accuracy needed for any forecasting, eg IUCN listing.
- Biodiversity measures with narrow confidence limits
(Simpson, Shannon, **NOT** species lists)
- In most Ecology texts, $\sim 75\%$ of chapters rely heavily on N_c ,
for mechanics or outcomes of competition, predation, etc.

N_c from: Mark Recapture **MR**;
Close Kin Mark Recapture **CKMR**

IMPERFECT MR:

- Sample sizes (for $\pm 10\%$ precision, need 80% true N_c !!!)
- 2+ samples
- Assume: no birth death immigration emigration

IMPERFECT CKMR:

- Sample size $\sqrt{\text{true } N_c}$ only gives $\pm 30\%$ accuracy – not $\pm 10\%$
- Individual or kin identification, often from worst possible DNA!
- Assume: family size mean and variance known independently

DnaDot

HOW:

- Based on MR, but no marking
- Pre-existing polymorphisms 'mark' separate groups in population
(eg, at this SNP, these ones have 'T', these ones have 'C')

ADVANTAGES:

- Single sample
- No need for independent knowledge of birth, death, immigration, emigration, family size mean and variance, etc
- Genotyping only good enough to estimate allele proportions, NOT to identify individuals and kin

DnaDot performs well (accurate & precise)

	Bias Estimate = $\pm 10\%$ of true N_c or better	Variability CL Confidence limits = $\pm 10\%$ or better
MR	No (1/3) Maybe (1/3) Yes (1/3)	No (4/5) Maybe (1/5)
CKMR	No (2/2)	No (4/4)
DnaDot	No (6/36) Yes (30/36)	Yes (36/36)

DnaDot Summary

Compared to older genetic and non-genetic measures:

- DnaDot avoids most pitfalls of previous estimates of population size N_c
- DnaDot is sufficiently accurate and precise for most uses of N_c
- Article has link to App requiring Excel input and output, no programming: Sherwin WB. 2024. DnaDot - Fixing Ecology and Evolution's Blind Spot, Population size. Ecological Indicators *****
- WB Sherwin EERC UNSW-Sydney W.Sherwin@unsw.edu.au

Coalescent effective population size


- **Estimating Historical Population Sizes:** Coalescent models can use genetic data to estimate changes in N_e over time, providing insights into past population dynamics.
- In coalescent models, N_e influences the expected time until two lineages coalesce.
- Show different SFS (based on different trajectories)
- Explain the coalescent in sketches
- Explain N_e (in this sense)
- Run different scenarios (explain impact of L , μ)
- Show some examples and how users can use their data (preparation)

DnaDot - Fixing Ecology and Evolution's Blind Spot, Population size N_c

Ecological Indicators 2024

WB Sherwin EERC UNSW-Sydney
W.Sherwin@unsw.edu.au

N_c from: Mark Recapture **MR**;
Close Kin Mark Recapture **CKMR**



N_c is **CRUCIAL** – abundance needed for:

- $\pm 10\%$ accuracy needed for any forecasting, eg IUCN listing.
- Biodiversity measures with narrow confidence limits
(Simpson, Shannon, **NOT** species lists)
- In most Ecology texts, $\sim 75\%$ of chapters rely heavily on N_c ,
for mechanics or outcomes of competition, predation, etc.

N_c from: Mark Recapture **MR**;
Close Kin Mark Recapture **CKMR**

IMPERFECT MR:

- Sample sizes (for $\pm 10\%$ precision, need 80% true N_c !!!)
- 2+ samples
- Assume: no birth death immigration emigration

IMPERFECT CKMR:

- Sample size $\sqrt{\text{true } N_c}$ only gives $\pm 30\%$ accuracy – not $\pm 10\%$
- Individual or kin identification, often from worst possible DNA!
- Assume: family size mean and variance known independently

DnaDot

HOW:

- Based on MR, but no marking
- Pre-existing polymorphisms 'mark' separate groups in population
(eg, at this SNP, these ones have 'T', these ones have 'C')

ADVANTAGES:

- Single sample
- No need for independent knowledge of birth, death, immigration, emigration, family size mean and variance, etc
- Genotyping only good enough to estimate allele proportions, NOT to identify individuals and kin

DnaDot performs well (accurate & precise)

	Bias Estimate = $\pm 10\%$ of true N_c or better	Variability CL Confidence limits = $\pm 10\%$ or better
MR	No (1/3) Maybe (1/3) Yes (1/3)	No (4/5) Maybe (1/5)
CKMR	No (2/2)	No (4/4)
DnaDot	No (6/36) Yes (30/36)	Yes (36/36)

DnaDot Summary

Compared to older genetic and non-genetic measures:

- DnaDot avoids most pitfalls of previous estimates of population size N_c
- DnaDot is sufficiently accurate and precise for most uses of N_c
- Article has link to App requiring Excel input and output, no programming: Sherwin WB. 2024. DnaDot - Fixing Ecology and Evolution's Blind Spot, Population size. Ecological Indicators *****
- WB Sherwin EERC UNSW-Sydney W.Sherwin@unsw.edu.au