

RWorksheet__Sanceda#4A

2024-10-14

#1.

#a. The data displays the Shoe Size, Height, and Gender.

```
ShoeSize <- c(6.5,9.0,8.5,8.5,10.5,7.0,9.5,9.0,13.0,7.5,10.5,8.5,12.0,10.5,13.0,11.5,8.5,5.0,10.0,6
```

```
Height <-c(66.0,68.0,64.5,65.0,70.0,64.0,70.0,71.0,72.0,64.0,74.5,67.0,71.0,71.0,77.0,72.0,59.0,62.0)
```

```
Gender <- c("F", "F", "F", "F", "M", "F", "F", "F", "M", "F", "M", "F", "M", "M", "M", "M", "F", "F", "M", "F", "F", "M")
```

```
Shoes_Tble <- data.frame(ShoeSize, Height, Gender)
```

Shoes_Tble

```
##      ShoeSize Height Gender
```

##	1	6.5	66.0	F
----	---	-----	------	---

```
## 2      9.0    68.0      F
```

```
## 3      8.5    64.5      F
```

##	4	8.5	65.0	F
----	---	-----	------	---

```
## 5      10.5    70.0      M
```

##	6	7.0	64.0	F
----	---	-----	------	---

##	7	9.5	70.0	F
----	---	-----	------	---

## 8	9.0	71.0	F
------	-----	------	---

##	9	13.0	72.0	M
----	---	------	------	---

##	9	18.0	72.0	H
##	10	7.5	64.0	F

##	10	1.0	91.0	I
##	11	10.5	74.5	M

##	11	10.5	74.5	M
##	12	8.5	67.0	F

##	12	8.5	67.0	F
##	13	12.0	71.0	M

##	13	12.0	71.0	M
##	14	10.5	71.0	M

##	14	10.5	71.0	M
##	15	13.0	77.0	M

##	15	13.0	77.0	M
##	16	11.5	72.0	M

##	16	11.5	72.0	M
##	17	8.5	50.0	F

##	17	8.5	59.0	F
##	18	5.0	60.0	F

##	18	5.0	62.0	F
##	19	10.0	72.0	M

##	19	10.0	72.0	M
##	20	0.5	33.0	F

##	20	6.5	66.0	F
----	----	-----	------	---

##	21	7.5	64.0	F
----	----	-----	------	---

```
## 22      8.5    67.0      M
```

```
## 23      10.5    73.0      M
```

```
## 24      8.5    69.0      F
```

```
## 25      10.5    72.0      M
```

```
## 26      11.0    70.0      M
```

```
## 27      9.0    69.0      M
```

```
## 28      13.0    70.0      M
```

#b.

```
Male <- subset(Shoes_Tble, Gender == "M", select = c(ShoeSize, Gender, Height))
Male
```

```
##      ShoeSize Gender Height
## 5         10.5      M   70.0
## 9         13.0      M   72.0
## 11        10.5      M   74.5
## 13         12.0      M   71.0
## 14         10.5      M   71.0
## 15         13.0      M   77.0
## 16         11.5      M   72.0
## 19         10.0      M   72.0
## 22          8.5      M   67.0
## 23         10.5      M   73.0
## 25         10.5      M   72.0
## 26         11.0      M   70.0
## 27          9.0      M   69.0
## 28         13.0      M   70.0
```

```
Female <- subset(Shoes_Tble, Gender == "F", select = c(ShoeSize, Gender, Height))
Female
```

```
##      ShoeSize Gender Height
## 1          6.5      F   66.0
## 2          9.0      F   68.0
## 3          8.5      F   64.5
## 4          8.5      F   65.0
## 6          7.0      F   64.0
## 7          9.5      F   70.0
## 8          9.0      F   71.0
## 10         7.5      F   64.0
## 12         8.5      F   67.0
## 17         8.5      F   59.0
## 18         5.0      F   62.0
## 20         6.5      F   66.0
## 21         7.5      F   64.0
## 24         8.5      F   69.0
```

#c.

```
Mean_Size <- mean(ShoeSize)
Mean_Size
```

```
## [1] 9.410714
```

```
Mean_Height <- mean(Height)
Mean_Height
```

```
## [1] 68.57143
```

#d. Yes, because the bigger or taller you are the more likely your shoe size is also bigger.

#2.

```
Months <- c("March", "April", "January", "November", "January", "September", "October", "September")

Factor_Mnths <- factor(Months)
Factor_Mnths
```

```
## [1] March      April       January    November   January    September  October
## [8] September  November   August     January    November   November   February
## [15] May        August     July       December   August     August     September
## [22] November   February   April
## 11 Levels: April August December February January July March May ... September
```

```
assign("Factor_Months_Vector",Factor_Mnths)
Factor_Months_Vector
```

```
## [1] March      April       January    November   January    September  October
## [8] September  November   August     January    November   November   February
## [15] May        August     July       December   August     August     September
## [22] November   February   April
## 11 Levels: April August December February January July March May ... September
```

#3.

```
summary(Months)
```

```
##      Length      Class      Mode
##         24 character character
```

```
summary(Factor_Months_Vector)
```

```
##      April      August  December  February   January      July      March      May
##         2         4         1         2         3         1         1         1
## November  October  September
##         5         1         3
```

#4.

```
Direction <- c("East", "West", "North")
Frequency <- c(1,4,3)
DirFreq <- data.frame(Direction, Frequency)
DirFreq
```

```
##      Direction Frequency
## 1         East         1
## 2         West         4
## 3         North         3
```

```
new_order_data <- factor(Direction ,levels = c("East","West","North"))
print(new_order_data)
```

```
## [1] East West North
## Levels: East West North
```

#5.

```
Excel <- read.table("import_march.csv", header = TRUE, sep = ",", stringsAsFactors = FALSE)
Excel
```

```
## Student Strategy.1 Strategy.2 Strategy.3
## 1 Male 8 10 8
## 2 4 8 6
## 3 0 6 4
## 4 Female 14 4 15
## 5 10 2 12
## 6 6 0 9
```