

THE STRING BORDISM OF BE_8 AND $BE_8 \times BE_8$ THROUGH DIMENSION 14

MICHAEL A. HILL

ABSTRACT. We compute the low dimensional String bordism groups $\tilde{\Omega}_k^{String} BE_8$ and $\tilde{\Omega}_k^{String}(BE_8 \times BE_8)$ using a combination of Adams spectral sequences together with comparisons to the Spin bordism cases.

1. INTRODUCTION

In this paper, we will answer a question posed by Hisham Sati about the low dimensional String bordism groups of spaces of particular interest to string theorists. We specifically compute the bordism groups of $K(\mathbb{Z}, 4)$; of BE_8 , the classifying space for principal bundles for the exceptional Lie group E_8 ; and of their cartesian squares. These computations have applications to various models string theory built out of E_8 bundles, many of which will be spelled out more fully in a future paper.

This computation generalizes computations of Stong of the Spin bordism groups of $K(\mathbb{Z}, 4)$ and of Edwards of the Spin bordism of $BE_8 \times BE_8$ [19, 9]. In particular, we see that the String bordism of $K(\mathbb{Z}, 4)$ injects into the Spin bordism of $K(\mathbb{Z}, 4)$ through dimension at least 14. This means that many of the cohomological invariants used to detect bordism classes of Spin manifolds apply equally well here. In particular, the comparison of Adams spectral sequences used to compute the 10th String bordism group of BE_8 shows that the Landweber-Stong invariant detects this String bordism class [15, 8].

Our main results are summarized in the following two theorems.

Theorem 1.1. *Through dimension 15, we have the following reduced String bordism groups.*

k	4	5	6	7	8	9	10	11	12	13	14
$\tilde{\Omega}_k^{String} BE_8$	\mathbb{Z}	0	0	0	\mathbb{Z}	$\mathbb{Z}/2$	$\mathbb{Z}/2$	0	\mathbb{Z}^2	0	0

The String bordism groups of $BE_8 \times BE_8$ contain two copies of the String bordism groups of BE_8 , together with the String bordism of the space $BE_8 \wedge BE_8$. These new groups are given in the following theorem.

Theorem 1.2. *Through dimension 15, we have the following reduced String bordism groups.*

1991 *Mathematics Subject Classification.* 57R90, 55N22, 55N34.
Dedicated to the memory of Robert Stong.

k	4	5	6	7	8	9	10	11	12	13	14
$\widetilde{\Omega}_k^{String}(BE_8 \wedge BE_8)$	0	0	0	0	\mathbb{Z}	0	$\mathbb{Z}/2$	0	$\mathbb{Z}^2 \oplus \mathbb{Z}/3$	0	$(\mathbb{Z}/2)^2$

2. COMPUTATIONAL REDUCTIONS

2.1. Reduction to tmf and $K(\mathbb{Z}, 4)$. Our actual computation will be of the tmf -homology of $K(\mathbb{Z}, 4)$. We proceed in this section to explain why this computation is sufficient.

The σ -orientation of Ando-Hopkins-Strickland is the primary tool. This is an E_∞ ring map $MString \rightarrow tmf$ that refines the Witten genus [2].

Theorem 2.1. *The σ -orientation is 15-connected.*

Proof. We argue this locally, showing that it is true at 2, at 3, and with 6 inverted.

For $p = 2$, we show this is true in cohomology. Since the σ orientation is a ring map, composition with the unit map shows that

$$\sigma^*: \mathbb{Z} = H^0(tm f) \rightarrow H^0(MString) = \mathbb{Z}$$

is an isomorphism.

Computations of Stong and Bahri-Mahowald show that

$$H^*(MString) = \mathcal{A} // \mathcal{A}(2) \oplus M,$$

for a module M that is 20-connected [20, 3]. Since σ^* is an isomorphism on H^0 , we conclude that

$$\mathcal{A} // \mathcal{A}(2) = H^*(tm f) \hookrightarrow H^*(MString).$$

In particular, the smallest degree element of the cokernel is in dimension 20, giving the result for $p = 2$.

The remaining cases are simpler. For $p = 3$, computations of Hovey and Ravenel show that the σ -orientation is 15-connected [14]. When 6 is a unit, there is no torsion in the homotopy of $MString$, and in degrees less than 17, there are polynomial generators in degree 8, 12, and 16. The σ -orientation sends the generator in degree 8 to c_4 and the one in degree 12 to c_6 . This gives the final case. \square

Bott and Samelson showed that the Postnikov section $E_8 \rightarrow K(\mathbb{Z}, 3)$ is 14-connected [5]. This implies that the Postnikov section $BE_8 \rightarrow K(\mathbb{Z}, 4)$ is an isomorphism in homotopy in degrees less than 16. In particular, we see that if $X^{[k]}$ denotes the k -skeleton of X , then

$$K(\mathbb{Z}, 4)^{[15]} \simeq BE_8^{[15]}.$$

Since the spectra tmf and $MString$ are (-1) -connected, the inclusion of the 15-skeleton induces an isomorphism in homology through degree 14. Combined with the previous theorem, we conclude the following.

Corollary 2.2. *As graded rings,*

$$\widetilde{tmf}_k K(\mathbb{Z}, 4) \cong \widetilde{\Omega}_k^{String} BE_8$$

for $k < 15$, where the isomorphism is the composite of the Postnikov section with the σ -orientation.

This is a substantial simplification, since tmf -homology at all primes is computable using the Adams spectral sequence.

2.2. Forms of the Adams Spectral Sequence. Hopkins and Mahowald have shown that $H^*(tmf) = \mathcal{A}/\mathcal{A}(2)$, where $\mathcal{A}(2)$ is the subalgebra of the Steenrod algebra generated by Sq^1 , Sq^2 , and Sq^4 . Standard change-of-rings arguments then show the following.

Theorem 2.3 ([12]). *If X is a finite spectrum, then there is an Adams spectral sequence of the form*

$$E_2 = \text{Ext}_{\mathcal{A}(2)}^{s,t} \left(\tilde{H}^*(X; \mathbb{F}_2), \mathbb{F}_2 \right) \Longrightarrow \widetilde{tmf}_{t-s} X_2^\wedge.$$

At $p = 3$, the cohomology of tmf is not a cyclic \mathcal{A} -module, so similar techniques no longer work. By considering instead a slightly refined version of the Adams spectral sequence, described by Baker and Lazarev [4], we can still build an analogous spectral sequence.

There is a Hopf algebra $\tilde{\mathcal{A}}(1)$ analogous to $\mathcal{A}(2)$ used to compute tmf -homology at $p = 3$. This is most easily expressed in the dual formulation:

$$\tilde{\mathcal{A}}(1)_* = \mathbb{F}_3[\xi_1]/\xi_1^3 \otimes E(\tau_0, \tau_1, a_2),$$

where ξ_1 , τ_0 , and τ_1 arise from the elements of the dual Steenrod algebra of the same name, and where the coproduct on a_2 is given by

$$\psi(a_2) = a_2 \otimes 1 + 1 \otimes a_2 + \xi_1 \otimes \tau_1 - \xi_1^2 \otimes \tau_0.$$

Theorem 2.4 ([11]). *If X is a finite spectrum, then there is an Adams spectral sequence of the form*

$$E_2 = \text{Ext}_{\tilde{\mathcal{A}}(1)}^{s,t} \left(\tilde{H}^*(X; \mathbb{F}_3), \mathbb{F}_3 \right) \Longrightarrow \widetilde{tmf}_{t-s} X_3^\wedge.$$

At $p > 3$, the computations are simplified by the lack of p -torsion and a splitting of tmf into spectra built out of BP .

Theorem 2.5 ([13, 17]). *At primes larger than 3, we have a splitting*

$$tmf_{(p)} = \bigvee BP\langle 2 \rangle.$$

Since $H^*(BP\langle 2 \rangle) = \mathcal{A}/E(Q_0, Q_1, Q_2)$, where Q_i is the i^{th} Milnor primitive, standard change-of-rings arguments allow us to compute each summand of $tmf_* X$ arising from this splitting.

Theorem 2.6 ([16]). *If X is a finite spectrum then there is an Adams spectral sequence of the form*

$$E_2 = \text{Ext}_{E(Q_0, Q_1, Q_2)}^{s,t} \left(\tilde{H}^*(X; \mathbb{F}_p), \mathbb{F}_p \right) \Longrightarrow \widetilde{BP\langle 2 \rangle}_{t-s} X_p^\wedge$$

At the prime 2, we will also have need of the ko -homology of $K(\mathbb{Z}, 4)$. Since $H^*(ko) = \mathcal{A}/\mathcal{A}(1)$, where $\mathcal{A}(1)$ is the subalgebra of the Steenrod algebra generated by Sq^1 and Sq^2 , a change-of-rings argument similar to that for tmf allows us to compute ko -homology.

Theorem 2.7. *If X is a finite spectrum, then there is a spectral sequence of the form*

$$E_2 = \text{Ext}_{\mathcal{A}(1)}^{s,t} \left(\tilde{H}^*(X; \mathbb{F}_2), \mathbb{F}_2 \right) \Longrightarrow \widetilde{ko}_{t-s} X_2^\wedge.$$

In cohomology, the effect of the canonical map $tmf \rightarrow ko$ is the quotient

$$H^*(ko) = \mathcal{A} // \mathcal{A}(1) \rightarrow \mathcal{A} // \mathcal{A}(2) = H^*(tmf).$$

This means that the induced map on Adams spectral sequences is given on E_2 terms by the canonical map

$$\mathrm{Ext}_{\mathcal{A}(2)} \left(\tilde{H}^*(X; \mathbb{F}_2), \mathbb{F}_2 \right) \rightarrow \mathrm{Ext}_{\mathcal{A}(1)} \left(\tilde{H}^*(X; \mathbb{F}_2), \mathbb{F}_2 \right)$$

induced by the inclusion of $\mathcal{A}(1)$ into $\mathcal{A}(2)$.

Many of our Adams spectral sequence arguments will be made more clear with pictures of the corresponding E_2 pages. In all of these charts, the horizontal axis represents $t - s$, the difference between the internal grading and the Ext degree, and this corresponds to the stem of the target. This means that all of the groups in a column reassemble to give a single homotopy group. The vertical direction represents s , the Ext degree, and a d_r differential decreases $t - s$ by 1 and increases s by r . Additionally, in the pictures that follow, a dot represents a copy of \mathbb{Z}/p , and vertical lines connecting dots represents multiplication by a class v_0 from the Adams spectral sequence of the sphere. This class detects multiplication by p , reflecting a non-trivial additive extension between the linked copies of \mathbb{Z}/p . In particular, a class which is v_0 -torsion free represents a copy of \mathbb{Z}_p .

Since $MString$ and BE_8 are finite type, we know that copies of \mathbb{Z}_p in the p -completion arise from copies of $\mathbb{Z}_{(p)}$ in the localization. We will therefore blur the distinction in what follows, stating without further comment that v_0 -towers yield copies of $\mathbb{Z}_{(p)}$.

2.3. Cohomology of $K(\mathbb{Z}, 4)$. The final ingredient is the cohomology of Eilenberg-MacLane spaces, worked out by Cartan and Serre [7, 18].

Theorem 2.8. *The cohomology of $K(\mathbb{Z}, n)$ with coefficients in \mathbb{F}_p is the free graded commutative unstable algebra over the Steenrod algebra generated by a class ι_n in degree n subject to the relation that $\beta\iota_n = 0$.*

In particular, $H^*(K(\mathbb{Z}, 4); \mathbb{F}_p)$ is generated by classes ι_4 and $\mathcal{P}^I \iota_4$ for all admissible I of excess less than 4 that do not end in β . This is a very harsh restriction.

Corollary 2.9. *For $p > 5$, the only cohomology classes of dimension less than 16 are ι_4 , ι_4^2 , and ι_4^3 .*


For $p = 5$, there are additionally the classes $\mathcal{P}^1 \iota_4$ and $\beta \mathcal{P}^1 \iota_4$ in degrees 12 and 13 respectively.

For $p = 3$, there are classes ι_4 , $\mathcal{P}^1 \iota_4$, $\beta \mathcal{P}^1 \iota_4$, ι_4^2 , $\iota_4^3 = \mathcal{P}^2 \iota_4$, $\iota_4 \mathcal{P}^1 \iota_4$, and $\iota_4 \beta \mathcal{P}^1 \iota_4$.

For $p = 2$, there are classes ι_4 , $Sq^2 \iota_4$, $Sq^3 \iota_4$, $\iota_4^2 = Sq^4 \iota_4$, $Sq^4 Sq^2 \iota_4$, $\iota_4 Sq^2 \iota_4$, $Sq^5 Sq^2 \iota_4$, $\iota_4 Sq^3 \iota_4$, $(Sq^2 \iota_4)^2 = Sq^6 Sq^2 \iota_4$, ι_4^3 , $Sq^6 Sq^3 \iota_4$, $Sq^2 \iota_4 Sq^3 \iota_4$, $(Sq^3 \iota_4)^2 = Sq^7 Sq^3 \iota_4$, $\iota_4 Sq^4 Sq^2 \iota_4$, $\iota_4^2 Sq^2 \iota_4$, $\iota_4^2 Sq^3 \iota_4$, and $\iota_4 Sq^5 Sq^2 \iota_4$.

We remark that for $p > 5$, the Bockstein spectral sequence collapses, showing that the integral cohomology of $K(\mathbb{Z}, 4)$ is p -torsion free through dimension 15.

The modules for $p = 3$ and $p = 2$ are best understood via pictures. In these, the vertical direction indicates the dimension, and dots represent basis elements. In the picture for $p = 3$, curved lines represent the action of \mathcal{P}^1 and straight lines represent the action of β . The module is presented in Figure 1(a). For $p = 2$, Theorem 2.3 shows us that we need only understand the cohomology as a module over $\mathcal{A}(2)$.

FIGURE 1. $H^k(K(\mathbb{Z}, 4); \mathbb{F}_p)$ for $k < 16$ as an \mathcal{A} -module

For this, straight lines indicate Sq^1 , curved lines indicate Sq^2 and brackets indicate Sq^4 . The module (with a slightly nicer basis) is depicted in Figure 1(b).

This collection of data is sufficient to carry out almost all of the required computations.

3. THE GROUPS $\tilde{\Omega}_k^{String}(BE_8)$ FOR $k < 15$

3.1. Computation with 6 Inverted. When 6 is inverted, the spectrum tmf has no torsion. This makes running the Atiyah-Hirzebruch spectral sequence for $\widehat{tmf}_k K(\mathbb{Z}, 4)$ much simpler.

Theorem 3.1. *For $k < 15$, we have*

$$\tilde{\Omega}_k^{String} BE_8 \otimes \mathbb{Z}[\frac{1}{6}] = \begin{cases} \mathbb{Z}[\frac{1}{6}] & \text{when } k = 4, 8, \\ \mathbb{Z}[\frac{1}{6}] \times \mathbb{Z}[\frac{1}{6}] & \text{when } k = 12, \\ 0 & \text{otherwise.} \end{cases}$$

Proof. Localized at primes larger than 5, there is no torsion in the integral homology of $K(\mathbb{Z}, 4)$ in our range. For degree reasons, the Atiyah-Hirzebruch spectral sequence then collapses, giving the result.

At $p = 5$, the Atiyah-Hirzebruch spectral sequence becomes less transparent:

$$H_{12}(K(\mathbb{Z}, 4); \mathbb{Z}[\frac{1}{6}]) = \mathbb{Z}/5 \oplus \mathbb{Z}[\frac{1}{6}],$$

and this propagates to give a possible 5-torsion class in $\tilde{\Omega}_{12}^{String} K(\mathbb{Z}, 4)$. Using instead the Adams spectral sequence, we see that in fact $\tilde{\Omega}_{12}^{String} K(\mathbb{Z}, 4)[\frac{1}{6}]$ is torsion free, since the class $\beta\mathcal{P}^1$ is Q^1 on integral classes and detects v_1 . At $p = 5$, the class v_1 is the class c_4 in tmf [17], showing that the 12th String bordism group is in fact torsion free. \square

Remark. The possible 5-torsion in $\tilde{\Omega}_{12}^{String} K(\mathbb{Z}, 4)$ is exactly analogous to the possible 3-torsion Stong encountered computing $\tilde{\Omega}_8^{Spin} K(\mathbb{Z}, 4)$. Stong elegantly avoided the use of the Adams spectral sequence by appealing to characteristic classes. A

similar argument applies here, relating the mod 5 reduction of p_2 to the first Wu class.

3.2. Computation at $p = 3$. The space $K(\mathbb{Z}, 4)$ splits stably at $p = 3$, due to the presence of the automorphism given by multiplication by -1 . This acts on i_4 by -1 , and the stable splitting is reflected in the cohomology by splitting the cohomology into ± 1 eigenspaces. In particular, the two summands of Figure 1(a) arise from different stable summands, and therefore they do not interact in the Adams spectral sequence.

The relevant Ext groups for the two modules making up $H^*(K(\mathbb{Z}, 4); \mathbb{F}_3)$ are easy to compute using the long exact sequences in Ext induced by the inclusions of the skeleta [11], and the analogue of the Adams E_2 term is given in Figure 2.




FIGURE 2. An Adams E_2 term for $\widetilde{tmf}_k K(\mathbb{Z}, 4)$, $k < 15$

We see that for degree reasons, there are no differentials and no possible extensions. In particular, there are no torsion summands in this range, and we conclude that at 3, the results are essentially the same as for large primes.

Theorem 3.2. *For $k < 15$, we have that*

$$\tilde{\Omega}_k^{String} BE_8 \otimes \mathbb{Z}_{(3)} = \begin{cases} \mathbb{Z}_{(3)} & \text{when } k = 4, 8, \\ \mathbb{Z}_{(3)}^2 & \text{when } k = 12, \\ 0 & \text{otherwise.} \end{cases}$$

3.3. Computation at $p = 2$. Using a program written by Robert Bruner [6], we compute

$$\text{Ext}_{\mathcal{A}(2)}^{s,t} \left(\tilde{H}^*(K(\mathbb{Z}, 4); \mathbb{F}_2), \mathbb{F}_2 \right).$$

The result is depicted in Figure 3. In this figure, lines of slope 1 are multiplication by η , the generator of $\pi_1^s(S^0)$. Since the stable homotopy groups of spheres are the cobordism groups of framed manifolds, we can interpret this class as a framed manifold: S^1 together with the non-bounding framing.




FIGURE 3. The Adams E_2 term for $\widetilde{tmf}_k K(\mathbb{Z}, 4)$, $k < 15$

There is the possibility for a d_2 differential in degrees 10 and 11. Using Stong's computation of $\tilde{\Omega}_{11}^{Spin}K(\mathbb{Z}, 4)$, we can show that there is such a differential.

Theorem 3.3 ([19]). *We have*

$$\tilde{\Omega}_{10}^{Spin}K(\mathbb{Z}, 4) = \mathbb{Z}/2 \oplus \mathbb{Z}/2 \text{ and } \tilde{\Omega}_{11}^{Spin}K(\mathbb{Z}, 4) = 0.$$

We can show our desired differential by mapping down to ko . There is a commutative square

$$\begin{array}{ccc} MString & \longrightarrow & MSpin \\ \sigma \downarrow & & \downarrow \hat{A} \\ tmf & \longrightarrow & ko. \end{array}$$

The \hat{A} -genus is the projection onto one of the summands of $MSpin$ described by Anderson, Brown, and Peterson [1]. Through dimension 10, the splitting looks like

$$MSpin = ko \vee ko[8] \vee ko[10],$$

where $ko[n]$ denotes the $(n - 1)$ -connected cover of ko .

Since $K(\mathbb{Z}, 4)$ begins in dimension 4, we conclude that the \hat{A} -genus induces an isomorphism

$$\tilde{\Omega}_k^{Spin}(K(\mathbb{Z}, 4)) \xrightarrow{\cong} \widetilde{ko}_k K(\mathbb{Z}, 4)$$

for $k \leq 11$.

The Adams E_2 term for $\widetilde{ko}_*K(\mathbb{Z}, 4)$ is very easy to compute directly, and is presented in Figure 4 through dimension 14.




FIGURE 4. The Adams E_2 term for $ko_*K(\mathbb{Z}, 4)$

It is similarly easy to see that the map on Adams E_2 terms induced by $tmf \rightarrow ko$ is injective through dimension 14.

Stong's computation that $\tilde{\Omega}_{11}^{Spin}K(\mathbb{Z}, 4)$ is 0 shows that the class ηx_{10} cannot survive the Adams spectral sequence. For degree reasons, the only way to remove this class is for it to support a differential, and the only possibility is

$$d_2(\eta x_{10}) = \eta^2 x_8.$$

Since the Adams spectral sequence for ko -homology is a spectral sequence of modules over the Adams spectral sequence for ko_* , we conclude that we also have a differential of the form

$$d_2(x_{10}) = \eta x_8.$$

Naturality of the Adams spectral sequence implies that the same must be true for tmf .

Corollary 3.4. *There is a d_2 differential in dimensions 10 and 11 in the Adams spectral sequence for $\widetilde{tmf}_*K(\mathbb{Z}, 4)$.*

In the homotopy of ko , there is a Massey product relation linking the generator of the 4 stem with the lower generators:

$$ko_4 = \mathbb{Z} \cdot \langle 2, \eta, \eta^2 \rangle.$$

This relationship propagates, and in particular, we know that the generator of 13 stem in the Adams E_2 term for $\widetilde{ko}_*K(\mathbb{Z}, 4)$ is linked by a bracket of the form $\langle 2, \eta, - \rangle$ to ηx_{10} . Adams differentials satisfy a kind of Leibnitz rule [16], and this allows us to conclude that there is a d_2 differential on the 13-dimensional generator, hitting the generator of the \mathbb{Z} -tower beginning in filtration 4 in dimension 12 modulo indeterminacy. For this bracket, the indeterminacy is the submodule generated by v_0 times each of the generators in this degree, and thus cannot cancel out the piece of the differential hitting the aforementioned generator.

Corollary 3.5. *In dimension 12, $\widetilde{\Omega}_k^{String}K(\mathbb{Z}, 4) \otimes \mathbb{Z}_{(2)}$ is torsion free.*

Using an analysis built out of the fact that $H^{13}(K(\mathbb{Z}, 4); \mathbb{Z}) = \mathbb{Z}/20$, Francis has shown that the target of the differential hits non-trivial multiples of all of the classes in that degree [10].

For degree reasons, there are no further differentials possible and no possible extensions. We conclude the following theorem.

Theorem 3.6. *For $k < 15$, we have*

$$\widetilde{\Omega}_k^{String}BE_8 \otimes \mathbb{Z}_{(2)} = \begin{cases} \mathbb{Z}_{(2)} & \text{when } k = 4, 8, \\ \mathbb{Z}/2 & \text{when } k = 9, 10, \\ \mathbb{Z}_{(2)}^2 & \text{when } k = 12, \\ 0 & \text{otherwise.} \end{cases}$$

Our analysis of the Adams spectral sequences for $\widetilde{ko}_*K(\mathbb{Z}, 4)$ and $\widetilde{tmf}_*K(\mathbb{Z}, 4)$ shows an additional result, since all of the differentials for $\widetilde{ko}_*K(\mathbb{Z}, 4)$ lift to differentials for $\widetilde{tmf}_*K(\mathbb{Z}, 4)$.

Corollary 3.7. *The map*

$$\widetilde{\Omega}_k^{String}(BE_8) \otimes \mathbb{Z}_{(2)} \rightarrow \widetilde{\Omega}_k^{Spin}(BE_8) \otimes \mathbb{Z}_{(2)}$$

is injective for $k < 14$.

Theorems 3.1, 3.2, and 3.6 together give a restatement of Theorem 1.1. In each degree, we simply find a finitely generated abelian group which has the correct localizations with respect to each prime, and this yields the table given in the introduction. Since the String bordism groups with 2 inverted are torsion free, Corollary 3.7 implies that integrally,

$$\widetilde{\Omega}_k^{String}(BE_8) \rightarrow \widetilde{\Omega}_k^{Spin}(BE_8)$$

is injective for $k < 14$.

In particular, in dimension 10, a dimension of particular interest to String theorists, the cobordism invariant discussed by Diaconescu-Moore-Witten can also be used to detect cobordism classes of String 10-manifolds equipped with a 4-dimensional cohomology class [8].

4. THE GROUPS $\tilde{\Omega}_k^{String}(BE_8 \times BE_8)$ FOR $k < 15$

The earlier analysis goes through *mutatis mutandis*, showing that it will suffice to compute $\widetilde{tmf}_k K(\mathbb{Z} \times \mathbb{Z}, 4)$. Since we have a stable splitting

$$K(\mathbb{Z} \times \mathbb{Z}, 4) = K(\mathbb{Z}, 4) \vee K(\mathbb{Z}, 4) \vee K(\mathbb{Z}, 4) \wedge K(\mathbb{Z}, 4),$$

the homology computation will be two copies of the results of the previous section together with a factor coming from the smash square of $K(\mathbb{Z}, 4)$.

The cohomology of this is easily determined. Let I denote the augmentation ideal of the graded algebra $H^*(X; \mathbb{F}_p)$ for any simply connected space X . Since we are working over a field, the Künneth theorem says that

$$H^*(X \times X; \mathbb{F}_p) = H^*(X; \mathbb{F}_p) \otimes H^*(X; \mathbb{F}_p).$$

The reduced cohomology of $X \wedge X$ is therefore given by the ideal $I \otimes I$ sitting in this algebra, and the action of the Steenrod algebra is given by the Cartan formula and the action on $H^*(X; \mathbb{F}_p)$. In particular, since the augmentation ideal begins in degree 4, there are very few possible classes in $I \otimes I$ below degree 16.

4.1. Computation with 6 Inverted. For degree reasons, the Atiyah-Hirzebruch spectral sequence collapses, and we quickly conclude the following theorem.

Theorem 4.1. *For $k < 15$, we have*

$$\tilde{\Omega}_k^{String}(BE_8 \wedge BE_8)[\tfrac{1}{6}] = \begin{cases} \mathbb{Z}[\tfrac{1}{6}] & \text{when } k = 8, \\ \mathbb{Z}[\tfrac{1}{6}]^2 & \text{when } k = 12, \\ 0 & \text{otherwise.} \end{cases}$$

4.2. The Case of $p = 3$. Since we have a stable splitting of the form

$$K(\mathbb{Z}, 4) = A \vee B,$$



where the bottom cell of A is in dimension 4 while that of B is in 8, we have a similar splitting

$$K(\mathbb{Z}, 4) \wedge K(\mathbb{Z}, 4) = (A \wedge A) \vee (A \wedge B) \vee (B \wedge A) \vee (B \wedge B).$$

Since we are looking only through dimension 15, the cells in dimensions larger than 9 in A play no role. The 9-skeleton of A is a desuspension of B , and this means that for purposes of our computation, each summand of the right-hand side is a suspension of spectra of the form $A \wedge A$. On these, there is an action of $\mathbb{Z}/2$ given by interchanging the factors. This again allows us to split each factor into eigenspaces. Working through all of the computations allows us to easily compute $\tilde{H}^*(K(\mathbb{Z}, 4) \wedge K(\mathbb{Z}, 4))$, and this is presented in Figure 5. We remark that here the bottom cell is in dimension 8, meaning that the figure contains more information than is necessary.

The above analysis shows that the splitting drawn for the modules is reflected topologically. This means that there are no possible differentials or extensions linking the summands. A fairly straightforward computation yields the relevant Ext groups, and the result is given in Figure 6.

We see in particular that there are no possible differentials or extensions. This allows us to conclude the following theorem.


FIGURE 5. $\tilde{H}^k(K(\mathbb{Z}, 4) \wedge K(\mathbb{Z}, 4); \mathbb{F}_3)$ for $k \leq 15$ FIGURE 6. An Adams E_2 term for $\widetilde{tmf}_*(K(\mathbb{Z}, 4) \wedge K(\mathbb{Z}, 4))$ at 3

Theorem 4.2. *Through dimension 15,*

$$\tilde{\Omega}_k^{String}(BE_8 \wedge BE_8) \otimes \mathbb{Z}_{(3)} = \begin{cases} \mathbb{Z}_{(3)} & \text{when } k = 8, \\ \mathbb{Z}_{(3)}^2 \oplus \mathbb{Z}/3 & \text{when } k = 12, \\ 0 & \text{otherwise.} \end{cases}$$

4.3. The Case of $p = 2$. For $p = 2$, the analysis for $K(\mathbb{Z}, 4) \wedge K(\mathbb{Z}, 4)$ is much simpler than that for $K(\mathbb{Z}, 4)$. Computing the action of the Steenrod algebra in the cohomology is very easy using the method described above. However, given the number of cells, it is also incredibly tedious. Bruner's Ext program package contains a program for computing the tensor product of two $\mathcal{A}(2)$ modules, allowing us to automate this computation. Combining this computation with the Ext program allows us to compute the relevant groups. These are depicted in Figure 7.

For degree reasons, there are no possible differentials or extensions, and we conclude the following theorem.

FIGURE 7. The Adams E_2 term for $\widetilde{tmf}_k(K(\mathbb{Z}, 4) \wedge K(\mathbb{Z}, 4))$, $k < 15$

Theorem 4.3. For $k < 15$,

$$\tilde{\Omega}_k^{String}(BE_8 \wedge BE_8) \otimes \mathbb{Z}_{(2)} = \begin{cases} \mathbb{Z} & \text{when } k = 8, \\ \mathbb{Z}/2 & \text{when } k = 10, \\ \mathbb{Z}^2 & \text{when } k = 12, \\ (\mathbb{Z}/2)^2 & \text{when } k = 14, \\ 0 & \text{otherwise.} \end{cases}$$

Just as before, Theorems 4.1, 4.2, and 4.3 together yield Theorem 1.2. Comparing the localizations yields the table from that theorem. We do remark that while the map

$$\tilde{\Omega}_k^{String} BE_8 \rightarrow \tilde{\Omega}_k^{Spin} BE_8$$

is injective for $k < 15$, the map

$$\tilde{\Omega}_k^{String}(BE_8 \times BE_8) \rightarrow \tilde{\Omega}_k^{Spin}(BE_8 \times BE_8)$$

is not.

REFERENCES

1. D. W. Anderson, E. H. Brown, Jr., and F. P. Peterson, *The structure of the Spin cobordism ring*, Ann. of Math. (2) **86** (1967), 271–298. MR MR0219077 (36 #2160)
2. Matthew Ando, Michael J. Hopkins, and Neil P. Strickland, *The sigma orientation is an H_∞ map*, Amer. J. Math. **126** (2004), no. 2, 247–334. MR MR2045503 (2005d:55009)
3. A. P. Bahri and M. E. Mahowald, *A direct summand in $H^*(MO\langle 8 \rangle, \mathbb{Z}_2)$* , Proc. Amer. Math. Soc. **78** (1980), no. 2, 295–298. MR MR550517 (81b:57029)
4. Andrew Baker and Andrej Lazarev, *On the Adams spectral sequence for R -modules*, Algebr. Geom. Topol. **1** (2001), 173–199 (electronic). MR MR1823498 (2002a:55011)
5. Raoul Bott and Hans Samelson, *Applications of the theory of Morse to symmetric spaces*, Amer. J. Math. **80** (1958), 964–1029. MR MR0105694 (21 #4430)
6. Robert R. Bruner, *Ext in the nineties*, Algebraic topology (Oaxtepec, 1991), Contemp. Math., vol. 146, Amer. Math. Soc., Providence, RI, 1993, pp. 71–90. MR MR1224908 (94a:55011)
7. Henri Cartan, *Sur les groupes d'Eilenberg-Mac Lane. II*, Proc. Nat. Acad. Sci. U. S. A. **40** (1954), 704–707. MR MR0065161 (16,390b)
8. Duiliu-Emanuel Diaconescu, Gregory Moore, and Edward Witten, *E_8 gauge theory, and a derivation of K -theory from M -theory*, Adv. Theor. Math. Phys. **6** (2002), no. 6, 1031–1134 (2003). MR MR1982693 (2004i:58062)
9. Steven R. Edwards, *On the spin bordism of $B(E_8 \times E_8)$* , Illinois J. Math. **35** (1991), no. 4, 683–689. MR MR1115992 (92d:57026)
10. John N. K. Francis, *Spin bordism of $BSpin$ and $K(\mathbb{Z}, 4)$: Integrality and index theory*, Undergraduate Thesis.
11. Michael A. Hill, *The 3-local tmf-homology of $B\Sigma_3$* , Proc. Amer. Math. Soc. **135** (2007), no. 12, 4075–4086 (electronic). MR MR2341960
12. Michael J. Hopkins and Mark Mahowald, *From elliptic curves to homotopy theory*, Available on the Hopf Archive, 1998.

13. Michael J. Hopkins and Haynes R. Miller, *Elliptic curves and stable homotopy I*, In preparation.
14. Mark A. Hovey and Douglas C. Ravenel, *The 7-connected cobordism ring at $p = 3$* , Trans. Amer. Math. Soc. **347** (1995), no. 9, 3473–3502. MR MR1297530 (95m:55008)
15. Peter S. Landweber and Robert E. Stong, *A bilinear form for Spin manifolds*, Trans. Amer. Math. Soc. **300** (1987), no. 2, 625–640. MR MR876469 (88h:57020)
16. Douglas C. Ravenel, *Complex cobordism and stable homotopy groups of spheres*, Pure and Applied Mathematics, vol. 121, Academic Press Inc., Orlando, FL, 1986.
17. Charles Rezk, *Supplimentary notes for math 512*, <http://www.math.uiuc.edu/rezk/papers.html>.
18. Jean-Pierre Serre, *Sur les groupes d'Eilenberg-MacLane*, C. R. Acad. Sci. Paris **234** (1952), 1243–1245. MR MR0046047 (13,675c)
19. R. E. Stong, *Appendix: calculation of $\Omega_{11}^{\text{Spin}}(K(Z, 4))$* , Workshop on unified string theories (Santa Barbara, Calif., 1985), World Sci. Publishing, Singapore, 1986, pp. 430–437. MR MR849112
20. Robert E. Stong, *Determination of $H^*(\text{BO}(k, \dots, \infty), \mathbb{Z}_2)$ and $H^*(\text{BU}(k, \dots, \infty), \mathbb{Z}_2)$* , Trans. Amer. Math. Soc. **107** (1963), 526–544. MR MR0151963 (27 #1944)

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF VIRGINIA, CHARLOTTESVILLE, VA 22904
E-mail address: mikehill@virginia.edu