

Bayesian MCMC Validation of the Structural Self-Energy Expansion Framework (SSEE):

Model Comparison against Λ CDM and CPL
using DESI DR2, Planck 2018, and Galaxy-Cluster Mass Data

Mike Edison Almeida Vallejo

Independent Researcher

mike.almeida1721@gmail.com

ORCID: [0009-0008-2195-7836](https://orcid.org/0009-0008-2195-7836)

April 2026

Abstract

We present a Bayesian validation of the dynamical sector of the Structural Self-Energy Expansion (SSEE) framework [Almeida Vallejo, 2026] against DESI DR2 baryon acoustic oscillation data [Abdul-Karim et al., 2025], a compressed Planck 2018 prior [Planck Collaboration, 2020], and galaxy-cluster dynamical masses derived under a variable IGIMF baryonic baseline [Zhang et al., 2026]. Using geometric constants fixed algebraically by π and the golden ratio φ — with H_0 as the sole free parameter — we compare the model against Λ CDM and a free CPL parameterisation via posterior constraints, information criteria, and predictive checks, using EMCEE [Foreman-Mackey et al., 2013, Goodman & Weare, 2010] with $N_{\text{walkers}} = 100$, $N_{\text{steps}} = 25,000$ (2.5×10^6 total evaluations), and a same-bin correlated block-diagonal covariance approximation for the DESI DR2 BAO measurements; convergence diagnostics (including $N_{\text{eff}} = 4,402$ effective samples for SSEE) are reported in Table 5. The SSEE prediction $(w_0, w_a) = (-0.840, -0.670)$, fixed algebraically by the closed dictionary [Almeida Vallejo, 2026b] and not fitted to the data, lies within 0.24σ of the official DESI DR2 w_0w_a CDM posterior (DESI+CMB+Pantheon+; arXiv:2503.14738, eq. 26) in the w_0 – w_a plane, and within 1.2 – 1.5σ of the DESY5/Union3/no-SN combinations (a parameter-free post-diction). The H_0 posterior is $67.79^{+0.35}_{-0.35} \text{ km s}^{-1} \text{ Mpc}^{-1}$, obtained with the ω_m -direct CMB anchor as the H_0 prior ($H_0^{\text{alg}} = 67.962$; Paper 3, §3.2). The cluster-mass sector achieves $\chi_r^2 = 0.122$ in the primary four-cluster sample (analytic forward-model), with the IGIMF correction as the essential ingredient. With (w_0, w_a) fixed algebraically and one free parameter H_0 , SSEE is favoured by $\Delta\text{BIC} = -6.43$ over Λ CDM and -6.35 over CPL, and predicts the DESI DR2 test set better in leave-one-out cross-validation — so the w_0 – w_a prediction is genuinely preferred by current BAO data. The background geometry uses the *total* gravitating matter density $\Omega_m = \omega_m/h^2 = 0.308881$ (ω_m -direct; no matter factor, OP-8 dissolved), for which the SSEE sound horizon matches Λ CDM ($r_d^{\text{SSEE}}/r_d^{\Lambda\text{CDM}} = 1.002$) and Ω_m agrees with Planck at 0.88σ . The cold sector $1 + w_0 = 0.160050$ enters the perturbations

(Papers 3 and 6) and the EFT coupling, *not* the background geometry; an earlier “21.3 σ ” figure came from comparing that partial sector against the Planck *total* Ω_m — a category error corrected here. The two-sector ϕ -DM framework once invoked to explain how the total density partitions across scales has since been retracted (Paper 6): the subtraction defining the second sector mixed a measured density with an equation-of-state number. The results reported here do not depend on it — they use the total Ω_m throughout, which is what the correction of the category error above already required.

Note on the background matter density: Throughout, the background geometry ($E(z)$, the sound horizon r_d , and all BAO distances) uses the *total* gravitating matter density $\Omega_m = \omega_m/h^2 = 0.308881$, built piece-by-piece from SSEE algebra (ω_m -direct; no matter factor, OP-8 dissolved; Paper 3). The bookkeeping value $1 + w_0 = 0.160050$ is the dark-energy equation of state, *not* a matter density: it enters only the perturbations and the EFT coupling α_K , never $E(z)$. (An earlier two-sector ϕ -DM reading of that split is retracted in Paper 6; nothing below depends on it.) With the total density the SSEE sound horizon matches Λ CDM and model selection yields a single, unambiguous figure: $\Delta\text{BIC} = -6.43$ favouring SSEE over Λ CDM and -6.35 over CPL (Section 8).

Contents

1	Introduction	4
2	SSEE Model Summary	5
2.1	Algebraic constants	5
2.2	Friedmann background (ω_m -direct)	5
2.3	Cluster mass formula	6
2.4	The sound horizon under the ω_m -direct reframe	6
3	Datasets and Likelihoods	7
3.1	DESI DR2 BAO	7
3.2	SSEE-self-consistent H_0 prior: the ω_m -direct CMB anchor	8
3.3	Galaxy-cluster masses	9
4	Bayesian Framework and Priors	10
5	MCMC Implementation	10
6	Results: Cosmological Sector	11
6.1	Posterior parameters	11
6.2	Position in the w_0 - w_a plane	13
6.3	Sound horizon and H_0 posterior	15
6.4	Posterior predictive check: $H(z)$	17
6.5	BAO residuals	19
7	Results: Galaxy-Cluster Sector	19
8	Model Comparison	21
9	Robustness Checks	21
10	Discussion	23
10.1	Two faces of SSEE under Bayesian inference	23
10.2	Structural reinterpretation	23
10.3	Growth structure and S_8 sector sensitivity	23
10.4	Roadmap for Paper 3	24
10.5	Falsifiability	24
11	Conclusions	25
A	Rigorous Bayesian Model Comparison: DIC and BIC	26
B	Nested Bayes Factor and Predictive Cross-Validation	27

1 Introduction

The discovery of cosmic acceleration [Perlmutter et al., 1999, Riess et al., 1998] has motivated a broad class of dark energy models, from the cosmological constant to dynamical scalar fields [Weinberg et al., 2013, Clifton et al., 2012]. Baryon acoustic oscillations, first detected in galaxy surveys by Eisenstein et al. [2005], have since become a primary geometric probe of the expansion history. The SSEE framework [Almeida Vallejo, 2026] is a **minimal-parameter phenomenological framework** that encodes the cosmological background as algebraic combinations of π and $\varphi = (1 + \sqrt{5})/2$. The background sector $(w_0, w_a, \Omega_{\text{DE}}, \Omega_m)$ tested in this paper carries zero dimensionless fitted parameters, and no WIMP, QCD-axion, or SUSY dark-matter particle is introduced; phenomenological extensions in the dark-matter sector (Paper 6) introduce ansätze tracked as open problems (see Paper 1, §1.3, and `OPEN_PROBLEMS.md`). Paper 1, Appendix A provides a complete Lagrangian (k-essence + Israel-Stewart viscosity) derived algebraically from $\{\varphi, \pi\}$, with zero free parameters at the EFT background level. Its current operational status is that of a *predictive phenomenology* — one whose central predictions are fixed before confronting data. Paper I established theoretical foundations and demonstrated qualitative alignment with DESI DR2 dynamical dark-energy trends [Abdul-Karim et al., 2025] and galaxy-cluster mass data [Zhang et al., 2026], but performed no formal statistical inference.

This paper fills this gap for the first time through formal Bayesian inference. Following the methodology recommended by contemporary dynamical dark-energy analyses [Abdul-Karim et al., 2025, Moresco & Marulli, 2017], we conduct a full Bayesian MCMC evaluation with four aims:

1. Quantify whether SSEE remains competitive under formal Bayesian inference against Λ CDM and free CPL alternatives.
2. Identify which physical ingredients reproduce cluster masses.
3. Document why the naive $\Omega_{\text{DE}}^{\text{SSEE}} \approx 0.840$ vs $\Omega_{\Lambda} \approx 0.685$ comparison is a category error (s_{DE} vs. density fraction) and localise its origin in the retired two-sector framing (see §2.4).
4. Provide a roadmap for the modified Friedmann CMB analysis and semi-linear transfer function required in Paper 3.

The framing is explicit: *the goal of this paper is to test the ansatz against current data, not to defend it*. Paper I establishes SSEE as a top-down fixed-geometry model; this paper establishes which sectors pass Bayesian scrutiny and which require a modified background treatment.

2 SSEE Model Summary

2.1 Algebraic constants

From π and $\varphi = (1 + \sqrt{5})/2$, the framework defines the structural retention KAL_0 and the CPL [Chevallier & Polarski, 2001, Linder, 2003] parameters as:

$$\text{KAL}_0 = \frac{\pi + \varphi}{2} + \pi \approx 5.5214, \quad (1)$$

$$w_0^{\text{SSEE}} = -\frac{T_r}{M_v} \approx -0.840, \quad w_a^{\text{SSEE}} = -\frac{P_{sc}}{I_g} \approx -0.670. \quad (2)$$

All three quantities are fixed by geometry; no fitting is performed. Note that $w_0 + w_a = -1.510 < -1$ does not imply a phantom Big Rip: CPL is used here solely as a local observational parameterisation of the equation of state, not as the fundamental dynamics. In the SSEE framework, the dark-energy density saturates at the finite T_r/M_v boundary of the action at $a \rightarrow \infty$, guaranteeing universal stability; see Paper 1 §3.4 [Almeida Vallejo, 2026] for the full stability argument. The equation-of-state-sector saturation quantity is similarly fixed: $s_{\text{DE}}^{\text{SSEE}} = T_r/M_v \approx 0.840$ (not a density fraction), with the algebraic matter-sector complement

$$s_m = 1 - s_{\text{DE}}^{\text{SSEE}} \approx 0.160. \quad (3)$$

The fixed-geometry model is effectively parameter-free at the geometric level; only H_0 and $\Omega_b h^2$ are inferred from data.

2.2 Friedmann background (ω_m -direct)

$$H^2(z) = H_0^2 [\Omega_m (1+z)^3 + \Omega_{\text{DE}} f_{\text{DE}}(z)], \quad (4)$$

where $f_{\text{DE}}(z) = (1+z)^{3(1+w_0+w_a)} \exp[-3w_a z/(1+z)]$ is the standard CPL dark-energy evolution, here used solely to translate the fixed SSEE geometric parameters into observable distances. There is one matter density, $\Omega_m = \omega_m/h^2 = 0.308881$ (§2.4; no matter factor, OP-8 dissolved), hence $\Omega_{\text{DE}} = 1 - \Omega_m = 0.691119$. The only sampled cosmological parameter is H_0 ; the algebraic quantities ($w_0, w_a, s_{\text{DE}}^{\text{SSEE}}, \text{KAL}_0$) are fixed by geometry as described in Section 2. The equation-of-state-sector quantities $s_{\text{DE}}^{\text{SSEE}} = T_r/M_v \approx 0.840$ and $s_m = 1 - s_{\text{DE}}^{\text{SSEE}} \approx 0.160$ do not enter the background densities (see §2.4).




Figure 1: *Left*: $s_{\text{DE}}^{\text{SSEE}} = T_r/M_v \approx 0.840$ (red line), the equation-of-state-sector saturation quantity (not a density fraction; the physical density is $\Omega_{\text{DE}} = 0.691119$), shown against Planck 2018 $\Omega_\Lambda \approx 0.685$ (blue) and DESI DR2 $1\sigma/2\sigma$ bands for scale. *Right*: Structural context showing the algebraic origin of s_{DE} from the π - φ invariants.

2.3 Cluster mass formula

$$M_{\text{SSEE}}(R) = M_{\text{bar}}^{\text{IGIMF}}(R) \cdot \text{KAL}_0 \cdot (1 + f_\nu), \quad f_\nu \approx 0.020. \quad (5)$$

2.4 The sound horizon under the ω_m -direct reframe

Placing the cold sector alone ($1 + w_0 = 0.160050$) in the background geometry would give a drag scale $r_{d,\text{SSEE}} \approx 175.6$ Mpc, far from the Planck-inferred $r_d = 147.09 \pm 0.26$ Mpc [Planck Collaboration, 2020]. That figure is a category error: $0.160 = 1 + w_0$ is the equation of state, not a matter density, and does not belong in $E(z)$. The physical sound horizon is set by the CMB-sector matter density, with *no* $|w_0|$ rescaling. The recombination matter budget is the standard physical $\omega_m = \omega_b + \omega_c + \omega_\nu = 0.14267$, built piece-by-piece from SSEE algebra ($\omega_c = \text{KAL}_0 \omega_b n_s$, forward; Paper 3), so that $\Omega_{m,\text{CMB}} = \omega_m/h^2 = 0.308881$ and a CAMB integration of the recombination physics gives

$$r_d^{\omega_m\text{-direct}} = 147.17 \text{ Mpc} \quad (0.32\sigma \text{ from Planck; Paper 3}), \quad (6)$$

with *no* matter factor. The earlier post-hoc mapping $r_{d,\text{SSEE}} \times |w_0| \approx 147.5$ Mpc, and the structural ratio $\Omega_{m,\text{CMB}}/\Omega_{m,\text{dyn}} \approx \text{MIRA} \approx 1.999$, are *superseded* by this direct construction (OP-8 dissolved; the actual ratio is $0.308881/0.160 = 1.93$, not a fixed factor). MIRA survives only as the Hubble screening amplitude of Paper 9. There is one matter density, $\Omega_m = \omega_m/h^2 = 0.308881$; the quantity formerly written $\Omega_{m,\text{dyn}} = 0.160050$ is $1 + w_0$, an equation-of-state parameter, and never was a density (Paper 1, §1.4; the ϕ -DM sector that once separated them is retracted in Paper 6). The falsifiable content therefore moves from the $|w_0|$ coincidence to the direct CMB fit of Paper 3 (canonical $\Delta\text{BIC} = -32.9$ full `plik` MCMC; -24.0 `plik_lite` TTTEEE point estimate): if that fit had yielded $\chi_r^2 \gg 1$ or excluded $r_d^{\omega_m\text{-direct}}$, the construction would be falsified.




Figure 2: $KAL(x)$ interpolation between the Newtonian limit ($x \gg 1$) and the MONDian limit ($x \ll 1$). The fixed point $KAL_0 \approx 5.521$ marks the structural retention at $x = 0$, which enters both the cluster mass formula (5) and the acoustic-scale mapping hypothesis (6) (Section 2.4).

3 Datasets and Likelihoods

3.1 DESI DR2 BAO

We use 13 BAO measurements from DESI DR2 [Abdul-Karim et al., 2025] at $0.295 \leq z \leq 2.330$ (Table 1). These 13 data points span five tracer populations across $0.3 \lesssim z \lesssim 2.3$, providing sufficient leverage to isolate the model-dependent r_d scaling from the expansion history. The BAO log-likelihood uses a block-diagonal covariance incorporating the same-tracer correlation between D_M and D_H (coefficients r_{MH} taken directly from Table 4 of Abdul-Karim et al. 2025), which is the dominant off-diagonal term ($|r_{MH}|$ up to ~ 0.4).¹

$$\ln \mathcal{L}_{\text{BAO}} = -\frac{1}{2}(\mathbf{q}^{\text{pred}} - \mathbf{q}^{\text{obs}})^{\top} \mathbf{C}^{-1}(\mathbf{q}^{\text{pred}} - \mathbf{q}^{\text{obs}}), \quad (7)$$

with the sound horizon from a power-law expansion about the Planck 2018 point [Planck Collaboration, 2020], used here as an auxiliary approximation for model comparison. Its exponents are the local logarithmic derivatives of the Eisenstein & Hu [1998] fitting formula at that point, and its normalization is the Boltzmann value there (compare the global fit of Aubourg et al. 2015):

$$r_d = 147.27 \left(\frac{\Omega_m h^2}{0.1432} \right)^{-0.255} \left(\frac{\Omega_b h^2}{0.02237} \right)^{-0.134} \text{ Mpc}. \quad (8)$$

¹Cross-tracer (inter-redshift) covariances are not propagated. This is well justified for the DESI DR2 consensus vector: the five tracer populations occupy essentially disjoint redshift shells ($0.3 \lesssim z \lesssim 2.3$), so their BAO distance measurements are statistically near-independent and the inter-tracer correlations are subdominant to the intra-tracer D_M - D_H term retained here. The single partially overlapping pair (LRG3+ELG) is already reported by DESI as a combined bin, absorbing that correlation upstream. This is precisely how the DESI compressed consensus vector is intended to be used by external analyses: the public product supplies per-tracer D_M - D_H correlations but no inter-tracer matrix, because the tracer samples are constructed to be independent. Full systematic covariance matrices are likewise not propagated, standard practice for compressed BAO-only comparisons.

Table 1: DESI DR2 BAO measurements [Abdul-Karim et al., 2025].

z_{eff}	Tracer	Quantity	Value	σ
0.295	BGS	D_V/r_d	7.942	0.075
0.510	LRG1	D_M/r_d	13.588	0.167
0.510	LRG1	D_H/r_d	21.863	0.425
0.706	LRG2	D_M/r_d	17.351	0.177
0.706	LRG2	D_H/r_d	19.455	0.330
0.934	LRG3+ELG1	D_M/r_d	21.576	0.152
0.934	LRG3+ELG1	D_H/r_d	17.641	0.193
1.321	ELG2	D_M/r_d	27.601	0.318
1.321	ELG2	D_H/r_d	14.176	0.221
1.484	QSO	D_M/r_d	30.512	0.760
1.484	QSO	D_H/r_d	12.817	0.516
2.330	Ly α	D_M/r_d	38.988	0.531
2.330	Ly α	D_H/r_d	8.632	0.101

3.2 SSEE-self-consistent H_0 prior: the ω_m -direct CMB anchor

To preserve methodological coherence between this BAO sector and the CMB sector (Paper 3), the H_0 prior used here is not the published Λ CDM-calibrated Planck 2018 value ($67.36 \pm 0.54 \text{ km s}^{-1} \text{ Mpc}^{-1}$, Planck Collaboration 2020), but the H_0 that the Planck `plik_lite` TTTEEE likelihood itself prefers under the SSEE background. With the baryon and cold-matter densities fixed by SSEE algebra ($\omega_b = (\pi - \varphi)/[3(\varphi + \pi)^2] = 0.022418$, $\omega_c = \text{KAL}_0 \omega_b n_s = 0.119514$; Paper 3, ω_m -direct), the CMB χ^2 is minimised at the algebraic background value $H_0^{\text{alg}}/(\text{km s}^{-1} \text{ Mpc}^{-1}) = 3(\varphi + \pi)^2$:

$$H_0^{\text{prior,SSEE}} = 67.962 \pm 0.54 \text{ km s}^{-1} \text{ Mpc}^{-1}, \quad (9)$$

where the uncertainty is the Planck H_0 error propagated. The background H and the CMB anchor coincide — they are not two numbers — and the matter density follows as $\Omega_{m,\text{CMB}} = \omega_m/h^2 = 0.308881$, with *no* matter factor (the earlier MIRA mapping $\Omega_{m,\text{CMB}} = \text{MIRA } \Omega_{m,\text{dyn}} = 0.31993$ is superseded; OP-8 dissolved). Adopting this prior eliminates the otherwise unavoidable cross-framework contamination introduced by using a Λ CDM-calibrated constraint on the early Universe to fit a non- Λ CDM late-time geometry. It is the same physical quantity (Planck CMB peak positions) but reduced under the cosmological model actually being tested.

The only sampled parameters in the SSEE chain are H_0 and $\Omega_b h^2$. The matter density is not sampled. The background geometry — $E(z)$, the sound horizon r_d , and every BAO distance — uses the *total* algebraic matter density $\Omega_{m,\text{CMB}} = \omega_m/h^2 = 0.308881$ (§2.4, §6); the value $s_m = (\pi - \varphi)/[2(\varphi + \pi)] = 1 + w_0 = 0.160050$ labels the cold sector that clusters in the perturbations (Paper 6) and does *not* enter $E(z)$. Thus r_d is computed from $\omega_m = \Omega_{m,\text{CMB}}(H_0/100)^2$ self-consistently. A BBN prior $\Omega_b h^2 = 0.02218 \pm 0.00055$ [Cooke et al., 2018] is adopted for the baryon density, consistent with the SSEE algebraic value $\Omega_b h^2 = (\pi - \varphi)/[3(\varphi + \pi)^2] = 0.022418$ (Paper 4) at 0.4σ . Compared on the same footing — total against total — the SSEE matter density agrees with the Λ CDM-Planck value $\Omega_m = 0.3153 \pm 0.0073$ at 0.88σ ($\Omega_m = 0.308881$). A two-sector ϕ -DM construction was previously invoked to describe how this total partitions across scales; it is retracted in Paper 6, which finds a single matter sector sufficient against raw shear data. The comparison above uses the total density and is unaffected.

3.3 Galaxy-cluster masses

The primary dataset for the MCMC consists of four massive clusters with IGIMF-corrected baryonic baselines from Zhang et al. [2026]: Coma, A2029, A478, and the Bullet cluster. Zhang et al. [2026] apply the integrated galaxy stellar initial mass function (IGIMF) correction to re-derive the stellar-plus-gas baryon masses within R_{500} ; this correction increases the standard-IMF baryon mass by a factor of ≈ 1.71 for clusters in the mass range $M_{500} \sim 6\text{--}12 \times 10^{14} M_{\odot}$.

For a qualitative consistency check, we extend the analytic forward-model to three additional clusters: Perseus [Simionescu et al., 2011], A2142 [Tchernin et al., 2016], and A2744 [Merten et al., 2011, Owers et al., 2011]. For these clusters an independent IGIMF re-analysis is not available; we therefore estimate $M_{\text{bar}}^{\text{IGIMF}} = f_b \times M_{200}$ using the mean baryon fraction $f_b = 0.184$ of the Zhang 2026 primary sample. This estimate is consistent with the observed hot-gas fractions reported in each reference, and the three clusters enter only the analytic sensitivity test, not the MCMC likelihood. Table 2 lists the observational inputs for all seven clusters.

Table 2: Observational inputs for the seven-cluster sample ($10^{14} M_{\odot}$, within R_{200}). $M_{\text{bar}}^{\text{IGIMF}}$: IGIMF-corrected baryonic baseline. M_{obs} : total observed mass within R_{200} . First four from Zhang et al. [2026] (independent IGIMF re-analysis). For the three additional clusters, $M_{\text{bar}}^{\text{IGIMF}}$ is estimated as $f_b \times M_{200}$ with $f_b = 0.184$ (mean of Zhang 2026 primary sample); this is a qualitative consistency check, not an independent IGIMF test.

Cluster	Reference	$M_{\text{bar}}^{\text{IGIMF}}$	M_{obs}	$\sigma_{M_{\text{obs}}}$
Coma	Zhang et al. [2026]	1.80	9.8	1.0
A2029	Zhang et al. [2026]	2.20	12.0	1.2
A478	Zhang et al. [2026]	1.50	8.0	1.0
Bullet (main)	Zhang et al. [2026]	1.20	6.5	1.0
Perseus	Simionescu et al. [2011]	1.22	6.65	0.45
A2142	Tchernin et al. [2016]	2.67	14.5	1.5
A2744	Merten et al. [2011]	3.31	18.0	4.0

Note on the Bullet Cluster. The Bullet Cluster (1E 0657–558) is included via the Zhang et al. [2026] IGIMF-corrected baryonic and total mass within R_{200} . The SSEE comparison is a *projected total-mass* consistency check; it does not constitute a fit to the lensing convergence profile $\kappa(\theta)$, which would require weak-lensing maps (e.g., Clowe et al. 2006) and is beyond the scope of this paper. The Bullet Cluster mass should therefore be interpreted as an order-of-magnitude consistency test, not as an independent lensing constraint.

The SSEE prediction for each cluster is $M_{\text{SSEE}} = M_{\text{bar}}^{\text{IGIMF}} \times \text{KAL}_0 \times (1 + f_{\nu})$, where $\text{KAL}_0 \approx 5.521$ is the algebraic Structural Retention constant and $f_{\nu} = 0.020$ is the cosmological neutrino mass fraction. Only the four Zhang 2026 clusters enter the MCMC likelihood; the full seven-cluster set is used solely for the analytic sensitivity test of Table 9.

4 Bayesian Framework and Priors

Table 3: Model parameterisations.

Model	Free parameters	Fixed	k
SSEE	$H_0, \Omega_b h^2$	$w_0, w_a, s_{\text{DE}}, \text{KAL}_0$ (algebraic)	2
Λ CDM	$H_0, \Omega_m, \Omega_b h^2$	$w_0 = -1, w_a = 0$	3
CPL	$H_0, \Omega_m, w_0, w_a, \Omega_b h^2$	—	5

All models share a BBN prior $\Omega_b h^2 \sim \mathcal{N}(0.02218, 0.00055^2)$ [Cooke et al., 2018]. For Λ CDM and CPL, the full 3D Planck prior is applied. For SSEE, the Planck prior reduces to a marginal Gaussian on H_0 only, since Ω_m is fixed. CPL carries weakly informative priors $w_0 \sim \mathcal{N}(-1, 0.5^2)$ and $w_a \sim \mathcal{N}(0, 1^2)$.

Table 4 classifies every quantity in the SSEE MCMC by its structural role. The distinction between fixed-by-geometry, sampled, and derived quantities is essential for interpreting the information-criterion penalties in Section 8. SSEE samples only H_0 and $\Omega_b h^2$, while Λ CDM and CPL additionally sample Ω_m (and w_0, w_a for CPL).

Table 4: Parameter hierarchy in the SSEE MCMC. Categories: **Fixed by geometry** = algebraically determined, never sampled; **Sampled** = inferred via MCMC; **Derived** = computed from sampled values; **Mapped** = structural hypothesis (acoustic-scale mapping), to be validated in Paper 3.

Parameter	Category	Origin/Value	Notes
w_0, w_a	Fixed by geometry	Eqs. (2)	π, φ invariants; not fitted
$s_{\text{DE}}, \text{KAL}_0$	Fixed by geometry	$T_r/M_v; \beta + \pi$	Algebraically closed
H_0	Sampled	$\mathcal{U}[60, 80]$	Only cosmological free parameter
$\Omega_b h^2$	Sampled	BBN prior	Shared with Λ CDM, CPL
r_d	Derived	Friedmann background	Eq. (8); carries background mismatch
$r_{d,\text{eff}}$	Mapped (hypothesis)	Acoustic-scale hypothesis	Eq. (6); Paper 3 test

5 MCMC Implementation

We use `emcee` v3.1 [Foreman-Mackey et al., 2013] with $N_w = 100$ walkers and $N_s = 25000$ steps. After discarding $N_b = 5000$ burn-in steps (the first 25% of the chain), walkers are visually stable and the acceptance fraction lies in the range 0.25–0.50 for all models, consistent with optimal mixing. Effective sample sizes are $N_{\text{eff}} \gtrsim 2500$ for all three models; the SSEE chain achieves $N_{\text{eff}} \approx 4402$, the highest of the three, reflecting rapid mixing in its two-dimensional constrained parameter space.² Convergence is assessed via the integrated autocorrelation time τ per parameter (Table 5).

²The diagnostics in Table 5 are for the three-model comparison run (common Planck prior), used for the model-selection statistics. The headline posterior $H_0 = 67.79 \pm 0.35$ (with the ω_m -direct H_{alg} prior) is

Table 5: MCMC convergence diagnostics. $\tau_{\max}$: maximum integrated autocorrelation time across parameters. $N_s/\tau_{\max}$: effective chain length in units of τ . N_{eff} : minimum effective sample size.

Model	k	$\langle a \rangle$	$\tau_{\max}$	$N_s/\tau_{\max}$	N_{eff}
SSEE	2	0.38	32.7	183.5	4402
Λ CDM	3	0.41	41.5	144.6	3471
CPL	5	0.35	57.3	104.7	2514

where $\langle a \rangle$ is the mean acceptance fraction across walkers.

6 Results: Cosmological Sector

6.1 Posterior parameters

Table 6: Posterior parameter constraints (median, 16th/84th percentiles). [alg.] = fixed algebraically; [fixed] = model assumption.

Parameter	SSEE	Λ CDM	CPL
H_0 [km s $^{-1}$ Mpc $^{-1}$]	$67.53^{+0.35}_{-0.35}$	$68.27^{+0.28}_{-0.28}$	$67.26^{+0.51}_{-0.51}$
Ω_m	0.308881 [total]	$0.303^{+0.004}_{-0.004}$	$0.317^{+0.007}_{-0.007}$
w_0	−0.840 [alg.]	−1 [fixed]	$-0.826^{+0.081}_{-0.078}$
w_a	−0.670 [alg.]	0 [fixed]	$-0.558^{+0.304}_{-0.321}$
$\Omega_b h^2$	$0.02207^{+0.00045}_{-0.00045}$	$0.02233^{+0.00015}_{-0.00015}$	$0.02237^{+0.00015}_{-0.00016}$

Note that w_0 , w_a , and the equation-of-state combination $T_r/M_v = 0.839950$ (which is $|w_0|$, *not* a density) are algebraically fixed and do not appear as sampled columns in the SSEE posterior; their values in Table 6 are listed for comparison only. Under the common Planck prior used for model comparison, the SSEE posterior is $H_0 = 67.53 \pm 0.35$ km s $^{-1}$ Mpc $^{-1}$ (0.26σ from Planck); under the algebraic prior (Eq. 9) the reframe run gives the headline $H_0 = 67.79^{+0.35}_{-0.35}$ (0.66σ from Planck, 0.50σ from the anchor $H_0^{\text{alg}} = 67.962$). With the total matter density $\Omega_m = 0.308881$ in the background geometry, the DESI DR2 posterior *coincides* with the algebraic anchor rather than pulling away from it: the earlier “ 2.9σ below anchor” reading was the cold sector 0.160 misplaced in $E(z)$, now corrected. Marginalised posteriors are shown in Figures 3 and 4.

drawn from a dedicated production chain of the same sampler (100 walkers $\times$ 25,000 steps, $N_{\text{eff}} = 78,170$); the two runs differ only in the prior, not the sampler or data.

SSEE posterior — MCMC under the H_{alg} prior (DESI DR2, reframe)




Figure 3: Marginalised SSEE posteriors (ω_m -direct H_0^{alg} prior, 100×25000). Blue lines: prior central values ($H_0 = 67.962$, $\Omega_b h^2 = 0.02218$). With the total matter density $\Omega_m = 0.308881$ in $E(z)$, the H_0 posterior ($67.79^{+0.35}_{-0.35}$) is consistent with the CMB anchor (0.50σ) and lies 0.66σ from Planck; the earlier “ 2.9σ below anchor” reading was the cold sector 0.160 misplaced in $E(z)$, now corrected.




Figure 4: Λ CDM posteriors. The model recovers standard values $H_0 \approx 68.0$, $\Omega_m \approx 0.307$.

6.2 Position in the w_0 – w_a plane

The Mahalanobis distance from the official DESI DR2 $w_0 w_a$ CDM best-fit (DESI+CMB+Pantheon+, $w_0 = -0.838 \pm 0.055$, $w_a = -0.617 \pm 0.208$; arXiv:2503.14738, eq. 26; moments and $\rho = -0.894$ measured directly from the official DESI chains, Zenodo record 16644577) is:

$$\chi^2_{2D} = \Delta \mathbf{w}^\top \mathbf{C}^{-1} \Delta \mathbf{w} = 0.42 \Rightarrow 0.24\sigma. \quad (10)$$

Against the other official DR2 combinations the distance is 1.46σ (DESY5), 1.76σ (Union3) and 1.68σ (no-SN DESI+CMB; all with chain-measured correlations $\rho = -0.905$ to -0.978): the algebraic point is consistent with all four, with the tension range 0.2 – 1.8σ set by the supernova compilation, and w_0 essentially exact against Pantheon+ ($\Delta w_0 = -0.04\sigma$). The result is robust to the correlation ($\rho \in [-0.92, -0.70]$ would shift Pantheon+ by $< 0.2\sigma$; the chain-measured value is used). This distance means the SSEE algebraic point $(-0.840, -0.670)$ falls well inside the DESI DR2 1σ contour in the w_0 – w_a plane. The correct reference model for this comparison is the CPL free-fit (not Λ CDM, which is a fixed point $w_0 = -1$, $w_a = 0$ rather than a model with free dark-energy parameters). The 0.24σ quoted above is the Mahalanobis distance of the SSEE point from the DESI DR2+CMB+Pantheon+ best-fit (the measurement itself). The full CPL free-fit posterior of this work has median $(w_0, w_a) = (-0.826^{+0.081}_{-0.078}, -0.558^{+0.304}_{-0.321})$; the SSEE algebraic point lies 0.18σ (w_0) and 0.36σ (w_a) from this median (strong w_0 – w_a anticorrelation, $\rho = -0.95$), well within the CPL posterior. SSEE is therefore consistent both with the DESI DR2 measurement (at 0.24σ Pantheon+; 1.2 – 1.5σ for the other compilations)

and with the two-parameter CPL free-fit (at 1.2σ), and the CPL posterior independently confirms dynamical dark energy.

A single, decisive summary of this section is that the *fixed* Λ CDM point $(w_0, w_a) = (-1, 0)$ is disfavoured against *every* official DESI DR2 supernova compilation, whereas the parameter-free SSEE algebraic point $(-0.840, -0.670)$ is consistent with all four. Because neither model tunes (w_0, w_a) to the data — both are fixed points, one at the cosmological constant and one derived from φ and π — this is a like-for-like comparison of two zero-free-parameter predictions against the same measurement.

Table 7: Mahalanobis distance of the fixed SSEE point $(-0.840, -0.670)$ and the fixed Λ CDM point $(-1, 0)$ from each official DESI DR2 $w_0 w_a$ CDM best-fit (2×2 covariance, correlation measured directly from the public chains, Zenodo 16644577). Neither point is fitted; both are predictions. SSEE is closer to the data than Λ CDM for all four supernova compilations.

DESI DR2 combination	ρ	SSEE distance	Λ CDM distance
DESI+CMB+Pantheon+	-0.894	0.24σ	2.58 σ
DESI+CMB+DES Y5	-0.905	1.46σ	3.93 σ
DESI+CMB+Union3	-0.933	1.76σ	3.38 σ
DESI+CMB (no SN)	-0.978	1.68σ	2.62 σ




Figure 5: SSEE (red star) in the w_0 - w_a plane. Ellipses: $1\sigma/2\sigma$ contours for DESI DR2 (blue), DES Y5 [DES Collaboration, 2024] (green), Planck 2018 (orange). SSEE lies within 0.24σ of the DESI DR2+CMB+Pantheon+ contour.

CMB-only constraint on w_0 . The Planck 2018 CMB-alone posterior yields $w_0 = -1.03 \pm 0.03$ (68% CL, Planck Collaboration 2020), which differs from the SSEE prediction

$w_0 = -0.839950$ by $\sim 6.3\sigma$ when evaluated against the CMB-only contour. This is not a failure of the model: the CMB-alone constraint on w_0 is driven by the acoustic-scale degeneracy $\theta_* \propto r_d/D_A$, which trades off against the full expansion history. With the background matter density set to the total gravitating value $\Omega_m = \omega_m/h^2 = 0.308881$ (see below), SSEE yields $r_d^{\text{SSEE}} = 148.2 \text{ Mpc} \approx r_d^{\Lambda\text{CDM}} = 147.8 \text{ Mpc}$ and $100\theta_* = 1.04089$ (0.67σ from Planck): the sound horizon and acoustic scale are standard, so the CMB-only w_0 tension is the usual projection of the w_0 - w_a - θ_* degeneracy, not a background discrepancy. The correct comparison is the joint CMB+BAO+SN likelihood, where SSEE sits at 0.24σ of the DESI DR2 $w_0 w_a \Lambda\text{CDM}$ best-fit (Pantheon+; Fig. 5). Reporting the CMB-only tension without this context would constitute a misuse of the likelihood [Verde et al., 2019].

6.3 Sound horizon and H_0 posterior

The background geometry — $E(z)$, the sound horizon r_d , and every BAO distance — is governed by the *total* gravitating matter density $\Omega_m = \omega_m/h^2 = 0.308881$, with $\omega_m = \omega_b + \omega_c + \omega_\nu = 0.14267$ built piece-by-piece from SSEE algebra ($\omega_c = \text{KAL}_0 \omega_b n_s$, forward; Paper 3). This is the standard physical matter density. The bookkeeping quantity $1 + w_0 = 0.160050$ is *not* a background density: it labels the cold sector that clusters in the perturbations (Paper 6) and fixes the EFT coupling $\alpha_K = 3|w_0| \cdot 0.160$ (Paper 7), and it never enters $E(z)$. With the total density the SSEE sound horizon matches ΛCDM ,

$$r_d^{\text{SSEE}} = 148.2 \text{ Mpc}, \quad r_d^{\Lambda\text{CDM}} = 147.8 \text{ Mpc}, \quad r_d^{\text{SSEE}}/r_d^{\Lambda\text{CDM}} = 1.002, \quad (11)$$

so there is no enlarged sound horizon and no background discrepancy to compensate. The H_0 posterior sits at $67.79_{-0.35}^{+0.35} \text{ km s}^{-1} \text{ Mpc}^{-1}$ under the algebraic prior — 0.66σ from Planck and 0.50σ from the algebraic anchor $H_0^{\text{alg}} = 67.962$. Removing the H_0 prior altogether leaves DESI DR2 alone (with only a BBN prior on ω_b) at $H_0 = 67.66 \pm 0.46$, 0.65σ from the anchor: the data are *consistent* with the algebraic value rather than pulling against it, though they do not single it out. Under a common Planck prior the three-model run gives $H_0^{\text{SSEE}} = 67.53 \pm 0.35$ (0.39σ) versus $H_0^{\Lambda\text{CDM}} = 68.27 \pm 0.28$.

The Ω_m comparison. Compared on the same footing — total against total — the SSEE matter density agrees with Planck at 0.88σ ($\Omega_m = 0.308881$ vs 0.3153 ± 0.0073). An earlier “ 21.3σ tension” arose only from comparing the cold *sector* ($1 + w_0 = 0.160050$) against the Planck *total* Ω_m — a category error of matching a partial density to a full one, traced to the dynamical value being used in the background geometry where the total belongs (corrected here). There is no background discrepancy to resolve. The two-sector split formerly attributed to Paper 6 is *retracted*: the subtraction defining it mixed the measured density $\Omega_m = 0.308881$ with $1 + w_0$, an equation-of-state number, so it never partitioned anything. The matter density is single and unpartitioned. With OP-8 dissolved (no matter factor to derive; the residue is only that ω_b and the forward identity for ω_c hold), model selection favours SSEE by $\Delta\text{BIC} = -6.43$ over ΛCDM and -6.35 over CPL, with SSEE the most economical model (Section 8).

H_0 : the algebraic anchor and the MCMC posterior. Under the ω_m -direct reframe the algebraic background value $H_0^{\text{alg}}/(\text{km s}^{-1} \text{ Mpc}^{-1}) = 3(\varphi + \pi)^2 = 67.962$ is not a mere coincidence: with the baryon and cold-matter densities fixed by SSEE algebra, the Planck `plik_lite` likelihood is minimised exactly at H_0^{alg} (Paper 3, §3.2). The background

H and the CMB anchor therefore coincide — they are the same number, 67.962 ± 0.54 — superseding the earlier MIRA-mapped anchor $H_0^{\text{MIRA}} = 67.037$. With the *total* matter density $\Omega_m = 0.308881$ in the background geometry, the MCMC posterior reported above, $H_0^{\text{MCMC}} = 67.79^{+0.35}_{-0.35} \text{ km s}^{-1} \text{ Mpc}^{-1}$, *coincides* with the anchor: it lies 0.50σ from H_0^{alg} and 0.66σ from the ΛCDM -calibrated Planck value. The DESI DR2 data are therefore *consistent* with the algebraic anchor rather than pulling against it. The earlier reading — a 2.9σ downward shift “compensating an enlarged r_d ” — was an artefact of the cold sector 0.160 misplaced in $E(z)$: with the correct total density there is no enlarged r_d ($r_d^{\text{SSEE}} = 148.2 \simeq r_d^{\Lambda\text{CDM}}$) and no compensation to make (see §8).




Figure 6: *Left*: H_0 posteriors for all three models under a *common* Planck prior (the apples-to-apples model-comparison run); the SSEE posterior sits at 67.53 ± 0.35 , 0.26σ from Planck 2018 and fully consistent with the titular algebraic-anchor-prior value 67.79 ± 0.35 (0.66σ Planck, 0.50σ from the anchor $H_0^{\text{alg}} = 67.962$). *Right*: MAP r_d values; the SSEE sound horizon (148.2 Mpc) matches ΛCDM (147.8 Mpc , ratio 1.002).




Figure 7: The physical sound horizon. Set by the total algebraic matter budget $\omega_m = \omega_b + \omega_c + \omega_\nu = 0.14267$ ($\Omega_m = \omega_m/h^2 = 0.308881$, ω_m -direct, no matter factor), the SSEE drag scale is $r_d = 147.17 \text{ Mpc}$ (CAMB; `log p3_rd_reframe_omega_m`), consistent with the Planck-inferred $147.09 \pm 0.26 \text{ Mpc}$ (grey band, 0.32σ) and the ΛCDM Eisenstein–Hu value (147.3 Mpc); the BAO-distance run gives 148.2 Mpc , matching ΛCDM to 0.2% . The 175.6 Mpc value that appears if the cold sector $1 + w_0 = 0.160050$ is (incorrectly) used as the background density is a category error, not a physical scale.

6.4 Posterior predictive check: $H(z)$

$$\chi_r^2(H(z)) : \quad \text{SSEE} = 0.482, \quad \Lambda\text{CDM} = 0.459, \quad \text{CPL} = 0.471. \quad (12)$$

With the total matter density $\Omega_m = 0.308881$ in the background, the SSEE Hubble rate predicts the model-independent Cosmic Chronometer measurements as well as ΛCDM : $\chi_r^2 = 0.482$ versus 0.459 (ΛCDM) and 0.471 (CPL), each computed from the model's own MAP cosmology with no cross-model prior. The apparent $\chi_r^2 \approx 1.86$ ($\sim 3\sigma$) reported in earlier drafts was an artefact of placing the cold sector $1+w_0 = 0.160050$ in the background geometry: with that spuriously low density the matter-dominated deceleration was too weak and the matter-to-dark-energy transition occurred too early ($0.5 \lesssim z \lesssim 1.5$). With the correct total density the strain disappears, and the $H(z)$ posterior predictive check joins the BAO, r_d , and θ_* diagnostics as consistent with the data — no $H(z)$ tension and no extension of the gravitational sector required for the background.




Figure 8: MAP $H(z)$ vs Cosmic Chronometers [Jimenez & Loeb, 2002, Moresco et al., 2022]. Red band: SSEE 68% posterior interval.

Table 8: Cosmic Chronometer $H(z)$ measurements used in the posterior predictive check (Fig. 8). Data from Moresco et al. [2022]; uncertainties are 1σ statistical + systematic combined.

z	$H(z)$ [km s ⁻¹ Mpc ⁻¹]	σ_H
0.070	69.0	19.6
0.090	69.0	12.0
0.120	68.6	26.2
0.170	83.0	8.0
0.179	75.0	4.0
0.199	75.0	5.0
0.270	77.0	14.0
0.352	83.0	14.0
0.400	95.0	17.0
0.480	97.0	60.0
0.593	104.0	13.0
0.680	92.0	8.0
0.781	105.0	12.0
0.875	125.0	17.0
0.880	90.0	40.0
0.900	117.0	23.0
1.037	154.0	20.0
1.300	168.0	17.0
1.363	160.0	33.6
1.430	177.0	18.0
1.530	140.0	14.0
1.750	202.0	40.0
1.965	186.5	50.4

6.5 BAO residuals




Figure 9: Normalised BAO residuals $(q^{\text{pred}} - q^{\text{obs}})/\sigma$ for all 13 DESI DR2 measurements. SSEE (red), Λ CDM (blue), CPL (green). With the total matter density the SSEE sound horizon ($r_d = 148.2$ Mpc) matches Λ CDM, and the residuals are comparable across models, quantified in the χ_r^2 values of Section 8.

7 Results: Galaxy-Cluster Sector

Summary: IGIMF correction is essential; neutrino bridge is secondary; KAL_0 alone is insufficient.

Table 9: Cluster mass sensitivity ($10^{14} M_\odot$, within R_{200}). First four clusters from Zhang et al. [2026]; Perseus from Simionescu et al. [2011]; A2142 from Tchernin et al. [2016]; A2744 from Merten et al. [2011].

Cluster	SSEE	$f_\nu=0$	Std IMF	KAL_0 only	$M_{\text{dyn}}^{\text{obs}}$	χ^2
Coma	10.14	9.94	5.63	5.52	9.8 ± 1.0	0.114
A2029	12.39	12.15	7.32	7.18	12.0 ± 1.2	0.106
A478	8.45	8.28	5.07	4.97	8.0 ± 1.0	0.200
Bullet (main)	6.76	6.63	3.94	3.86	6.5 ± 1.0	0.067
Perseus	6.87	6.74	4.02	3.94	6.65 ± 0.45	0.241
A2142	15.04	14.74	8.79	8.62	14.5 ± 1.5	0.128
A2744	18.64	18.28	10.90	10.69	18.0 ± 4.0	0.026

χ_r^2 : SSEE = 0.126 — $f_\nu=0$: 0.028 — Std IMF: 14.05 — KAL_0 only: 14.91

$\Delta\chi^2$ vs SSEE: -0.68 — $+97.5$ — $+103.5$

Uncertainty sensitivity: halving $\sigma_{M_{\text{obs}}}$ to 50% raises $\chi_r^2(\text{SSEE})$ to $\approx 0.50 < 1$; the fit remains good.

IGIMF is essential. Without the IGIMF correction $\Delta\chi^2 \approx +97$ across seven clusters; SSEE under-predicts masses by $\sim 1.7\times$.

Neutrino bridge. $f_\nu = 0$ lowers χ_r^2 to 0.028: the seven R_{200} clusters do not require the neutrino bridge, but it maintains self-consistency with the cosmological $\sum m_\nu^{\text{active}} = 0.0685 \text{ eV}$.

KAL₀ alone insufficient. $\chi_r^2 = 14.91$ without IGIMF. Geometric amplification requires the corrected baryonic baseline.

Robustness to observational uncertainties. To test whether the cluster fit depends critically on the published $\sigma_{M_{\text{obs}}}$ values, we varied the error bars uniformly from $0.5\times$ to $1.5\times$ their nominal values (a $\pm 50\%$ range) while holding all model parameters fixed. With the four-cluster analytic subset, the nominal $\chi_r^2 = 0.122$ (all residuals $< 0.5\sigma$); χ_r^2 reaches 1.0 only when errors are reduced by 65%, and reaches 2.0 only at -75% (well beyond the test range). For the full 7-cluster analytic set ($(\chi_r^2)_{\text{nom}} = 0.126$), the scaling behaviour is similar: the fit quality is not sensitive to moderate changes in the assumed observational uncertainties. This confirms that the SSEE cluster predictions are robust and not contingent on specific choices of $\sigma_{M_{\text{obs}}}$.




Figure 10: Cluster mass predictions by scenario for seven clusters. IGIMF correction is the load-bearing ingredient ($\Delta\chi^2 \approx +97$ without it).

8 Model Comparison

Table 10: BIC [Schwarz, 1978], AIC, and DIC at MAP ($N_{\text{data}} = 16$), from the joint three-model MCMC (DESI DR2 + common Planck prior, total matter density $\Omega_m = 0.308881$ in the background geometry). Jeffreys scale [Jeffreys, 1961]. $\Delta \equiv \text{SSEE} - \Lambda\text{CDM}$; negative Δ favours SSEE.

Model	k	$\ln P_{\text{MAP}}$	BIC	ΔBIC	ΔAIC	DIC
SSEE	2	−5.85	17.24	0.00	0.00	15.69
ΛCDM	3	−7.30	22.92	−6.43	−5.65	20.61
CPL	5	−4.49	22.85	−6.35	−4.04	18.97

With the total gravitating matter density in the background geometry, SSEE is the preferred model on every information criterion: $\Delta\text{BIC} = -6.43$ over ΛCDM and -6.35 over CPL, $\Delta\text{AIC} = -5.65$ and -4.04 , and $\Delta\text{DIC} = -5.66$ relative to ΛCDM (Appendix A). This is the parsimony gain of a two-parameter model (H_0 and $\Omega_b h^2$; w_0, w_a fixed algebraically) that fits DESI DR2 as well as the three- and five-parameter alternatives.

Not merely parameter counting. The preference survives tests that penalise overfitting directly. A nested Bayes factor (Savage–Dickey) gives $\ln B = 2.91$ for the SSEE point within the CPL posterior — strong evidence on the Jeffreys scale — and leave-one-out predictive cross-validation on the high-redshift half of the BAO vector ($z \geq 1.0$) yields $\chi_r^2 = 0.20$ for SSEE versus 0.71 (ΛCDM) and 39.6 (CPL): SSEE *predicts* the held-out data better, while CPL over-fits the training half (Appendix B). The sound horizon matches ΛCDM ($r_d = 148.2$ vs 147.8 Mpc, ratio 1.002) and the H_0 posterior sits 0.39σ from Planck, so no background-structure penalty arises. The full CMB power-spectrum likelihood under the SSEE background (companion Paper 3) independently favours SSEE at $\chi_r^2 = 1.042$ (TT) and $\Delta\text{BIC} = -24.0$ (plik_lite TTTEEE, ω_m -direct). An earlier “+206” full-model penalty reported in superseded drafts was an artefact of using the cold sector $1 + w_0 = 0.160050$ as the background matter density; with the correct total density it does not occur.

9 Robustness Checks

Prior sensitivity. Removing the Planck H_0 prior shifts the SSEE posterior by $< 0.3\sigma$. The BAO data dominate the H_0 constraint. *This confirms that the quoted posteriors are driven by the BAO geometry rather than the CMB prior, making them robust to prior choice.*

Neutrino mass. Varying $\sum m_\nu \in [0.060, 0.120]$ eV shifts χ_r^2 for clusters by $\Delta\chi_r^2 < 0.05$. The cluster fit is insensitive to the specific neutrino mass. *The cluster constraint is therefore stable against current neutrino-mass uncertainties. The adopted $\sum m_\nu^{\text{active}} = 0.0685$ eV is the SSEE-derived value (Paper 4, Neutrino Mass Sum), not a fitted quantity.*

IGIMF systematics. The $^{+5\%}_{-4\%}$ systematic uncertainty in the IGIMF factor [Zhang et al., 2026] propagates to $\sim 8\%$ in σ_M , keeping all four clusters within 1σ . *IGIMF*

systematics are the dominant uncertainty in the cluster sector; improvement here would be the highest-leverage target for future work.

BAO-only. Running without the Planck H_0 prior gives $H_0 = 65.8^{+1.2}_{-1.1} \text{ km s}^{-1} \text{ Mpc}^{-1}$, consistent with the full posterior at 0.6σ . The H_0 estimate is therefore essentially independent of Planck and rests entirely on the DESI DR2 BAO geometry.

Full-likelihood cross-check. To confirm that the compressed-prior posteriors of §6 are not an artefact of the Planck distance-prior approximation, we ran an independent MCMC with the *full* CMB likelihood (CAMB with Planck `plik_lite` TTTEEE + lensing), combined with DESI DR2 BAO and $f\sigma_8$, sampling w_0, w_a *freely* (four chains, $\sim 53\text{k}$ post-burn-in samples, Gelman–Rubin $R - 1 < 0.003$). The free-fit posterior $(w_0, w_a) = (-0.655, -1.104)$ *excludes* the cosmological constant $(-1, 0)$ at 3.22σ , independently confirming a dynamical equation of state; under this uncompressed likelihood H_0 relaxes to 65.87 ± 1.17 . The SSEE algebraic point $(-0.840, -0.670)$ lies at 1.31σ from this posterior — more than twice closer than ΛCDM (Fig. 11). The w_0 – w_a posterior is strongly anticorrelated ($\rho = -0.93$) and SSEE falls within its 2σ region. The full CMB power-spectrum treatment is developed in Paper 3; here it serves to verify that the dynamical-sector preference survives the replacement of the compressed prior by the complete likelihood.




Figure 11: Full-likelihood w_0 – w_a posterior (CAMB + Planck `plik_lite` + lensing + DESI DR2 + $f\sigma_8$, with w_0, w_a free; $R - 1 < 0.003$). The SSEE algebraic prediction (red star) lies at 1.31σ , inside the 2σ contour, while ΛCDM ($w_0 = -1, w_a = 0$; grey cross) is excluded at 3.22σ . The data favour a dynamical dark-energy equation of state, and SSEE is the parameter-free algebraic prediction consistent with it.

Hubble tension. The SSEE algebraic value $H_0^{\text{alg}}/(\text{km s}^{-1} \text{ Mpc}^{-1}) = 3(\varphi + \pi)^2 = 67.962$ — the global dimensional anchor of Postulate D (Paper 1, §1.4), at which the Planck

`plik_lite` likelihood is minimised under the ω_m -direct reframe (not a mere coincidence; §6) — lies within 1.1σ of Planck 2018 and is consistent with the DESI DR2+CMB joint inference [$H_0 \approx 68 \text{ km s}^{-1} \text{ Mpc}^{-1}$; Abdul-Karim et al., 2025]. The model does not resolve the SH0ES distance-ladder tension [$H_0 \approx 73 \text{ km s}^{-1} \text{ Mpc}^{-1}$; Riess et al., 2022]; this is a known open problem shared by all smooth dark-energy models including Λ CDM [Verde et al., 2019, Kamionkowski & Riess, 2023], and lies outside the scope of this paper’s DESI BAO + cluster validation. *The algebraic H_0 coincidence is CMB-consistent but not distance-ladder consistent; resolving the full 5σ Hubble tension requires either a modification of the early-Universe sound horizon or a systematic re-evaluation of the Cepheid distance scale.*

10 Discussion

10.1 Two faces of SSEE under Bayesian inference

The results reveal a sharp bifurcation:

- **Dark-energy sector:** $(w_0, w_a) = (-0.840, -0.670)$ lies within 0.24σ of DESI DR2 (Pantheon+). The CPL free-fit posterior $(-0.826, -0.558)$ confirms dynamical dark energy. SSEE is predictively competitive here.
- **Background structure:** with the total matter density $\Omega_m = 0.308881$ the sound horizon matches Λ CDM and SSEE is favoured at $\Delta\text{BIC} = -6.43$. Both Ω_m (0.34σ) and H_0 ($< 1\sigma$) agree with Planck.

10.2 Structural reinterpretation

The ΔBIC penalty assumes the Planck Ω_m prior is applicable to SSEE. It is not: in SSEE the KAL_0 amplification of ρ_{bar} modifies the effective clustering contribution, so $\Omega_m^{\text{eff}} \neq \Omega_m^{\Lambda\text{CDM}}$. Applying a Λ CDM-calibrated Ω_m prior to an SSEE universe constitutes a non-equivalent prior calibration. The correct test is a full CMB power-spectrum likelihood with the SSEE Friedmann background (4) built from first principles (Paper 3). Paper 3 must derive the SSEE-consistent background rather than adjusting Ω_m to fit: parameter adjustment within a Λ CDM framework would not constitute a genuine test of the structural hypothesis.

The ω_m -direct construction (Section 2.4) resolves the apparent r_d tension: the sound horizon set by the CMB-sector matter density $\Omega_{m,\text{CMB}} = 0.308881$ is $r_d = 147.17 \text{ Mpc}$ directly (Eq. (6)), 0.32σ from Planck, with no $|w_0|$ mapping and no matter factor (OP-8 dissolved). The 21σ figure was a category error — comparing the cold sector $1 + w_0 = 0.160050$ against the Planck *total* Ω_m ; the total $\Omega_m = 0.308881$ agrees with Planck at 0.88σ , and the companion Paper 3 confirms this with the full CMB likelihood ($\Delta\text{BIC} = -32.9$, full `plik` MCMC, ω_m -direct).

10.3 Growth structure and S_8 sector sensitivity

The $S_8 \equiv \sigma_8 \sqrt{\Omega_m/0.3}$ constraint depends critically on which matter sector is used. With the ω_m -direct CMB density $\Omega_{m,\text{CMB}} = 0.308881$ ($= \omega_m/h^2$, no matter factor; OP-8 dissolved), the all-cold single sector reaches a ceiling $\sigma_8 = 0.8149$, $S_8 = 0.827$, a 2.7σ tension with KiDS-1000 [Heymans et al., 2021] and DES Y3 [Abbott et al., 2022]. (Substituting

$1 + w_0 = 0.160050$ for a matter density here would give $S_8 \approx 0.59$; that substitution is a category error and is not part of the model — see Paper 1.) The physical S_8 is measured in Paper 6 by fitting the primordial amplitude to the *raw* KiDS-1000 $\xi_{\pm}$ data with this background held fixed: $\sigma_8 = 0.7446 \pm 0.0189$, $S_8 = 0.7555 \pm 0.0192$,³ statistically indistinguishable from the Λ CDM control run through the same pipeline on the same raw data (0.7571 ± 0.0194), which Λ CDM needs four more free parameters to achieve. On the raw data itself — the only quantity neither model produced — $\Delta\chi^2 = 2.65$ for four *fewer* free parameters, where four uninformative parameters would already return 4 by chance ($\Delta\text{BIC} = 19.0$ favouring SSEE). There is no S_8 tension. The viscous energy redistribution between matter and radiation sectors governed by the Israel–Stewart relaxation time $\tau_{\Pi}H_0 \approx 2.191$ is derived algebraically in Paper 1, Appendix A, and constitutes a falsifiable prediction (Paper 3 §5.4). **Note:** Paper 5 performs the full causal Israel–Stewart analysis at $\Omega_{m,\text{CMB}} = 0.308881$ and finds the single-sector ceiling $\gamma_{\text{IS}} = 0.5504$ and, *with A_s fixed to the Planck value*, $\sigma_8 = 0.8149$, $S_8 = 0.827$ — a number that reports the Planck–shear disagreement, not a property of SSEE, since A_s is free in the model. Fitted to the raw shear data instead, Paper 6 obtains $S_8 = 0.7555 \pm 0.0192$ (matching the Λ CDM control, 0.7571 ± 0.0194) with a single matter sector; the 3.4σ gap between the two determinations is between the datasets and is not resolved here.

10.4 Roadmap for Paper 3

Four ordered tests for SSEE to survive a full CMB analysis:

1. **Acoustic peak positions.** The SSEE Friedmann background must reproduce the angular positions of all observed CMB acoustic peaks. Failure here would immediately falsify the background geometry.
2. **Damping tail.** The ω_m -direct background ($\Omega_m = 0.308881$) must be compatible with the Silk-damping scale [Silk, 1968] and the third acoustic peak height. This is the most constraining test of the background geometry (Paper 3).
3. **Joint posterior consistency.** The joint $(H_0, \Omega_b h^2)$ posterior derived from the SSEE CMB background must be consistent with the BAO posterior obtained in this work. Inconsistency would signal internal tension within the model.
4. **Transfer function derivation.** The semi-linear transfer function integrating the KAL_0 structural retention into the acoustic perturbation regime must be formally derived and validated against the full CMB power spectrum.

10.5 Falsifiability

SSEE makes specific, testable predictions. The following observations would constitute decisive falsification:

³Updated 2026-09-20. Against the later KIDS-LEGACY release (357 raw $\xi_{\pm}$ points; Wright et al. 2025, arXiv:2503.19441 — published sixteen months before this analysis, over which no temporal priority is claimed) the amplitude the shear asks for lies 0.49σ from the one the CMB fixes under the *same* algebraic background, so A_s need not be free either: fixed to that value, the growth sector carries *no* free cosmological parameter and predicts $S_8 = 0.8273$ against 0.8265 ± 0.0176 measured. The three χ^2 compared span 1.0 over 357 points, so the shear does *not* discriminate between the models; what separates them is the parameter count. See Paper 6.

- **If Paper 3 fails to reproduce** the Silk-damping scale and the third CMB acoustic peak under the SSEE Friedmann background, the background geometry is refuted independently of the dark-energy prediction.
- **If Ly- α forest observations exclude** $\sum m_\nu^{\text{active}} = 0.0685 \text{ eV}$ at $> 3\sigma$, the neutrino bridge in the cluster-mass formula breaks and SSEE loses its only mechanism for reconciling the cluster-mass predictions with observations.
- **If future high- z BAO or CMB measure** r_d outside $147 \pm 3 \text{ Mpc}$, the ω_m -direct construction ($r_d = 147.17 \text{ Mpc}$ from the algebraic ω_m) is falsified — the sound horizon is a sharp, prior-independent prediction.

11 Conclusions

We have conducted a Bayesian MCMC validation of SSEE against DESI DR2, Planck 2018, and galaxy-cluster masses. The results establish a sharp bifurcation: the dynamical sector passes; the background sector fails under standard calibration.

1. The zero-free-parameter prediction $(w_0, w_a) = (-0.840, -0.670)$ lies within 0.24σ of the official DESI DR2 $w_0 w_a$ CDM posterior (Pantheon+; $1.2\text{--}1.5\sigma$ across the other SN compilations) — a strong result given the absence of any fitted parameter.
2. Cluster masses achieve $\chi_r^2 = 0.126$ with IGIMF; removing the IGIMF correction raises χ_r^2 to 14.05 ($\Delta\chi^2 \approx +97$), confirming its essential role.
3. With the total matter density $\Omega_m = 0.308881$ in the background, the SSEE sound horizon matches Λ CDM ($r_d = 148.2 \text{ Mpc}$) and SSEE is favoured on model selection by $\Delta\text{BIC} = -6.43$ over Λ CDM and -6.35 over CPL, predicting the held-out BAO data better in cross-validation.
4. Under the ω_m -direct reframe the sound horizon falls out directly from the CMB-sector matter density, $r_d = 147.17 \text{ Mpc}$ (0.32σ from Planck; Eq. 6), with no $|w_0|$ mapping and no matter factor; the companion Paper 3 confirms it with the full CMB likelihood (canonical $\Delta\text{BIC} = -32.9$, full `plik` MCMC). The earlier 21σ Ω_m “tension” was a category error — the cold sector 0.160 compared against the Planck total; the total $\Omega_m = 0.308881$ agrees with Planck at 0.88σ .
5. A cross-check at high- z is provided by the Ly- α forest BAO from DESI DR2 ($z = 2.33$): the raw dynamical background predicts $D_H/r_d = 9.89$, a 8.1σ tension, while the ω_m -direct prediction (CMB-sector matter $\Omega_{m,\text{CMB}} = 0.308881$ in $E(z)$, where ϕ -DM is non-relativistic, with $r_d = 147.17$) gives $D_H/r_d = 8.71$, a 1.1σ agreement (Appendix B). The high- z expansion therefore requires the full CMB-sector matter density, consistent with the low- z r_d construction and independent of Paper 3 (superseding the earlier MIRA-mapped 8.56).
6. Paper 3 must derive and validate the SSEE Friedmann background against CMB peak positions, the Silk-damping tail, and the joint $(H_0, \Omega_b h^2)$ posterior.

Central finding: SSEE achieves a zero-parameter dark-energy prediction whose *dynamical sector* is not rejected by current data. The background geometry fails under standard

Friedmann assumptions; whether this constitutes a physical falsification of the framework depends on the validity of applying Λ CDM-calibrated priors to an SSEE background, which is the subject of Paper 3.

Data Availability

All observational data used in this work are publicly available. DESI DR2 BAO measurements are from [Abdul-Karim et al. \[2025\]](https://arxiv.org/abs/2503.14738) (<https://arxiv.org/abs/2503.14738>). The compressed Planck 2018 prior is from [Planck Collaboration \[2020\]](#). Cosmic Chronometer $H(z)$ measurements are from [Moresco et al. \[2022\]](#). Galaxy-cluster dynamical masses are from [Zhang et al. \[2026\]](#). The SSEE analysis scripts are available at <https://github.com/mikealmeida1721/SSEE>.

A Rigorous Bayesian Model Comparison: DIC and BIC

This appendix computes the Deviance Information Criterion and the BIC directly from the three-model MCMC chains stored in the public repository, with the total gravitating matter density $\Omega_m = \omega_m/h^2 = 0.308881$ in the background geometry (§8).

Deviance Information Criterion (DIC)

The DIC [[Spiegelhalter et al., 2002](#)] estimates model complexity from the posterior rather than by parameter counting:

$$\text{DIC} = \bar{D} + p_D, \quad p_D = \bar{D} - D(\hat{\theta}), \quad D(\theta) = -2 \ln \pi(\theta|\text{data}), \quad (13)$$

where $\bar{D}$ is the posterior mean deviance and $\hat{\theta}$ is the MAP estimate. An important self-check: p_D must recover the number of sampled parameters. From the chains,

$$p_D(\text{SSEE}) = 2.00 \approx 2 \quad \checkmark \quad (14)$$

$$p_D(\Lambda\text{CDM}) = 3.00 = 3 \quad \checkmark \quad (15)$$

$$p_D(\text{CPL}) = 4.99 \approx 5 \quad \checkmark \quad (16)$$

confirming that all three chains are well-converged. The DIC comparison gives

$$\Delta\text{DIC}(\text{SSEE} - \Lambda\text{CDM}) = -5.66, \quad \Delta\text{DIC}(\text{SSEE} - \text{CPL}) = -3.27, \quad (17)$$

both favouring SSEE (lower DIC), consistent with the BIC and AIC of the main text.

Information criteria summary

Table 11: Bayesian model comparison from the three-model MCMC (total matter density in the background). $N_{\text{data}} = 16$ (13 BAO + 3 Planck). Negative Δ (BIC, AIC, DIC) favours SSEE relative to Λ CDM. Jeffreys criterion: $|\Delta| > 5 = \text{strong}$.

Criterion	SSEE	Λ CDM	CPL	$\Delta_{\text{SSEE}-\Lambda\text{CDM}}$
p_D	2.00	3.00	4.99	—
DIC	14.95	20.61	18.97	−5.66
BIC	16.49	22.92	22.85	−6.43
AIC	14.95	20.60	18.99	−5.65

Conclusion. Every information criterion favours SSEE: it fits DESI DR2 as well as Λ CDM and CPL while fixing w_0, w_a algebraically, so the parsimony of a two-parameter background is rewarded across BIC, AIC, and DIC. With the total matter density in the geometry the sound horizon matches Λ CDM and no background-structure penalty arises; the full CMB likelihood (Paper 3) independently favours SSEE at $\Delta\text{BIC} = -32.9$ (full `plik` MCMC).

B Nested Bayes Factor and Predictive Cross-Validation

This appendix provides two additional model-comparison criteria that are independent of the k -counting schemes discussed in Appendix A: the *Savage–Dickey density ratio* (exact Bayes factor for nested models) and a *predictive cross-validation* on the DESI DR2 BAO data.

B.1 Savage–Dickey density ratio

SSEE is nested within CPL: the SSEE dynamic sector is CPL evaluated at the algebraic point $(w_0, w_a) = (-0.840, -0.670)$. For nested models, the marginal Bayes factor for the null hypothesis $\mathcal{H}_0 : (w_0, w_a) = (-0.840, -0.670)$ versus the unconstrained CPL is given exactly by the Savage–Dickey density ratio [Verdinelli & Wasserman \[1995\]](#), [Trotta \[2008\]](#):

$$B_{\text{SSEE/CPL}} = \frac{p(w_0^S, w_a^S \mid \text{data}, \text{CPL})}{\pi(w_0^S, w_a^S \mid \text{CPL})} = 18.34, \quad \ln B = +2.91. \quad (18)$$

The CPL posterior is evaluated at (w_0^S, w_a^S) using a Scott-bandwidth Gaussian KDE of $N = 500\,000$ thinned CPL chain samples (columns 3–4 of the 5-dimensional CPL chain, shape 2.5×10^6). The marginal prior on (w_0, w_a) is the truncated Gaussian $\mathcal{N}(-1, 0.5^2) \times \mathcal{N}(0, 1^2)$ clipped to the MCMC sampling box $w_0 \in (-2.5, 0.5)$, $w_a \in (-3, 2)$; truncation factors are (0.9973, 0.9759), negligible corrections.

On the Jeffreys scale [Trotta \[2008\]](#) $\ln B = +2.91$ (> 2.3) constitutes *strong* evidence in favour of the SSEE dynamic sector within the CPL framework. The CPL posterior median ($w_0^{\text{CPL}} = -0.826 \pm 0.080$, $w_a^{\text{CPL}} = -0.557 \pm 0.314$) places the SSEE algebraic point at $(0.18\sigma, 0.36\sigma)$, consistent with the 0.24σ joint tension quoted in Section 6.2.

B.2 Ly- α predictive audit ($z = 2.33$)

The Ly- α point at $z = 2.33$ provides the sharpest direct test of SSEE’s background expansion. At $z \sim 2$ radiation is negligible and the ratio $D_H/r_d = c / (H_0 E(z) r_d)$ directly encodes the product $H_0 E(z) r_d$. We evaluate this quantity under two SSEE scenarios and compare to the DESI DR2 measurement $D_H/r_d = 8.632 \pm 0.101$ (Table 1).

Scenario A — raw ssee background ($s_m = 0.160050$, $r_d = 175.6$ Mpc). With the algebraic $H_0 = 67.962 \text{ km s}^{-1} \text{ Mpc}^{-1}$ and $s_m = 0.160050$, the CPL dark-energy fraction at $z = 2.33$ is $f_{\text{DE}}(2.33) = 0.648$, giving $E(2.33) = 2.540$ and $D_H(2.33) = 1737$ Mpc. The predicted ratio is:

$$\left. \frac{D_H}{r_d} \right|_{z=2.33}^{\text{SSEE, raw}} = \frac{1737}{175.6} = 9.89 \quad (12.5\sigma \text{ above DESI}), \quad (19)$$

demonstrating the category error directly: using the cold sector $s_m = 0.160050$ in $E(z)$ reduces matter domination at high z , lowers $H(z)$, and enlarges D_H to a 12.5σ discrepancy. This is the same misplacement that produced the spurious background tensions in superseded drafts; the correct total density (Scenario B) resolves it.

Scenario B — ω_m -direct CMB-sector background ($\Omega_{m,\text{CMB}} = 0.308881$, $r_d = 147.17$ Mpc). At $z = 2.33$ the full matter density sets the expansion, so $E(z)$ takes the algebraic value $\Omega_{m,\text{CMB}} = \omega_m/h^2 = 0.308881$ (free-streaming suppresses small-scale clustering, not the homogeneous density). Taking the sound horizon $r_d = 147.17$ Mpc directly from ω_m (CAMB; no $|w_0|$ mapping) gives $E(2.33) = 3.443$, $D_H(2.33) = 1281$ Mpc, and:

$$\left. \frac{D_H}{r_d} \right|_{z=2.33}^{\omega_m\text{-direct}} = \frac{1281}{147.17} = 8.71 \quad (0.7\sigma \text{ from DESI}). \quad (20)$$

This supersedes the earlier MIRA-mapped value ($\Omega_m = 0.320$, $D_H/r_d = 8.56$): the matter factor is dissolved (OP-8). The D_M/r_d counterpart at $z = 2.33$ is recovered within 0.1σ . The full 13-point diagonal fit gives $\chi^2 = 12.0$, matching ΛCDM ($\chi^2 = 12.0$): the two backgrounds describe DESI DR2 equally well. The largest SSEE residual is at $z = 0.51$ (D_M/r_d pull -2.1σ); ΛCDM shows a comparable largest residual ($+1.7\sigma$ at $z = 0.706$, D_H/r_d) with the same total χ^2 , confirming the residual pattern is a DESI DR2 feature rather than an SSEE-specific failure.

Implication. Eqs. (19)–(20) show that the high- z expansion requires the full CMB-sector matter density $\Omega_{m,\text{CMB}} = 0.308881$, where the cold sector counts as non-relativistic matter, rather than the EoS-sector value $s_m = 0.160050$: retaining it in $E(z)$ with $r_d = 147.17$ would predict $D_H/r_d = 11.8$ (a 31σ discrepancy). This is the same total-versus-dynamical distinction that sets the low- z sound horizon (§2.4), now with *no* matter factor (OP-8 dissolved) and with no second matter sector (retracted, Paper 6). The Savage–Dickey ratio in B.1 ($\ln B = +2.91$) then holds as a test of the *dynamical sector* under this ω_m -direct background.

Acknowledgements

The author thanks the DESI Collaboration for public DR2 data and P. Kroupa and collaborators for the IGIMF cluster re-analysis.

References

- Almeida Vallejo, M. E. (2026). *A Minimal-Parameter Framework for Cosmological Dynamics and Galaxy Cluster Mass Discrepancies via SSEE*. Zenodo preprint.
- Abdul-Karim, M., et al. (DESI Collaboration) (2025). DESI DR2 Results II. *Phys. Rev. D* **112**, 083515. arXiv:2503.14738.
- Planck Collaboration (2020). Planck 2018 results VI. *A&A* **641**, A6.
- Zhang, D., Zonoozi, A. H., & Kroupa, P. (2026). Revisiting the missing mass problem in MOND for nearby galaxy clusters. *Phys. Rev. D* **113**, 043027. arXiv:2602.06082.
- Foreman-Mackey, D., et al. (2013). *emcee*: The MCMC Hammer. *PASP* **125**, 306.
- Moresco, M., et al. (2022). Unveiling the universe with emerging cosmological probes. *Living Rev. Relativ.* **25**, 6.
- Moresco, M., & Marulli, F. (2017). Raising the bar: new constraints on $H(z)$. *MNRAS* **471**, L82.
- Eisenstein, D. J., & Hu, W. (1998). Baryonic features in the matter transfer function. *ApJ* **496**, 605.
- Aubourg, É., et al. (2015). Cosmological implications of baryon acoustic oscillation measurements. *Phys. Rev. D* **92**, 123516, arXiv:1411.1074.
- Almeida Vallejo, M. E. (2026). *SSEE Structural Constant Dictionary & Prediction Register*. Zenodo. <https://doi.org/10.5281/zenodo.20684908> Closed algebraic constant dictionary; parameter-free postdiction, no claim of temporal priority over public data.
- Clowe, D., et al. (2006). A direct empirical proof of the existence of dark matter. *ApJL* **648**, L109.
- Simionescu, A., et al. (2011). Baryons at the Edge of the X-ray-brightest Galaxy Cluster. *Science* **331**, 1576.
- Tchernin, C., et al. (2016). An exploration of the X-ray and SZ properties of galaxy clusters in the 0.1–0.4 redshift range. *A&A* **595**, A42.
- Merten, J., et al. (2011). Creation of cosmic structure in the complex galaxy cluster merger Abell 2744. *MNRAS* **417**, 333.
- Owers, M. S., et al. (2011). The Dissection of Abell 2744: A Rich Cluster Growing Through Major and Minor Mergers. *ApJ* **728**, 27.
- Spiegelhalter, D. J., Best, N. G., Carlin, B. P., & van der Linde, A. (2002). Bayesian measures of model complexity and fit. *J. Roy. Statist. Soc. B* **64**, 583.
- Verdinelli, I., & Wasserman, L. (1995). Computing Bayes factors using a generalization of the Savage–Dickey density ratio. *J. Am. Statist. Assoc.* **90**, 614.
- Trotta, R. (2008). Bayes in the sky: Bayesian inference and model selection in cosmology. *Contemp. Phys.* **49**, 71. arXiv:0803.4089.

- Riess, A. G., et al. (2022). A Comprehensive Measurement of the Local Value of the Hubble Constant with 1 HST Programs Using the Distance Ladder. *ApJL* **934**, L7. arXiv:2112.04510.
- Chevallier, M. & Polarski, D. 2001, Accelerating universes with scaling dark matter. *Int. J. Mod. Phys. D* **10**, 213. arXiv:gr-qc/0009008.
- Linder, E.V. 2003, Exploring the expansion history of the universe. *Phys. Rev. Lett.* **90**, 091301. arXiv:astro-ph/0208512.
- Perlmutter, S., et al. (1999). Measurements of Ω and Λ from 42 High-Redshift Supernovae. *ApJ* **517**, 565. arXiv:astro-ph/9812133.
- Riess, A. G., et al. (1998). Observational Evidence from Supernovae for an Accelerating Universe and a Cosmological Constant. *AJ* **116**, 1009. arXiv:astro-ph/9805200.
- Weinberg, D. H., et al. (2013). Observational probes of cosmic acceleration. *Phys. Rep.* **530**, 87. arXiv:1201.1084.
- Clifton, T., et al. (2012). Modified gravity and cosmology. *Phys. Rep.* **513**, 1. arXiv:1106.3999.
- Verde, L., Treu, T., & Riess, A. G. (2019). Tensions between the early and late Universe. *Nature Astron.* **3**, 891. arXiv:1907.10625.
- Kamionkowski, M., & Riess, A. G. (2023). The Hubble tension and early dark energy. *Ann. Rev. Nucl. Part. Sci.* **73**, 153. arXiv:2211.04492.
- Goodman, J., & Weare, J. (2010). Ensemble samplers with affine invariance. *Commun. Appl. Math. Comput. Sci.* **5**, 65.
- Schwarz, G. (1978). Estimating the dimension of a model. *Ann. Statist.* **6**, 461.
- Jeffreys, H. (1961). *Theory of Probability*, 3rd ed. Oxford University Press.
- Cooke, R. J., Pettini, M., & Steidel, C. C. (2018). One Percent Determination of the Primordial Deuterium Abundance. *ApJ*, 855, 102.
- DES Collaboration (2024). The Dark Energy Survey: Cosmology Results with 1500 deg² of Weak Gravitational Lensing and Galaxy Clustering (DES Y5). arXiv:2401.02929.
- Deffayet, C., Esposito-Farèse, G., & Vikman, A. (2010). Covariant Galileon. *Phys. Rev. D*, **79**, 084003.
- Gubitosi, G., Piazza, F., & Vernizzi, F. (2013). The Effective Field Theory of Dark Energy. *JCAP*, 2013(02), 032.
- Heymans, C., et al. (2021). KiDS-1000 Cosmology: Multi-probe weak gravitational lensing and spectroscopic galaxy clustering constraints. *A&A*, 646, A140.
- Abbott, T. M. C., et al. (DES Collaboration) (2022). Dark Energy Survey Year 3 Results: Cosmological Constraints from Galaxy Clustering and Weak Lensing. *Phys. Rev. D*, **105**, 023520.

- Eisenstein, D. J., et al. (2005). Detection of Baryon Acoustic Oscillations in the Correlation Function of a Sample of 46,748 Luminous Red Galaxies. *ApJ*, 633, 560.
- Jimenez, R., & Loeb, A. (2002). Constraining Cosmological Parameters Based on Relative Galaxy Ages. *ApJ*, 573, 37.
- Silk, J. (1968). Cosmic Black-Body Radiation and Galaxy Formation. *ApJ*, 151, 459.
- Hu, W., & Sugiyama, N. (1996). Small-Scale Cosmological Perturbations: An Analytic Approach. *ApJ*, 471, 542.
- Amendola, L. (2000). Coupled quintessence. *Phys. Rev. D*, 62, 043511.