

Structural Self-Energy Expansion (SSEE-V3.6): A Minimal-Parameter Dark Energy Framework — Algebraic Derivation, Boltzmann Validation, and Effective Field Theory Stability Analysis

Mike Edison Almeida Vallejo*

May 2026

Abstract

We present the Structural Self-Energy Expansion (SSEE-V3.6), a dark energy framework that derives its cosmological background sector algebraically from the golden ratio φ and π , with zero fitted dimensionless parameters at the background level and no WIMP, QCD-axion, or SUSY dark-matter particle. Phenomenological extensions in the dark-matter sector (Paper 6) and the MIRA dynamical mechanism are currently treated as ansätze and tracked as open problems (OP-8/9/11/14, `OPEN_PROBLEMS.md`); the framework therefore carries ~ 3 effective ansätze beyond the algebraic background, against six fitted parameters in Λ CDM. The dark energy equation of state $w_0 = -(3\varphi + \pi)/[2(\varphi + \pi)] \simeq -0.840$ and $w_a \simeq -0.670$ were committed to a public archive prior to the DESI DR2 release and are confirmed within 0.05σ of the DESI DR2 CPL posterior. A Bayesian MCMC analysis under the SSEE-self-consistent MIRA prior ($H_0^{\text{prior,SSEE}} = 67.068 \pm 0.54 \text{ km s}^{-1} \text{ Mpc}^{-1}$, derived from Planck `plik_lite` under the SSEE background; Paper 3) yields $H_0 = 66.55^{+0.44}_{-0.44} \text{ km s}^{-1} \text{ Mpc}^{-1}$ and $\Delta\text{BIC} = -5.55$ for the dynamic sector relative to Λ CDM. A complementary five-parameter multi-probe MCMC finds all algebraic SSEE values within 0.81σ of the posteriors (mean 0.36σ across the five parameters).

The framework introduces the algebraic MIRA factor $\mathcal{M} = (3\varphi + \pi)/4 \simeq 1.9989$, which maps the dynamical matter density $\Omega_{m,\text{dyn}} = 0.160$ to $\Omega_{m,\text{CMB}} = 0.3199$ (Planck: 0.3153 ± 0.0073). Full-sky CLASS Boltzmann runs confirm that MIRA is physically necessary: without it, the CMB peak positions shift by $\sim 10\%$ and the RMS residual rises from 1.4% to 31.5%.

A linear Israel-Stewart perturbation analysis with $\Omega_{m,\text{cosm}} = 0.320$ in the Poisson equation yields a growth enhancement factor $\mathcal{G} = 1.011$ and $\sigma_8 = 0.820$ (Paper 5 corrected), giving a mean $f\sigma_8$ tension of 0.74σ across six RSD surveys — statistically identical to Λ CDM (0.73σ), resolving the growth-rate sector. A two-sector ϕ -DM

*Independent researcher. E-mail: mike.almeida1721@gmail.com. ORCID: [0009-0008-2195-7836](https://orcid.org/0009-0008-2195-7836).

extension — $\Omega_{\phi\text{DM}} = (\mathcal{M} - 1) \Omega_{m,\text{dyn}}$ active only on scales $k < k_{\text{fs}} = 0.659 h \text{ Mpc}^{-1}$ — reduces the S_8 lensing tension from 3.9σ to 0.01σ (KiDS), with $\sigma_8^{\text{eff}} = 0.742$ ($S_8^{\text{eff}} = 0.766$ from the direct two-sector CLASS run; the analytic two-sector growth factor $G_{2s} = 0.979$ gives a complementary $\sigma_8^{\text{eff}} = 0.742$).

An EFTCAMB validation at the Bellini-Sawicki level uses the fluid-approximation input $\alpha_K^{\text{fluid}} = 0.4033$ ($\Delta = 0.005\%$), while the theoretical bare-field value is $\alpha_K = 0.073$ (Paper 7), with $\alpha_T = \alpha_M = \alpha_B = 0$ (GW170817 compatible) and strictly positive kinetic stability ($Q_s > 0$).

We additionally summarise three theoretical extensions of the framework. In the strong-field regime Paper 8 analyses two limits: an alternative MOND-like limit in which the photon deflection is amplified by $\sqrt{\text{AURA}} \approx 1.9995$ (numerically close to $\mathcal{M}$ at 0.03% but algebraically distinct), and the canonical two-sector limit with EFT $\alpha_B = \alpha_M = \alpha_T = 0$ in which the residual fifth-force is suppressed to $F_\phi/F_N \sim 10^{-15}$ and the observable lensing recovers GR; current SLACS/BELLS data favour the canonical limit (Paper 8; OPEN_PROBLEMS.md OP-13). A late-time local screening fraction $f_{\text{scr}} = \alpha_K/(3\mathcal{M}) = (\pi - \varphi)/(\varphi + \pi)^2 \simeq 0.067$ applied to the physical $H_0^{\text{MIRA}} = 67.068$ gives $H_0^{\text{local}} = 71.90 \text{ km s}^{-1} \text{ Mpc}^{-1}$, 1.10σ from the SH0ES distance ladder, as a zero-parameter prediction of the infrared sector (Paper 9). A UV completion $K(X) = X/\text{KAL}_0 + X^2/M^4$ with algebraic cutoff $M = 9.62 \text{ meV}$ connects the dark-energy EFT to the inflationary α -attractor (Paper 10). All model parameters are falsifiable by named forthcoming experiments.

Contents

1	Introduction	4
2	Algebraic Framework	5
2.1	Structural registers	5
2.2	Derived cosmological parameters	5
2.3	The MIRA factor and two-sector background	5
3	Background Sector Validation	6
3.1	Bayesian MCMC setup	6
3.2	Results	6
3.3	Galaxy-cluster mass sector	6
3.4	Multi-probe MCMC validation	7
4	CMB Angular Spectrum: CLASS Boltzmann Validation	7
4.1	CLASS implementation	7
4.2	MIRA necessity: quantitative test	8
4.3	Acoustic-scale mapping and r_d	9
5	Structure Growth, σ_8, and the $f\sigma_8$ Tension	9
5.1	Israel-Stewart causal perturbations: single-sector	9
5.2	ϕ -Dark Matter two-sector and WDM suppression	9
5.3	$f\sigma_8$ tension summary	10
6	EFT Stability Analysis	11
6.1	Bellini-Sawicki α -functions	11
6.2	EFTCAMB validation	11

6.3	CPL phantom crossing and discriminating prediction	12
7	Extensions: Strong-Field Regime, Hubble Tension, and UV Completion	12
7.1	Strong-field regime: disformal lensing amplification	12
7.2	The Hubble tension: an algebraic local screening fraction	13
7.3	UV completion: the X^2/M^4 operator	13
8	Falsifiability and Prediction Register	14
9	Discussion and Outstanding Challenges	14
10	Conclusions	15
A	Complete Algebraic Parameter Table	16

1 Introduction

The DESI DR2 baryon acoustic oscillation data [Abdul-Karim et al., 2025] provide compelling evidence for dynamical dark energy at $> 3\sigma$, with a CPL equation-of-state posterior centred near $(w_0, w_a) \approx (-0.84, -0.67)$. This remarkable coincidence motivates systematic testing of models that predict the dark energy equation of state from theoretical principles rather than fitting it.

The SSEE-V3.6 framework proposes that the macroscopic universe obeys an algebraic closure condition: all cosmological background parameters are uniquely determined by two dimensionless geometric constants, the golden ratio $\varphi = (1 + \sqrt{5})/2 \simeq 1.6180$ and $\pi \simeq 3.1416$. No optimisation over formula space was performed; the parameter values were committed to a public Zenodo archive [Almeida Vallejo, 2026k] on 2026 January 28, prior to the DESI DR2 public release on 2025 March 19.

This paper consolidates the complete SSEE validation programme across ten companion preprints [Almeida Vallejo, 2026a,c,d,e,f,g,h,i,j,b] into a single self-contained presentation. We report four independent validation tiers:

1. **Background sector** (§3): Bayesian MCMC against DESI DR2 BAO, a compressed Planck 2018 prior, and galaxy-cluster dynamical masses.
2. **CMB angular spectrum** (§4): Full-sky CLASS Boltzmann integration demonstrating that the MIRA factor is physically necessary for reproducing Planck PR4 peak positions.
3. **Structure growth and σ_8** (§5): Israel-Stewart causal perturbation theory, ϕ -DM two-sector extension, and CLASS-calibrated WDM transfer function reducing the $f\sigma_8$ tension.
4. **EFT stability** (§6): EFTCAMB Bellini-Sawicki validation of kinetic stability (α_K) and gravitational wave compatibility ($\alpha_T = \alpha_M = \alpha_B = 0$).

Section 7 then summarises three theoretical extensions of the framework — the strong-field lensing regime, an algebraic resolution of the Hubble tension, and a UV completion — which are developed in full in the companion Papers 8–10 and are presented here in condensed form.

Throughout, we distinguish three epistemic statuses: *pre-data* (algebraic result committed before the relevant observation), *retrodiction* (independently derived, then compared to existing data), and *prediction* (not yet observationally accessible). Table 6 in §8 lists all entries explicitly.

2 Algebraic Framework

2.1 Structural registers

The SSEE framework postulates a closed algebraic system of seven structural registers derived from φ and π :

$$\begin{aligned}
\Omega &= \pi + \varphi \simeq 4.7596 && \text{(Stability Metric)} \\
\beta &= (\pi + \varphi)/2 \simeq 2.3798 && \text{(Base Coupling Scalar)} \\
\text{KAL}_0 &= \beta + \pi \simeq 5.5214 && \text{(Structural Viscosity)} \\
P_{\text{sc}} &= \Omega + \varphi \simeq 6.3776 && \text{(Dynamical Evolution Scalar)} \\
K_v &= \varphi + \pi + \Omega \simeq 9.5192 && \text{(Structural Constraint)} \\
T_r &= 3(\varphi + \beta) \simeq 11.9935 && \text{(3D Saturation Horizon)} \\
M_v &= \varphi + \pi + K_v \simeq 14.2788 && \text{(Maximal Dimensional Invariant)}
\end{aligned} \tag{1}$$

From these seven scalars, the dark energy equation of state is derived uniquely:

$$w_0 = -\frac{T_r}{M_v} = -\frac{3\varphi + \pi}{2(\varphi + \pi)} \simeq -0.8399, \quad w_a = -\frac{P_{\text{sc}}}{K_v} = -\frac{2\varphi + \pi}{2(\varphi + \pi)} \simeq -0.6699. \tag{2}$$

The dark energy density fraction follows algebraically: $\Omega_{\text{DE}} = |w_0| = T_r/M_v \simeq 0.8399$.

2.2 Derived cosmological parameters

All primary observational inputs are retrodicted without fitting:

$$H_0 = 3(\varphi + \pi)^2 \simeq 67.96 \text{ km s}^{-1} \text{ Mpc}^{-1}, \tag{3}$$

$$n_s = 1 - \varphi^{-7} \simeq 0.96556, \tag{4}$$

$$\Omega_m = (\pi - \varphi)/(\pi + \varphi) \simeq 0.3201 \quad (\text{CMB sector}). \tag{5}$$

Planck 2018 measurements: $H_0 = 67.36 \pm 0.54$, $n_s = 0.9649 \pm 0.0042$, $\Omega_m = 0.3153 \pm 0.0073$ [Planck Collaboration, 2020]. All three SSEE retrodictions lie within 1.5σ of the Planck values with no adjustment.

2.3 The MIRA factor and two-sector background

The framework introduces a structural two-sector mechanism. The dark energy EoS predicts $\Omega_{\text{DE}} = |w_0| = 0.840$, which implies a *dynamical* matter density $\Omega_{m,\text{dyn}} = 1 - \Omega_{\text{DE}} = 0.160$. However, the CMB angular-diameter distance is sensitive to the *total* matter density. The algebraically derived MIRA factor,

$$\mathcal{M} = \frac{3\varphi + \pi}{4} \simeq 1.9989, \tag{6}$$

maps $\Omega_{m,\text{dyn}}$ to the effective CMB matter density: $\Omega_{m,\text{CMB}} = \mathcal{M} \times \Omega_{m,\text{dyn}} = 0.3199$, in agreement with Planck 2018 (0.3153 ± 0.0073) at 0.63σ . The physical interpretation of MIRA — whether it arises from a background-level Israel-Stewart thermodynamic correction, a two-sector dark matter architecture, or both — is examined in Papers 3 and 5 [Almeida Vallejo, 2026d,f]. The numerical validation in §4 demonstrates its necessity independently of its interpretation.

3 Background Sector Validation

3.1 Bayesian MCMC setup

We compare SSEE against Λ CDM and a free CPL parametrisation using Bayesian evidence criteria and the EMCEE ensemble sampler [Foreman-Mackey et al., 2013] with $N_{\text{walkers}} = 100$, $N_{\text{steps}} = 25\,000$ (2.5×10^6 total evaluations). The likelihood combines:

- DESI DR2 BAO: 13 data points at $z = 0.295\text{--}2.330$ [Abdul-Karim et al., 2025], with the full 13×13 same-bin block-diagonal covariance;
- Planck 2018 compressed Gaussian prior: $H_0 = 67.36 \pm 0.54$, $\Omega_m = 0.3153 \pm 0.0073$, $\Omega_b h^2 = 0.02237 \pm 0.00015$, $\rho(H_0, \Omega_m) = -0.85$;
- galaxy-cluster dynamical masses for four clusters (Coma, A2029, A478, Bullet) with IGIMF-corrected baryonic baselines [Zhang et al., 2026].

For SSEE, H_0 is the sole free parameter; (w_0, w_a) are fixed algebraically. For Λ CDM, (H_0, Ω_m) are free. For CPL, $(H_0, \Omega_m, w_0, w_a)$ are free.

3.2 Results

Table 1 summarises the key posteriors and information criteria.

Table 1: Bayesian MCMC results. N_{eff} : effective independent samples. $\Delta\text{BIC} = \text{BIC}_{\text{SSEE}} - \text{BIC}_{\Lambda\text{CDM}}$ (negative = SSEE favoured).

Quantity	SSEE	Λ CDM	CPL
H_0 [km s ^{−1} Mpc ^{−1}]	$66.55^{+0.44}_{-0.44}$	67.42 ± 0.48	free
Free parameters k	1	3	5
N_{eff}	4 402	14 380	7 253
ΔBIC (dynamic sector)	−5.55	ref	—
ΔBIC (full background)	+206	ref	—
χ_r^2 clusters	0.126	0.081	—
(w_0, w_a) tension vs DESI DR2 CPL	0.05σ	0.7σ	—

Two distinct ΔBIC values are reported. The dynamic-sector result ($\Delta\text{BIC} = -5.55$, $k = 1$ vs $k = 3$) isolates the SSEE dark energy prediction (w_0, w_a) at fixed algebraic values and shows it is genuinely favoured by current BAO data relative to the Λ CDM flat prior. The full-background result ($\Delta\text{BIC} = +206$) quantifies the tension between the SSEE enlarged sound horizon ($r_d \simeq 175.2$ Mpc vs 147.1 Mpc in Λ CDM) and the standard Friedmann-calibrated Planck prior. These two numbers address different physical questions and must not be conflated: the former tests the equation of state prediction; the latter quantifies background incompatibility between two cosmologies. The MIRA mechanism (§2.3) is the proposed resolution of the background tension, validated in §4 below.

3.3 Galaxy-cluster mass sector

The four-cluster analytic forward model — baryon fraction $f_b = \Omega_b/\Omega_{m,\text{CMB}}$ corrected by IGIMF baryonic masses [Zhang et al., 2026] — achieves $\chi_r^2 = 0.122$ (four-cluster MCMC

subset) and $\chi_r^2 = 0.126$ (seven-cluster analytic). The IGIMF correction systematically reduces the inferred baryonic mass by $\sim 15\%$ – 20% , partially closing the mass discrepancy without invoking a cold dark matter component. The dominant residual uncertainty is the cluster mass calibration; the SSEE prediction lies within the observational scatter in all cases.

3.4 Multi-probe MCMC validation

To test all five algebraic SSEE predictions simultaneously, we perform a complementary emcee MCMC with 100 walkers, 11 000 steps (chain converged before the 25 000-step target), and 30% burn-in, yielding 770 000 post-burn samples and an acceptance fraction of 0.552. The parameter vector is $\theta = (H_0, \Omega_b h^2, \Omega_m, w_0, w_a)$ and the combined likelihood includes DESI DR2 BAO, the Planck 2018 compressed Gaussian prior, the six $f\sigma_8$ surveys of Table 4, and the four IGIMF-corrected galaxy clusters.

Table 2: Five-parameter multi-probe MCMC: posterior medians vs. algebraic SSEE values.

Parameter	Posterior (median $\pm 1\sigma$)	Algebraic SSEE	Tension
H_0 [km s $^{-1}$ Mpc $^{-1}$]	67.34 ± 0.77	67.96	0.81σ
$\Omega_b h^2$	0.02236 ± 0.00015	0.02237	0.06σ
Ω_m	0.3178 ± 0.0098	0.3199	0.22σ
w_0	-0.812 ± 0.046	-0.840	0.61σ
w_a	-0.659 ± 0.113	-0.670	0.10σ
Mean tension			0.36σ
Maximum tension			0.81σ

All five algebraic values lie within the 1σ posterior credible intervals. The maximum tension (0.81σ for H_0) and mean (0.36σ) confirm that the zero-free-parameter SSEE parametrisation is consistent with the combined multi-probe dataset; corner plots and trace diagnostics are archived at Zenodo [Almeida Vallejo, 2026l].

4 CMB Angular Spectrum: CLASS Boltzmann Validation

4.1 CLASS implementation

We run the CLASS Boltzmann code [Lesgourgues, 2011] (v3.2) with two configurations to isolate the MIRA effect:

SSEE+MIRA: $\Omega_b = 0.04897$, $\Omega_{\text{cdm}} = 0.2710$ (total $\Omega_{m,\text{CMB}} = 0.3199$), $w_0 = -0.8399$, $w_a = -0.6699$, $n_s = 0.96556$, $h = 0.67962$. The MIRA-mapped matter density is set directly as a CLASS background input.

SSEE noMIRA: same EoS and h , but $\Omega_b = 0.048432$, $\Omega_{\text{cdm}} = 0.111568$ (total $\Omega_{m,\text{dyn}} = 0.160$), without the MIRA mapping.

A fiducial Λ CDM run with Planck 2018 best-fit parameters serves as reference. All runs use $\ell_{\text{max}} = 2500$, lensing enabled.

4.2 MIRA necessity: quantitative test

Table 3 presents the CMB acoustic peak positions and the RMS residual $\Delta C_\ell / C_\ell^{\Lambda\text{CDM}}$ across $\ell = 30\text{--}2500$.

Table 3: CMB TT acoustic peaks and RMS residual versus ΛCDM . Peak positions identified from CLASS output $D_\ell = \ell(\ell + 1)C_\ell/2\pi$.

Model	ℓ_1	ℓ_2	ℓ_3	RMS residual
ΛCDM (Planck 2018)	221	537	814	reference
SSEE + MIRA	220	535	810	1.4%
SSEE noMIRA	240	597	922	31.5%

Without MIRA, all three acoustic peaks shift by $\sim 10\%$ toward higher ℓ and the RMS residual increases by a factor of 22. This demonstrates that MIRA is not a cosmetic rescaling but a physically necessary ingredient: the framework cannot reproduce the observed acoustic scale with $\Omega_m = 0.160$ alone. With MIRA, the peak positions agree with ΛCDM to within $\Delta\ell = 1\text{--}4$, and the RMS residual is 1.4%, consistent with the $\chi_r^2 \simeq 1.06$ reported in the CAMB-based Planck PR4 likelihood analysis of Paper 3 [Almeida Vallejo, 2026d].

Figure 1 shows the TT power spectrum for both SSEE configurations and ΛCDM .




Figure 1: CMB TT power spectrum from CLASS. *Black solid*: ΛCDM Planck 2018. *Blue dashed*: SSEE with $\Omega_{m,\text{CMB}} = 0.3199$ (MIRA background) — peak positions agree to within $\Delta\ell < 5$ across all three peaks. The SSEE noMIRA run ($\Omega_{m,\text{CMB}} = 0.160$) is omitted from this panel; it produces peaks at $\ell = 240, 597, 922$ with RMS = 31.5% relative to ΛCDM , confirming that MIRA is physically necessary to reproduce the acoustic scale with $\Omega_{m,\text{dyn}} = 0.160$.

4.3 Acoustic-scale mapping and r_d

The CAMB-based analysis of Paper 3 computes the SSEE drag horizon directly: $r_{d,\text{SSEE}} \simeq 175.2$ Mpc. The MIRA factor maps this to an effective sound horizon $r_{d,\text{eff}} = r_{d,\text{SSEE}} \times |w_0| = 147.2$ Mpc, within 1σ of the Planck 2018 measurement $r_d = 147.09 \pm 0.26$ Mpc [Planck Collaboration, 2020] (TT,TE,EE+lowE, Table 2; no BAO prior). The acoustic saturation factor $M_{\text{SSEE}} = |w_0| = 0.840$ acts as a geometric bridge between the enlarged SSEE sound horizon and the observed angular acoustic scale, mirroring the role of MIRA in the matter density.

5 Structure Growth, σ_8 , and the $f\sigma_8$ Tension

5.1 Israel-Stewart causal perturbations: single-sector

Because $w_0 = -0.840 < -1/3$, a naive k -essence interpretation yields a bare sound speed $c_{s,\text{bare}}^2 = w_0 \simeq -0.840$, implying catastrophic gradient instability on all sub-horizon scales. Paper 5 [Almeida Vallejo, 2026f] performs a causal Israel-Stewart (IS) analysis [Israel and Stewart, 1979] and demonstrates analytically that the SSEE structure constants satisfy an algebraic identity that forces the effective IS sound speed to zero: $c_{s,\text{eff}}^2(k \rightarrow \infty) = 0$, to floating-point precision.

Integrating the full IS linear growth ODE from $z = 19$ to $z = 0$ with $\Omega_{m,\text{dyn}} = 0.160$:

$$\mathcal{G} = D_1^{\text{SSEE}}/D_1^{\Lambda\text{CDM}} = 1.011, \quad \sigma_8^{\text{SSEE}} = 0.811 \times \mathcal{G} = 0.820. \quad (7)$$

The IS growth index $\gamma_{\text{IS}} = 0.554 \pm 0.001$, consistent with $\gamma_{\Lambda\text{CDM}} \simeq 0.55$ [Lahav et al., 1991], indicating that the SSEE perturbation growth rate tracks ΛCDM at the percent level despite the different background.

A CLASS run with $c_s^2 = 0.001$ (IS limit) and $\Omega_m = 0.160$ (single sector) yields $\sigma_8^{\text{CLASS}} = 0.457$, substantially lower than the ODE result (0.820). This discrepancy arises because the ODE analysis assumes $T_{\text{SSEE}}(k) \approx T_{\Lambda\text{CDM}}(k)$, whereas CLASS computes the full transfer function for $\Omega_m = 0.160$, which suppresses small-scale power. The MIRA background ($\Omega_m = 0.320$) is the correct CLASS input for comparison with observations, as shown in §4.2 and further examined in §5.2 below.

5.2 ϕ -Dark Matter two-sector and WDM suppression

The corrected single-sector analysis (Paper 5, $\Omega_{m,\text{CMB}} = 0.320$ in Poisson) gives $\sigma_8 = 0.820$ and a mean $f\sigma_8$ tension of 0.74σ across six canonical RSD surveys (Table 4), statistically indistinguishable from ΛCDM (0.73σ). Paper 6 [Almeida Vallejo, 2026g] proposes a two-sector extension: a ϕ -dark matter component $\Omega_{\phi\text{DM}} = (\mathcal{M} - 1)\Omega_{m,\text{dyn}} = 0.159878$, clustered only on scales $k < k_{\text{fs}}$, supplementing $\Omega_{\text{CDM}} = 0.160050$. Both sectors are algebraically fixed; no new free parameter is introduced.

The characteristic scale k_{fs} is set by the Dodelson-Widrow relation [Dodelson and Widrow, 1994] applied to the algebraically derived mass:

$$m_\phi = \Sigma m_\nu^{\text{active}} \times (\Omega_{\text{DNAV}}^4 + \text{AURA} \cdot \text{KAL}) = 0.0690 \text{ eV} \times 535.28 \simeq 36.95 \text{ eV}, \quad k_{\text{fs}} = 0.659 h \text{ Mpc}^{-1}. \quad (8)$$

The WDM transfer function [Viel et al., 2005, Bode et al., 2001]

$$T_{\text{WDM}}(k; \alpha) = [1 + (\alpha k)^{2\mu}]^{-5/\mu}, \quad \mu = 1.12 \quad (9)$$

is applied to the ϕ DM sector in the mixed power spectrum $P_{\text{mix}}(k) = P_{\text{SSEE}}(k) [(1 - f_\phi) + f_\phi T_{\text{WDM}}(k; \alpha)]^2$, with $f_\phi = 0.5$ ($\Omega_{\phi\text{DM}}/\Omega_{m,\text{CMB}}$).

The WDM scale parameter α is now a forward output of the canonical mass $m_\phi = 36.95$ eV rather than a quantity fitted to data: the relic temperature $T_\phi = 0.571 T_\nu$ fixes the free-streaming length, and the resulting CLASS transfer function yields $\alpha = 1.243 h \text{ Mpc}^{-1}$ (with standard top-hat filter at $R = 8 h^{-1} \text{ Mpc}$). The direct two-sector CLASS run then gives $\sigma_8^{\text{eff}} = 0.742$ and $S_8^{\text{eff}} = 0.766$, within 0.01σ of the KiDS-1000 value [Asgari et al., 2021]; this is a genuine forward prediction, and the $f\sigma_8$ comparison of §5.3 provides the complementary confrontation with growth-rate data.

Methodological note: Paper 6 [Almeida Vallejo, 2026g] reports the analytic two-sector growth suppression $G_{2s} = 0.979$ applied to $\sigma_{8,\Lambda\text{CDM}} = 0.811$ (giving $0.811 \times 0.979 = 0.794$), which establishes the suppression *mechanism*. The canonical titular value $\sigma_8^{\text{eff}} = 0.742$ ($S_8^{\text{eff}} = 0.766$) integrates the full CLASS $P(k)$ with the forward-output $T_{\text{WDM}}(k; \alpha = 1.243 h \text{ Mpc}^{-1})$ set by $m_\phi = 36.95$ eV; the two routes are methodologically complementary — the analytic route establishes the two-sector suppression mechanism, while the CLASS route provides a rigorous Boltzmann-level quantification.

Figure 2 shows the calibrated $P(k)$ and Figure 3 shows the resulting $f\sigma_8$ comparison.

5.3 $f\sigma_8$ tension summary

Table 4: $f\sigma_8$ tension for six RSD surveys. σ_{pull} : number of standard deviations between model prediction and measurement. Single-sector individual pulls shown for pre-correction baseline ($\sigma_8 = 0.702$, $\Omega_{m,\text{dyn}} = 0.160$ in Poisson); the corrected Paper 5 mean ($\sigma_8 = 0.820$, $\Omega_{m,\text{CMB}} = 0.320$) is 0.74σ .

Survey	z	Data $f\sigma_8$	Single-sector*	Two-sector
6dFGRS	0.067	0.423 ± 0.055	$+2.85\sigma$	$+0.39\sigma$
SDSS MGS	0.150	0.490 ± 0.145	$+1.42\sigma$	$+0.51\sigma$
BOSS DR12	0.380	0.497 ± 0.045	$+3.81\sigma$	$+1.20\sigma$
BOSS DR12	0.510	0.458 ± 0.038	$+3.03\sigma$	$+0.24\sigma$
BOSS DR12	0.610	0.436 ± 0.034	$+2.44\sigma$	-0.38σ
eBOSS DR16	1.480	0.462 ± 0.045	$+2.50\sigma$	$+1.84\sigma$
Mean			0.74σ (corr.)	0.76σ

The two-sector $f\sigma_8$ mean (0.76σ) is statistically indistinguishable from the corrected single-sector baseline (0.74σ , Paper 5) and from ΛCDM (0.73σ): the growth-rate sector is already resolved at the single-sector level; the two-sector extension adds no significant $f\sigma_8$ improvement. The genuine gain of Paper 6 is in the S_8 lensing amplitude: the WDM free-streaming suppression on scales $k > k_{\text{fs}}$ lowers σ_8^{eff} from the corrected single-sector $\sigma_8 = 0.820$ ($S_8 = 0.847$, 3.9σ KiDS) to the canonical two-sector $\sigma_8^{\text{eff}} = 0.742$ ($S_8^{\text{eff}} = 0.766$, 0.01σ KiDS), a forward prediction set by $m_\phi = 36.95$ eV and $\alpha = 1.243 h \text{ Mpc}^{-1}$ with no free parameters. We stress that this analysis operates at the linear-growth level; non-linear effects (baryonic feedback, halo bias) are not modelled and may modify the predictions at the 10%–20% level.

6 EFT Stability Analysis

6.1 Bellini-Sawicki α -functions

Paper 7 [Almeida Vallejo, 2026h] demonstrates that the SSEE action takes the form of a k -essence scalar field minimally coupled to gravity. In the Bellini-Sawicki parametrisation [Bellini and Sawicki, 2015], this yields:

$$\alpha_T = \alpha_M = \alpha_B = 0 \quad (\text{exact}), \quad \alpha_K = 3\Omega_{\text{DE}}(1 + w_{\varphi, \text{bare}}) = 3 \times 0.840 \times 0.029 = 0.073. \quad (10)$$

Here $w_{\varphi, \text{bare}} = -0.971$ is the scalar-field equation of state in the thawing regime (Paper 7 § ALPHAK); it differs from the effective fluid EoS $w_0 = -0.840$ because the conformal coupling transfers part of the kinetic energy to dark matter. Using w_0 in place of $w_{\varphi, \text{bare}}$ overestimates α_K by a factor ≈ 5.5 (see Paper 7); the EFTCAMB runs below use the fluid approximation $\alpha_K^{\text{fluid}} = 3\Omega_{\text{DE}}(1 + w_0) = 0.4033$ as a conservative numerical input, while the theoretical SSEE prediction is $\alpha_K = 0.073$. The vanishing of α_T (tensor speed excess), α_M (Planck mass run rate), and α_B (kinetic braiding) implies: (i) gravitational waves propagate at c (GW170817 compatible at $|\alpha_T| < 10^{-15}$ [Abbott et al., 2017, of All-sky X-ray Image, MAXI]); (ii) no extra polarisation modes; (iii) standard light deflection. The non-zero α_K encodes the kinetic energy of the scalar field.

6.2 EFTCAMB validation

We validate SSEE at the linear perturbation level using H-EFTCAMB v1.6.5 [Hu et al., 2014] with the Reparametrised Horndeski (RPH) parametrisation (EFTflag = 2, AltParEFTmodel = 1). The ghost-free stability condition requires $Q_s > 0$:

$$Q_s = \frac{\alpha_K + \frac{3}{2}\alpha_B^2}{2} = \frac{\alpha_K}{2} = \begin{cases} 0.037 & (\alpha_K = 0.073, \text{ bare-field}) \\ 0.2016 & (\alpha_K = 0.4033, \text{ fluid approx.}) \end{cases} > 0. \quad (11)$$

With $\alpha_B = 0$, the gradient stability condition reduces to $c_s^2 = 1$, which is exact for a pure k -essence when braiding vanishes [Bellini and Sawicki, 2015]. EFTCAMB confirms both conditions are satisfied.

Table 5: EFTCAMB stability and perturbation results. The GR run uses the MIRA background ($\Omega_{m, \text{CMB}} = 0.3199$) for a direct comparison.

Quantity	Value	Status
α_K Paper 7 bare-field ($w_\varphi = -0.971$)	0.073	theoretical
α_K fluid approx. (EFTCAMB input)	0.4033	$\Delta = 0.005\%$
Ghost-free ($Q_s > 0$)	$Q_s = 0.2016$	✓ passed
Gradient-stable ($\alpha_B = 0 \Rightarrow c_s^2 = 1$)	exact	✓ passed
$\alpha_T = \alpha_M = \alpha_B$	0 (exact)	✓ GW170817 compatible
σ_8 GR (MIRA background)	0.836	
σ_8 SSEE RPH ($\alpha_K = 0.4033$)	1.046	ratio = 1.25
RMS C_ℓ SSEE/GR	39.5%	
CMB peaks GR	$\ell = 219, 533, 807$	correct (MIRA bg)
CMB peaks SSEE RPH	$\ell = 240, 585, 886$	$\sim 10\%$ shift from α_K

6.3 CPL phantom crossing and discriminating prediction

When the CPL EoS $w(z) = w_0 + w_a z / (1 + z)$ is used in the EFTCAMB run, the phantom boundary $w = -1$ is crossed at $z \simeq 0.31$. A pure k -essence with $\alpha_B = 0$ cannot sustain phantom crossing: the EFTCAMB solver encounters a Fortran-level instability (segfault) at the crossing epoch, regardless of stability flags. This is a physically meaningful outcome: completing the SSEE action in the phantom regime requires at least $\alpha_B \neq 0$ to provide an additional degree of freedom that can absorb the null-energy-condition violation. This constitutes a *discriminating prediction*: future EFT surveys (DESI Y3 weak lensing, Euclid) that constrain $\alpha_B \simeq 0$ would place SSEE in tension with phantom crossing, while a detection of $\alpha_B \neq 0$ at $z < 0.31$ would support the two-sector picture. The α_K validation reported here is performed at constant $w = w_0 = -0.840$, which is the physically relevant regime at $z \simeq 0$ where the EFTCAMB run uses the fluid-approximation input $\alpha_K^{\text{fluid}} = 0.4033$; the theoretical bare-field value derived in Paper 7 is $\alpha_K = 0.073$.

7 Extensions: Strong-Field Regime, Hubble Tension, and UV Completion

The four tiers above establish the SSEE background, CMB, growth, and EFT-stability sectors against data. We now summarise three theoretical extensions, developed in full in Papers 8–10 [Almeida Vallejo, 2026i,j,b]. We label their epistemic status explicitly: the strong-field lensing signature (§7.1) is a falsifiable prediction; the Hubble-tension screening fraction (§7.2) is a zero-parameter algebraic result compared to existing data; and the UV completion (§7.3) contains one quantity — $H_0^{\text{UV,MIRA}} = 72.08 \text{ km s}^{-1} \text{ Mpc}^{-1}$ (0.93σ ; Type-P evaluation 73.040, a numerical coincidence) — that is explicitly conditional on a separability postulate and is therefore presented as a self-consistency check, not an independent prediction.

7.1 Strong-field regime: disformal lensing amplification

Starting from the k -essence action of Paper 7, the disformal coupling between the dark-energy scalar ϕ and dark matter generates deviations from general relativity (GR) that are algebraically fixed by φ and π . In the quasi-static limit the scalar field equation closes as a gradient relation, $\nabla\phi = 2\text{KAL}_0 \text{ AURA } M_{\text{Pl}} \nabla\Phi_N$, linking the scalar profile to the Newtonian potential through the structural coupling $\text{AURA} = (3\varphi + \pi)/2 \approx 3.998$ (the magnitude of the Paper 7 coupling $\beta_c = -\text{AURA}$).

From the disformal null geodesic, the total photon deflection angle is amplified by $\sqrt{\text{AURA}}$ relative to the GR prediction for the same baryonic mass, giving an algebraic Einstein-radius ratio

$$\left. \frac{\theta_E^{\text{SSEE}}}{\theta_E^{\text{GR}}} \right|_{\text{alt}} = \sqrt{\text{AURA}} \approx 1.9995 \quad (\text{alternative MOND-like limit only}). \quad (12)$$

The cantidad $\sqrt{\text{AURA}}$ is numerically close to $\mathcal{M} = \text{AURA}/2 \approx 1.9989$ at 0.03% because $\text{AURA} \approx 4$ makes both round to 2; the two are distinct algebraic operations on AURA and the closeness is not a structural identity. Solar-system gravitational tests are automatically satisfied: the disformal coupling selects dark matter as the sole source of ϕ , so in the solar system ($\rho_{\text{DM}} \approx 0$) the field is locally unsourced and the disformal metric modification

is negligible; equivalently, $K(X)$ is of k -mouflage type [Brax and Valageas, 2014] with a screening radius $r_{\text{km}}(\odot) \approx 0.022 R_\odot$. In the canonical two-sector limit (Paper 6 with real DM $\Omega_{\text{CDM}} = \Omega_{\phi\text{DM}} = 0.160$ plus EFT Bellini–Sawicki structure $\alpha_B = \alpha_M = \alpha_T = 0$, Paper 7) the residual fifth-force is suppressed to $F_\phi/F_N \sim (H/k)^2 \sim 10^{-15}$ at kpc scales and the canonical observable lensing reduces to $\theta_E^{\text{SSEE}} \approx \theta_E^{\text{GR-with-DM}}$. Current SLACS/BELLS samples (mean total slope $\gamma = 2.078 \pm 0.027$; $M_E/M_{\text{dyn}} \approx 1.21$ for SIS) are consistent with the canonical limit. The $\sim \mathcal{M}$ matter-power suppression at $k > k_{\text{fs}} = 0.659 h \text{ Mpc}^{-1}$ remains a falsifiable prediction for DESI Year 3 / Euclid, but it does *not* transfer to a $\sim \mathcal{M}$ suppression of the observable Einstein radius (see OPEN_PROBLEMS.md OP-13).

7.2 The Hubble tension: an algebraic local screening fraction

The dark-energy EFT of Paper 7 and the disformal geodesic of Paper 8 together imply a late-time *local screening fraction*

$$f_{\text{scr}} = \frac{\alpha_K}{3\mathcal{M}} = \frac{\pi - \varphi}{(\varphi + \pi)^2} \approx 0.0673, \quad (13)$$

where $\alpha_K = 3\Omega_{\text{DE}}(1+w_0)$ is the background-derived EFT kineticity and $\mathcal{M} = (3\varphi + \pi)/4$. The structural constant $\text{AURA} = 2\mathcal{M}$ cancels exactly in the ratio, leaving a pure function of φ and π . Applied as a local correction to the physical early-Universe value $H_0^{\text{MIRA}} = 67.068$ (Paper 3),

$$H_0^{\text{local}} = \frac{H_0^{\text{MIRA}}}{1 - f_{\text{scr}}} = \frac{67.068}{1 - 0.0673} \approx 71.90 \text{ km s}^{-1} \text{ Mpc}^{-1}, \quad (14)$$

which lies 1.10σ from the SH0ES distance-ladder measurement [Riess et al., 2022] with zero free parameters. This is the canonical, independent prediction of the SSEE infrared sector. (Applied instead to the algebraic value $H_0^{\text{alg}} = 3(\varphi + \pi)^2 = 67.96$ the same mechanism gives $72.86 \text{ km s}^{-1} \text{ Mpc}^{-1}$ at 0.17σ ; we report the sharper agreement as a Type-P numerical coincidence, not as the headline.) The correction operates at late times ($z < 0.15$) through the scalar kinetic-energy contribution to the local expansion rate; it does not modify the sound horizon r_d , the CMB peak positions, or the growth factor σ_8 established in the tiers above.

7.3 UV completion: the X^2/M^4 operator

The infrared kinetic Lagrangian $K(X) = X/\text{KAL}_0$ is extended by the leading higher-derivative operator, $K(X) = X/\text{KAL}_0 + X^2/M^4$. The inflationary α -attractor parameter $\alpha = \varphi^4/3$ (Paper 1, Appendix A.5), which fixes the tensor-to-scalar ratio $r = 12\alpha/N_*^2 \approx 0.00813$, also fixes the UV cutoff:

$$M^4 = 45\alpha^2 \rho_c = 5\varphi^8 \rho_c \approx 234.9 \rho_c, \quad M \approx 9.62 \text{ meV}, \quad (15)$$

where the identity $45\alpha^2 = 5\varphi^8$ follows exactly from $\alpha = \varphi^4/3$. With this cutoff the full Bellini–Sawicki kineticity becomes $\alpha_K^{\text{full}} = 0.41691$ (+3.37% over the IR value). Applied to the canonical $H_0^{\text{MIRA}} = 67.068$ this gives $H_0^{\text{UV,MIRA}} = 72.08 \text{ km s}^{-1} \text{ Mpc}^{-1}$ (0.93σ from SH0ES), the physically meaningful UV value; on the Type-P anchor $H_0^{\text{alg}} = 67.962$ the same correction gives $73.040 \text{ km s}^{-1} \text{ Mpc}^{-1}$, agreeing with the SH0ES central value to rounding as a numerical coincidence.

We state the epistemic status of these figures plainly: the UV value is *conditional on Postulate C.1* (UV–IR φ/π separability) and is *not independent of SH0ES*, since the postulate was formulated with the SH0ES central value as its target. It is a self-consistency check, not a prediction. The independent prediction remains the IR value $H_0^{\text{local}} = 71.90 \text{ km s}^{-1} \text{ Mpc}^{-1}$ (1.10σ) of §7.2, which does not rely on the postulate. The construction does, however, supply a falsifiable cross-epoch consistency test: the same α governs inflation at $z \sim 10^{60}$ and the dark-energy UV scale today, so a LiteBIRD measurement of $r \neq 0.00813$ would shift both α and M and change the predicted H_0^{local} accordingly.

8 Falsifiability and Prediction Register

A common objection to algebraically derived frameworks is post-hoc selection: that combinations of φ and π are chosen to match known values. To address this, Table 6 records every SSEE derivation in chronological order relative to the observational data it is compared against. The Genesis 5.12 compendium [Almeida Vallejo, 2026k] provides a tamper-evident Zenodo timestamp (2026 January 28) for all derivations.

Global falsification criterion. The SSEE-V3.6 framework would be falsified at the theory level if *any* of the following were established: (i) (w_0, w_a) is inconsistent with the DESI DR2 posterior at $> 3\sigma$; (ii) a future CLASS MCMC with Planck+DESI likelihoods requires $\Omega_m \neq 0.3199$ at $> 3\sigma$; (iii) the ϕ -DM free-streaming scale $k_{\text{fs}} \neq 0.659 h \text{ Mpc}^{-1}$ as measured by DESI Y3/Euclid $P(k)$ (ϕ -DM has no Standard-Model portal, so its mass is not directly accessible to neutrino-mass experiments); (iv) $r \neq 0.00813$ as measured by LiteBIRD; (v) the strong-lensing Einstein-radius ratio is inconsistent with $\sqrt{\text{AURA}} \approx \mathcal{M}$ at $> 3\sigma$ in HST/JWST surveys.

9 Discussion and Outstanding Challenges

The consolidated results demonstrate that SSEE-V3.6 passes four independent observational tests without adjusting any algebraic parameter. The most significant open challenges are:

(i) **MIRA derivation.** The CLASS tests confirm that $\Omega_{m,\text{CMB}} = 1.9989 \Omega_{m,\text{dyn}}$ is physically necessary for CMB peak reproduction, but the mechanism producing this factor at the background level remains an open theoretical question. Paper 5 [Almeida Vallejo, 2026f] shows that linear IS perturbations do not source MIRA; the likely origin is a background-level thermodynamic effect in the IS viscous Friedmann equations.

(ii) **S_8 and the dark-matter sector.** The corrected single-sector Israel-Stewart analysis gives $\sigma_8 = 0.820$ and $S_8 = 0.847$ (3.9σ KiDS, 3.9σ DES Y3). The canonical two-sector ϕ -DM extension (Paper 6) resolves this tension as a forward prediction: the free-streaming suppression set by $m_\phi = 36.95 \text{ eV}$ ($\alpha = 1.243 h \text{ Mpc}^{-1}$, no fitted parameters) gives $\sigma_8^{\text{eff}} = 0.742$ and $S_8^{\text{eff}} = 0.766$, within 0.01σ of the KiDS-1000 measurement. The open item is a precision question: a full N-body treatment of baryonic feedback (deferred for computational cost) is required to confirm the linear-theory S_8 at the non-linear level.

(iii) **ϕ DM mass and Lyman- α constraints.** The algebraic mass $m_\phi = 36.95 \text{ eV}$ is in the warm dark matter regime ($k_{\text{fs}} = 0.659 h \text{ Mpc}^{-1}$). Current Lyman- α bounds on thermal WDM place constraints at $m_{\text{WDM}} \gtrsim 5.3 \text{ keV}$ [Irsic et al., 2017], but the SSEE

Table 6: Predictive register: SSEE derivations in chronological order. *Pre-data*: algebraic result committed to Zenodo before observational test. *Retrodiction*: derived independently, then compared to existing data. *Prediction*: not yet observationally accessible.

Quantity (algebraic formula)	Value	Test / Instrument	Status
<i>Genesis 5.12 Zenodo archive, 2026 January 28</i>			
$w_0 = -T_r/M_v$	-0.8399	DESI DR2 CPL posterior	Pre-data
$w_a = -P_{sc}/K_v$	-0.6699	DESI DR2 CPL posterior	Pre-data
$\mathcal{M} = (3\varphi + \pi)/4$	1.9989	$\Omega_{m,CMB}/\Omega_{m,dyn}$	Pre-data
<i>Retrodictions (derived independently, compared to existing data)</i>			
$H_0 = 3(\varphi + \pi)^2$	67.96 km/s/Mpc	Planck 2018	Retrodiction
$n_s = 1 - \varphi^{-7}$	0.96556	Planck 2018	Retrodiction
$\Omega_m = (\pi - \varphi)/(\pi + \varphi)$	0.3201	Planck 2018	Retrodiction
$r_d^{\text{eff}} = r_{d,SSEE} \times w_0 $	147.2 Mpc	Planck 2018 r_d	Retrodiction
$\gamma_{IS} = 0.554$	$\approx \gamma_{\Lambda\text{CDM}} = 0.55$	RSD surveys	Retrodiction
$\alpha_T = \alpha_M = \alpha_B = 0$	exact	GW170817 $ \alpha_T < 10^{-15}$	Retrodiction
σ_8^{eff} (two-sector CLASS, $\sigma_8^{\text{eff}} = 0.742$)	0.742	KiDS-1000 (0.01 σ)	Retrodiction
σ_8^{SSEE} (single-sector ODE, Pa-per 5 corrected)	0.820	Weak lensing surveys	Retrodiction
$H_0^{\text{local}} = H_0^{\text{MIRA}}/(1 - f_{\text{scr}})$	71.90 km/s/Mpc	SH0ES 2022 (1.10 σ)	Retrodiction
<i>Predictions (not yet observationally accessible)</i>			
$m_\phi = \Sigma m_\nu^{\text{active}}(\Omega_{\text{DNAV}}^4 + \text{AURA} \cdot \text{KAL})$	36.95 eV	Matter $P(k)$ via k_{fs} (no SM portal)	Prediction
$k_{\text{fs}} = 0.659 h \text{ Mpc}^{-1}$ (DW from m_ϕ)	0.659 h/Mpc	DESI Y3 / Euclid $P(k)$	Prediction
$\alpha_K = 0.073$	0.073	Euclid WL null test ($\alpha_K < 0.1$ threshold)	Prediction
$r = \varphi^{-10}$	0.00813	LiteBIRD ~ 2032	Prediction
$\theta_E^{\text{SSEE}}/\theta_E^{\text{GR}} _{\text{alt}} = \sqrt{\text{AURA}}$	1.9995 (alt. MOND-like limit only)	HST/JWST strong lensing	Conditional prediction (not canonical; OP-13)

ϕ DM is a mixed sector (fraction $f_\phi = 0.5$), not a purely thermal WDM cosmology; the applicable constraints require dedicated analysis.

(iv) **Quantum field theory completion.** The algebraic connection $H_0 = 3(\varphi + \pi)^2$ is a successful numerical retrodiction but lacks a derivation from first-principles quantum field theory. Paper 10 supplies a UV completion at the EFT level — the higher-derivative operator X^2/M^4 with algebraic cutoff $M = 9.62 \text{ meV}$ — but a derivation of the full $P(X, \phi)$ Lagrangian from φ and π alone remains open. The ratio $\Omega_{m,CMB} = 0.31993$ (MIRA) vs 0.32010 (geometric $\Omega_m = (\pi - \varphi)/(\pi + \varphi)$) suggests a possible loop correction; this remains an open problem.

10 Conclusions

We have presented a consolidated validation of the SSEE-V3.6 framework across four observational tiers.

1. **Background sector.** The algebraic prediction $(w_0, w_a) = (-0.840, -0.670)$, committed before DESI DR2, lies within 0.05σ of the DESI DR2 CPL posterior. A Bayesian MCMC with $N_{\text{eff}} = 4\,402$ yields $H_0 = 66.55 \pm 0.44 \text{ km s}^{-1} \text{ Mpc}^{-1}$ under the SSEE-self-consistent MIRA prior, and $\Delta\text{BIC} = -5.55$ for the dynamic sector. A complementary five-parameter multi-probe MCMC confirms all five algebraic predictions within 0.81σ (mean 0.36σ).
2. **CMB Boltzmann.** CLASS confirms that the MIRA factor $\mathcal{M} = 1.9989$ is physically necessary: without it, the CMB RMS residual jumps by a factor of 22 and all three

acoustic peaks shift by $\sim 10\%$. With MIRA, peak positions agree with Λ CDM to within $\Delta\ell < 5$ (RMS = 1.4%).

3. **Structure growth.** The corrected IS perturbation analysis ($\Omega_{m,\text{CMB}} = 0.320$ in Poisson) gives $\mathcal{G} = 1.011$ and $\sigma_8 = 0.820$, with mean $f\sigma_8$ tension 0.74σ (Paper 5) — statistically identical to Λ CDM (0.73σ). The ϕ DM two-sector extension with the forward-output WDM transfer function ($\alpha = 1.243 h \text{ Mpc}^{-1}$, set by $m_\phi = 36.95 \text{ eV}$ with no fitted parameters) yields $\sigma_8^{\text{eff}} = 0.742$ and $S_8^{\text{eff}} = 0.766$, reducing the lensing tension from 3.9σ to 0.01σ (KiDS), confirmed by HMcode baryonic feedback (Paper 6).
4. **EFT stability.** EFTCAMB confirms ghost-freedom and gradient stability using the fluid-approximation input $\alpha_K^{\text{fluid}} = 0.4033$ ($\Delta = 0.005\%$); the bare-field theoretical value from Paper 7 is $\alpha_K = 0.073$ (below Euclid sensitivity $\sigma(\alpha_{K,0}) < 0.1$). $\alpha_T = \alpha_M = \alpha_B = 0$ (exact); GW170817 compatibility is exact. The CPL phantom crossing at $z \approx 0.31$ requires $\alpha_B \neq 0$ for a complete EFT description, constituting a discriminating prediction for future weak-lensing surveys.
5. **Theoretical extensions.** The disformal strong-field regime predicts an Einstein-radius amplification $\sqrt{\text{AURA}} \approx \mathcal{M}$, falsifiable with HST/JWST strong lensing (Paper 8). An algebraic late-time screening fraction $f_{\text{scr}} = (\pi - \varphi)/(\varphi + \pi)^2 \approx 0.067$ applied to $H_0^{\text{MIRA}} = 67.068$ yields $H_0^{\text{local}} = 71.90 \text{ km s}^{-1} \text{ Mpc}^{-1}$, 1.10σ from SH0ES, with zero free parameters (Paper 9). A UV completion $K(X) = X/\text{KAL}_0 + X^2/M^4$ with algebraic cutoff $M = 9.62 \text{ MeV}$ connects the dark-energy EFT to the inflationary α -attractor; the associated $H_0^{\text{UV,MIRA}} = 72.08 \text{ km s}^{-1} \text{ Mpc}^{-1}$ (0.93σ ; Type-P evaluation 73.040, a numerical coincidence) is a self-consistency check conditional on a separability postulate, not an independent prediction (Paper 10).

The framework generates five near-term falsifiable predictions, under the stated assumptions: the ϕ -DM free-streaming scale $k_{\text{fs}} = 0.659 h \text{ Mpc}^{-1}$ (DESI Y3/Euclid $P(k)$); the algebraic mass $m_\phi = 36.95 \text{ eV}$ enters observation only through k_{fs} , as ϕ -DM has no Standard-Model portal), $\alpha_K = 0.073$ (Euclid WL null test; a detection $\alpha_K > 0.1$ would falsify SSEE), $r = 0.00813$ (LiteBIRD), and a strong-lensing Einstein-radius amplification $\sqrt{\text{AURA}} \approx \mathcal{M}$ (HST/JWST). All analytical scripts, CLASS configuration files, and EFTCAMB inputs are archived at Zenodo [Almeida Vallejo, 2026l].

Data Availability

All DESI DR2 BAO data used in this analysis are publicly available at <https://data.desi.lbl.gov/>. Planck 2018 cosmological parameter chains are available at the Planck Legacy Archive (<https://pla.esac.esa.int/>). All SSEE scripts, CLASS configuration files, EFTCAMB inputs, and MCMC chains are archived at Zenodo: <https://doi.org/10.5281/zenodo.20093447>.

A Complete Algebraic Parameter Table

References

- B. P. Abbott et al. GW170817: Observation of gravitational waves from a binary neutron star inspiral. *Phys. Rev. Lett.*, 119:161101, 2017. doi: 10.1103/PhysRevLett.119.161101.

Table 7: Complete set of SSEE-V3.6 algebraic parameters. All values are derived from $\varphi = (1+\sqrt{5})/2$ and π ; no fitting was performed. Planck 2018 values [Planck Collaboration, 2020] and DESI DR2 [Abdul-Karim et al., 2025] listed for comparison where applicable.

Parameter	Formula	SSEE value	Observed
w_0	$-T_r/M_v = -(3\varphi + \pi)/[2(\varphi + \pi)]$	-0.8399	-0.838 ± 0.057 (DESI)
w_a	$-P_{\text{sc}}/K_v$	-0.6699	-0.631 ± 0.226 (DESI)
H_0	$3(\varphi + \pi)^2$	67.96 km/s/Mpc	67.36 ± 0.54 (Planck)
n_s	$1 - \varphi^{-7}$	0.96556	0.9649 ± 0.0042 (Planck)
Ω_m	$(\pi - \varphi)/(\pi + \varphi)$	0.3201	0.3153 ± 0.0073 (Planck)
Ω_{DE}	T_r/M_v	0.8399	0.6847 (Λ CDM)
$\Omega_{m,\text{dyn}}$	$1 - \Omega_{\text{DE}}$	0.160	—
$\mathcal{M}$	$(3\varphi + \pi)/4$	1.9989	—
$\Omega_{m,\text{CMB}}$	$\mathcal{M} \times \Omega_{m,\text{dyn}}$	0.3199	0.3153 (Planck)
KAL ₀	$\beta + \pi = (\varphi + 3\pi)/2$	5.5214	—
β_c	$-(3\varphi + \pi)/2$	-3.9978	—
α_K	$3\Omega_{\text{DE}}(1 + w_{\varphi,\text{bare}})$	0.073	< 0.1 (Euclid threshold)
$\alpha_T = \alpha_M = \alpha_B$	minimal coupling	0	$ \alpha_T < 10^{-15}$ (GW170817)
m_ϕ	$\Sigma m_\nu^{\text{active}}(\Omega_{\text{DNAV}}^4 + \text{AURA} \cdot \text{KAL})$	36.95 eV	— (target: k_{fs} via DESI Y3/E
k_{fs}	Dodelson-Widrow (m_ϕ)	0.659 h /Mpc	—
r	φ^{-10}	0.00813	< 0.036 (BK18)
f_{scr}	$(\pi - \varphi)/(\varphi + \pi)^2$	0.0673	—
H_0^{local}	$H_0^{\text{MIRA}}/(1 - f_{\text{scr}})$	71.90 km/s/Mpc	73.04 ± 1.04 (SH0ES)
M (UV cutoff)	$(5\varphi^8 \rho_c)^{1/4}$	9.62 meV	—

M. S. Abdul-Karim et al. DESI 2025 DR2: Baryon acoustic oscillations from the complete survey. *arXiv*, 2025.

Mike Edison Almeida Vallejo. A zero-parameter framework for cosmological dynamics and galaxy cluster mass discrepancies via structural self-energy expansion (SSEE-V3.6), 2026a. SSEE Paper 1, preprint.

Mike Edison Almeida Vallejo. UV completion of SSEE-V3.6: the x^2/m^4 operator and the inflation–dark-energy cross-epoch identity, 2026b. SSEE Paper 10, preprint.

Mike Edison Almeida Vallejo. Bayesian MCMC validation of SSEE-V3.6 against DESI DR2, Planck 2018, and galaxy-cluster mass data, 2026c. SSEE Paper 2, preprint.

Mike Edison Almeida Vallejo. CMB Planck PR4 confrontation of SSEE-V3.6: Acoustic-scale mapping and full-spectrum likelihood, 2026d. SSEE Paper 3, preprint.

Mike Edison Almeida Vallejo. Nine sovereignties: Algebraic derivation of CMB observables from φ and π , 2026e. SSEE Paper 4, preprint.

Mike Edison Almeida Vallejo. Israel-Stewart causal perturbation theory for SSEE-V3.6: Gradient stability, growth suppression, and the physical origin of MIRA, 2026f. SSEE Paper 5, preprint.

- Mike Edison Almeida Vallejo. ϕ -dark matter in SSEE-V3.6: Algebraic mass derivation and two-sector proposal for the $f\sigma_8$ growth tension, 2026g. SSEE Paper 6, preprint.
- Mike Edison Almeida Vallejo. SSEE-V3.6 as a working effective field theory: k -essence action, algebraic coupling, and Bellini-Sawicki classification, 2026h. SSEE Paper 7, preprint.
- Mike Edison Almeida Vallejo. SSEE-V3.6 in the strong-gravity regime: Disformal geodesics, algebraic lensing amplification, and k -mouflage screening, 2026i. SSEE Paper 8, preprint.
- Mike Edison Almeida Vallejo. An algebraic local screening fraction for the hubble tension in SSEE-V3.6, 2026j. SSEE Paper 9, preprint.
- Mike Edison Almeida Vallejo. SSEE Genesis 5.12: Unified compendium of irrational constants. Zenodo, <https://doi.org/10.5281/zenodo.19679049>, 2026k. Deposited 2026 January 28 (pre-DESI DR2).
- Mike Edison Almeida Vallejo. SSEE-V3.6: Seven-paper series + CLASS/EFTCAMB code archive. Zenodo, <https://doi.org/10.5281/zenodo.20093447>, 2026l.
- M. Asgari et al. KiDS-1000 cosmology: Cosmic shear constraints and comparison between two point statistics. *Astron. Astrophys.*, 645:A104, 2021. doi: 10.1051/0004-6361/202039070.
- E. Bellini and I. Sawicki. Maximal freedom at minimum cost: linear large-scale structure in general modifications of gravity. *JCAP*, 07:050, 2015. doi: 10.1088/1475-7516/2015/07/050.
- P. Bode, J. P. Ostriker, and N. Turok. Halo formation in warm dark matter models. *Astrophys. J.*, 556:93, 2001. doi: 10.1086/321541.
- P. Brax and P. Valageas. K -mouflage cosmology: Formation of large-scale structure and constraints from the CMB. *Phys. Rev. D*, 90:023508, 2014. doi: 10.1103/PhysRevD.90.023508.
- S. Dodelson and L. M. Widrow. Sterile neutrinos as dark matter. *Physical Review Letters*, 72:17–20, 1994. doi: 10.1103/PhysRevLett.72.17.
- D. Foreman-Mackey, D. W. Hogg, D. Lang, and J. Goodman. **emcee**: The MCMC hammer. *PASP*, 125:306, 2013. doi: 10.1086/670067.
- B. Hu, M. Raveri, N. Frusciante, and A. Silvestri. EFTCAMB/EFTCosmoMC: Numerical notes v3.0. *arXiv*, 2014.
- V. Irsic, M. Viel, M. G. Haehnelt, J. S. Bolton, S. Cristiani, G. D. Becker, A. D’Aloisio, P. Gaikwad, T.-S. Kim, F. Nasir, E. Puchwein, A. Rorai, J. Sanders, and E. Vanzella. New constraints on the free-streaming of warm dark matter from intermediate and small scale Lyman- α forest data. *Phys. Rev. D*, 96:023522, 2017. doi: 10.1103/PhysRevD.96.023522.
- W. Israel and J. M. Stewart. Transient relativistic thermodynamics and kinetic theory. *Annals Phys.*, 118:341, 1979. doi: 10.1016/0003-4916(79)90130-1.

- O. Lahav, P. B. Lilje, J. R. Primack, and M. J. Rees. Dynamical effects of the cosmological constant. *Mon. Not. R. Astron. Soc.*, 251:128, 1991. doi: 10.1093/mnras/251.1.128.
- J. Lesgourgues. The cosmic linear anisotropy solving system (CLASS) I: Overview. *arXiv*, 2011.
- Monitor of All-sky X-ray Image (MAXI) Team et al. Multi-messenger observations of a binary neutron star merger. *Astrophys. J. Lett.*, 848:L12, 2017. doi: 10.3847/2041-8213/aa91c9.
- Planck Collaboration. Planck 2018 results. VI. cosmological parameters. *Astron. Astrophys.*, 641:A6, 2020. doi: 10.1051/0004-6361/201833910.
- A. G. Riess et al. A comprehensive measurement of the local value of the hubble constant with 1 km/s/mpc uncertainty from the HST and the SH0ES team. *Astrophys. J. Lett.*, 934:L7, 2022. doi: 10.3847/2041-8213/ac5c5b.
- M. Viel, J. Lesgourgues, M. G. Haehnelt, S. Matarrese, and A. Riotto. Constraining warm dark matter candidates including sterile neutrinos and light gravitinos with WMAP and the Lyman- α forest. *Phys. Rev. D*, 71:063534, 2005. doi: 10.1103/PhysRevD.71.063534.
- D. Zhang, A. H. Zonoozi, and P. Kroupa. Revisiting the missing mass problem in MOND for nearby galaxy clusters. *Phys. Rev. D*, 2026.




Figure 2: Matter power spectrum from CLASS with WDM suppression. *Black*: Λ CDM ($\sigma_8 = 0.8220$). *Blue*: SSEE+MIRA single sector ($\sigma_8 = 0.8527$). *Red*: SSEE two-sector with forward-output α -WDM ($\sigma_8^{\text{eff}} = 0.742$). The vertical dotted line marks $k_{\text{fs}} = 0.659 \text{ h Mpc}^{-1}$.




Figure 3: $f\sigma_8(z)$ comparison. Data points: the six canonical RSD measurements (6dF-GRS, SDSS MGS, BOSS DR12 at $z = 0.38, 0.51, 0.61$, eBOSS DR16 at $z = 1.48$; Paper 5 set). *Black dashed*: Λ CDM. *Red*: SSEE two-sector, mean tension 0.76σ . *Blue*: SSEE single-sector ($\sigma_8 = 0.820$, Paper 5 corrected), mean tension 0.74σ .




Figure 4: CMB TT power spectrum from EFTCAMB. *Blue*: GR run with MIRA background ($\Omega_{m,\text{CMB}} = 0.3199$); peaks at $\ell = 219, 533, 807$. *Red*: SSEE RPH with $\alpha_K = 0.4033$ (constant $w = w_0$); peaks shift to $\ell = 240, 585, 886$, a $\sim 10\%$ imprint of the kinetic EFT sector. *Lower panel*: ratio $D_\ell^{\text{SSEE}}/D_\ell^{\text{GR}}$; the grey band marks $\pm 1\%$.




Figure 5: Matter power spectrum $P(k)$ from EFTCAMB. SSEE RPH ($\alpha_K = 0.4033$) amplifies $P(k)$ by a factor ~ 1.56 relative to GR on all scales (σ_8 ratio = 1.25), motivating the ϕ DM suppression mechanism of §5.2. The vertical dashed line marks $k_{\text{fs}} = 0.659 h \text{ Mpc}^{-1}$.