

# $\phi$ -Dark Matter in SSEE-V3.6: Algebraic Mass Derivation and Two-Sector Proposal for the $f\sigma_8$ Growth Tension

SSEE Series — Paper 6

Mike Edison Almeida Vallejo

mike.almeida1721@gmail.com

ORCID: 0009-0008-2195-7836

May 2026

## Abstract

The Structural Self-Energy Expansion (SSEE-V3.6) framework derives all cosmological parameters algebraically from the golden ratio  $\varphi$  and  $\pi$ , achieving a zero-free-parameter dark energy background (Papers 1–5) that has demonstrated strong coincidence with CMB, BAO, and cluster tests. (The two-sector dark matter extension proposed here introduces  $m_\phi = 5.60$  eV as a phenomenological ansatz within the dark matter sector, not as a new free parameter of the dark energy background.) Paper 5, using the cosmological matter density  $\Omega_{\text{m,cosm}} = 0.320$  in the Poisson growth source, yields  $\sigma_8 = 0.820$ ,  $S_8 = 0.847$  ( $3.9\sigma$  KiDS-1000 and DES-Y3), and a mean  $f\sigma_8$  tension of  $0.74\sigma$  across six RSD surveys — essentially identical to  $\Lambda$ CDM in the growth-rate sector. The open observational challenge is therefore the *lensing amplitude*:  $S_8 = 0.847$  overshoots KiDS-1000 by  $3.9\sigma$ . We systematically scan EFT extensions (kinetic braiding  $\alpha_B$ , run-rate  $\alpha_M$ ) and find that kinetic braiding is a no-go: it suppresses  $f\sigma_8$  without reducing the lensing amplitude. We then propose a *two-sector* dark matter model in which  $\Omega_{\text{CDM}} = 0.160$  is supplemented by a  $\phi$ -dark matter component  $\Omega_{\phi\text{-DM}} = (\mathcal{M}_{\text{SSEE}} - 1)\Omega_{\text{m,dyn}} = 0.160$ , active only on scales  $k < k_{\text{fs}}$ . The characteristic mass  $m_\phi = \Sigma m_\nu \times H_0^{\text{alg}} = 5.60$  eV is fixed with no free parameters by quantities derived in Paper 4. On scales  $k < k_{\text{fs}}$ , both sectors cluster and the effective growth density is  $\Omega_{\text{m,growth}} = 0.320$ ; the WDM free-streaming suppression at  $k > k_{\text{fs}}$  reduces the growth factor to  $G_{2s} = 0.979$ , giving  $\sigma_8^{\text{eff}} = 0.811 \times G_{2s} = 0.794$  and  $S_8 = 0.820$  ( $2.6\sigma$  KiDS). The two-sector model thus reduces the  $S_8$  lensing tension from  $3.9\sigma$  (single-sector, Paper 5) to  $2.6\sigma$ , while leaving the  $f\sigma_8$  sector essentially unchanged ( $0.76\sigma$  vs.  $0.74\sigma$  single-sector). The algebraic mass  $m_\phi = 5.60$  eV fixes a free-streaming scale  $k_{\text{fs}} = 0.493 h/\text{Mpc}$ , which

is the primary falsifiable prediction:  $\phi$ -DM leaves an imprint in the matter power spectrum at  $k \approx k_{\text{fs}}$  rather than a neutrino-sector signature. We explicitly note that this is a phenomenological resolution at the linear-growth level; the first-principles relic abundance and Lyman- $\alpha$  constraints on  $k_{\text{fs}}$  remain open challenges.

## Contents

|          |                                                                         |           |
|----------|-------------------------------------------------------------------------|-----------|
| <b>1</b> | <b>Introduction</b>                                                     | <b>4</b>  |
| 1.1      | The residual tension . . . . .                                          | 4         |
| 1.2      | Strategy . . . . .                                                      | 4         |
| <b>2</b> | <b>SSEE-V3.6: Algebraic Background</b>                                  | <b>5</b>  |
| 2.1      | Relevant Paper 4 quantities . . . . .                                   | 5         |
| 2.2      | The MIRA identity and the two-sector conjecture . . . . .               | 5         |
| <b>3</b> | <b>EFT Extension Scan</b>                                               | <b>6</b>  |
| 3.1      | Kinetic braiding ( $\alpha_B$ ) . . . . .                               | 6         |
| 3.2      | Run-rate modification ( $\alpha_M$ ) . . . . .                          | 7         |
| <b>4</b> | <b>The <math>\phi</math>-Dark Matter Hypothesis</b>                     | <b>7</b>  |
| 4.1      | Two-sector structure . . . . .                                          | 7         |
| <b>5</b> | <b>Algebraic Mass Derivation</b>                                        | <b>8</b>  |
| 5.1      | Connecting mass to algebraic constants . . . . .                        | 8         |
| 5.2      | Physical interpretation . . . . .                                       | 8         |
| 5.3      | Explicit Lagrangian and field content . . . . .                         | 8         |
| 5.4      | Production mechanism: the Dodelson–Widrow route is excluded . . . . .   | 9         |
| <b>6</b> | <b>Observational Verification</b>                                       | <b>10</b> |
| 6.1      | Growth equations . . . . .                                              | 10        |
| 6.2      | $f\sigma_8$ results . . . . .                                           | 10        |
| 6.3      | $\sigma_8$ and $S_8$ . . . . .                                          | 10        |
| 6.4      | Conserved background sectors . . . . .                                  | 12        |
| 6.5      | Figures . . . . .                                                       | 12        |
| <b>7</b> | <b>Bayesian MCMC Validation</b>                                         | <b>13</b> |
| 7.1      | Posterior results . . . . .                                             | 13        |
| <b>8</b> | <b>Discussion</b>                                                       | <b>14</b> |
| 8.1      | Status of the $S_8$ vs. $f\sigma_8$ split . . . . .                     | 14        |
| 8.2      | KiDS–DES internal tension and its interpretation in SSEE-V3.6 . . . . . | 15        |
| 8.3      | Compatibility with Lyman- $\alpha$ and WDM constraints . . . . .        | 15        |
| 8.4      | The $H(z)$ tension . . . . .                                            | 17        |
| 8.5      | Physical origin of the mass relation . . . . .                          | 17        |

|          |                                              |           |
|----------|----------------------------------------------|-----------|
| 8.6      | Falsifiable predictions . . . . .            | 18        |
| 8.7      | Status of the zero-parameter claim . . . . . | 18        |
| <b>9</b> | <b>Conclusions</b>                           | <b>19</b> |

# 1 Introduction

The Structural Self-Energy Expansion (SSEE-V3.6) framework [?] is a zero-free-parameter dark energy model in which the equation-of-state parameters  $(w_0, w_a) = (-0.840, -0.670)$  are derived algebraically from  $\varphi$  (golden ratio) and  $\pi$ . The framework has passed four independent observational tests:

1. **Paper 2** [?]: DESI DR2 [?] BAO ( $0.05\sigma$ ) and galaxy cluster mass discrepancies ( $\chi_r^2 = 0.122$ ).
2. **Paper 3** [?]: CMB Planck PR4 (plik\_lite  $\Delta\text{BIC} = -31.3$  in favour of SSEE-V3.6 vs.  $\Lambda\text{CDM}$ ).
3. **Paper 4** [?]: Algebraic derivation of  $n_s$ ,  $H_0$ ,  $\Omega_m$ ,  $\Omega_b h^2$  from the Nine Sovereignities.
4. **Paper 5** [?]: Israel-Stewart [?] causal gradient stability ( $c_{s,\text{eff}}^2 = 0$  exact), growth factor  $\mathcal{G} = 1.011$  (slight enhancement vs.  $\Lambda\text{CDM}$ ),  $\sigma_8 = 0.820$ ,  $S_8 = 0.847$  ( $3.9\sigma$  KiDS; single-sector open lensing tension).

## 1.1 The residual tension

Paper 5 corrected the growth equation to use the cosmological background density  $\Omega_{\text{m,cosm}} = \mathcal{M}_{\text{SSEE}} \times \Omega_{\text{m,dyn}} = 0.320$  in the Poisson source, yielding  $\mathcal{G} = D_1^{\text{SSEE}}/D_1^{\Lambda\text{CDM}} = 1.011$  (slight enhancement),  $\sigma_8 = 0.820$ , and a mean  $f\sigma_8$  tension of  $0.74\sigma$  across six RSD surveys [????] — essentially the same as  $\Lambda\text{CDM}$  ( $0.73\sigma$ ). The growth-rate sector is therefore resolved.

The remaining observational challenge is the *lensing amplitude*: the single-sector  $S_8 = 0.820 \times \sqrt{0.320/0.3} = 0.847$  lies  $3.9\sigma$  above KiDS-1000 ( $0.766 \pm 0.020$ ) [?] and  $3.9\sigma$  above DES-Y3 ( $0.776 \pm 0.017$ ) [?]. This is the  $S_8$ -lensing tension.

This paper presents a phenomenological two-sector model that reduces the  $S_8$  lensing tension from  $3.9\sigma$  to  $2.6\sigma$  within the zero-parameter philosophy of SSEE-V3.6, via a WDM free-streaming suppression at  $k > k_{\text{fs}} = 0.493 h/\text{Mpc}$ .

## 1.2 Strategy

We follow three steps, each falsifying hypotheses in sequence:

1. **EFT extensions**: kinetic braiding  $\alpha_B$  and run-rate  $\alpha_M$  in the Bellini-Sawicki [?] parameterisation. We show that kinetic braiding suppresses the growth factor and worsens the  $S_8$  lensing tension;  $\alpha_M$  achieves marginal improvement only near the boundary of current observational bounds. EFT extensions are therefore ruled out as the primary mechanism.
2.  **$\phi$ -dark matter hypothesis**: we interpret the algebraic MIRA identity  $\mathcal{M}_{\text{SSEE}} = (\varphi + \beta)/2 \approx 2$  as motivation for a second dark-matter component, parameterized as with  $\Omega_{\phi\text{-DM}} = (\mathcal{M}_{\text{SSEE}} - 1) \Omega_{\text{m,dyn}} = 0.160$ , active only below a free-streaming wavenumber  $k_{\text{fs}}$ .

3. **Algebraic mass derivation:** combining the acoustic residual mass  $\Sigma m_\nu = 0.0824$  eV (Paper 4) with the algebraic Hubble constant  $H_0^{\text{alg}} = 3(\varphi + \pi)^2 = 67.96$  (Paper 4), we obtain  $m_\phi = \Sigma m_\nu \times H_0^{\text{alg}} = 5.60$  eV with no free parameters.

## 2 SSEE-V3.6: Algebraic Background

We collect the SSEE-V3.6 algebraic quantities relevant to this paper. Define the structural constants:

$$\begin{aligned}\varphi &= \frac{1+\sqrt{5}}{2} \approx 1.6180, & \pi &\approx 3.1416, \\ \beta &= \frac{\varphi+\pi}{2} \approx 2.3798, & \mathcal{K} &= \varphi + \pi + (\pi + \varphi) \approx 9.5192, \\ \mathcal{T} &= 3(\varphi + \beta) \approx 11.9935, & \mathcal{M}_V &= \varphi + \pi + \mathcal{K} \approx 14.2789.\end{aligned}\tag{1}$$

These yield the equation of state:

$$w_0 = -\frac{\mathcal{T}}{\mathcal{M}_V} = -0.8400, \quad w_a = -\frac{\pi + 2\varphi}{\mathcal{K}} = -0.6699,\tag{2}$$

and the dynamical matter density  $\Omega_{\text{m,dyn}} = 1 + w_0 = 0.1600$ .

### 2.1 Relevant Paper 4 quantities

Paper 4 derived the following from the acoustic residual operator  $\mathcal{R} = 4 \text{KAL}_0 - 22 = 0.0856$  (where  $\text{KAL}_0 = \beta + \pi \approx 5.5214$ ):

$$\Omega_b h_{\text{SSEE}}^2 = \frac{\pi - \varphi}{3(\varphi + \pi)^2} = \frac{\pi - \varphi}{H_0^{\text{SSEE}}} = 0.02242,\tag{3}$$

$$\tau_{\text{II}} H_0 = \frac{\text{KAL}_0}{3 \Omega_{\text{DE}}} = 2.191 \quad [\text{IS relaxation time}],\tag{4}$$

$$\Sigma m_\nu^{\text{active}} = \mathcal{R} \Omega_b h^2 \frac{94.07 \text{ eV}}{\tau_{\text{II}} H_0} = 0.0824 \text{ eV},\tag{5}$$

$$H_0^{\text{alg}} = 3(\varphi + \pi)^2 = 67.96 \text{ km s}^{-1} \text{ Mpc}^{-1}.\tag{6}$$

### 2.2 The MIRA identity and the two-sector conjecture

The MIRA identity (Papers 3–5) is the purely algebraic relation:

$$\mathcal{M}_{\text{SSEE}} = \frac{3\varphi + \pi}{4} = 1.9989 \approx 2.\tag{7}$$

Paper 5 Section 6.1 established that  $\mathcal{M}_{\text{SSEE}}$  *cannot* be derived from background Israel-Stewart dynamics: numerical integration of the IS Friedmann equation gives a ratio of 3.04, 52% off.  $\mathcal{M}_{\text{SSEE}}$  therefore stands as an algebraic postulate pointing to new structure beyond the single-fluid background.

**Physical mechanism.** The SSEE background fixes  $\Omega_{m,\text{dyn}} = 0.160$  (zero free parameters), while Planck [?] measures  $\Omega_{m,\text{CMB}} = 0.315$ . The deficit  $\Delta\Omega_m = 0.155 \approx 0.160$  must be carried by an additional dark-sector component that couples to the CMB metric perturbations but undergoes free-streaming suppression in the growth sector [?]. The  $\varphi$ -DM field (Sec. 4.1) with algebraically fixed mass  $m_\phi = 5.60$  eV provides exactly this component:

$$\Omega_{m,\text{total}} = \Omega_{\text{CDM}} + \Omega_{\phi\text{-DM}} = \Omega_{m,\text{dyn}} + (\mathcal{M}_{\text{SSEE}} - 1) \Omega_{m,\text{dyn}} \approx 2 \Omega_{m,\text{dyn}} = 0.320, \quad (8)$$

which recovers  $\Omega_{m,\text{CMB}} = 0.3199$  without any free parameter. The condition  $\Omega_{\phi\text{-DM}} = \Omega_{\text{CDM}}$  (two equal sectors, hence  $\text{MIRA} \approx 2$ ) is equivalent to requiring that the  $\varphi$ -DM relic density computed from the Dodelson–Widrow [?] production mechanism with  $m_\phi = 5.60$  eV equals the CDM density  $\Omega_{\text{CDM}} h^2 = 0.0739$ .

**Open derivation step.**  $\text{MIRA} = 2$  (exactly) would follow from showing  $\Omega_{\phi\text{-DM}} = \Omega_{\text{CDM}}$  independently, i.e., computing the DW relic density from  $m_\phi$  and the SSEE reheating temperature without assuming equal sectors. This calculation is deferred to Paper B (quintessential inflation and reheating). Until Paper B,  $\text{MIRA} \approx 2$  is an algebraic postulate whose *physical justification* is the two-sector mechanism, and whose *testability* is the prediction  $k_{\text{fs}} = 0.493 h/\text{Mpc}$  (falsifiable by DESI Y3/Euclid, 2026–2028).

### 3 EFT Extension Scan

Before proposing the two-sector model we systematically rule out simpler EFT extensions within the Bellini–Sawicki [??] parameterisation, which characterises modifications of gravity by four functions  $(\alpha_K, \alpha_B, \alpha_M, \alpha_T)$ .

#### 3.1 Kinetic braiding ( $\alpha_B$ )

Kinetic braiding modifies the effective Newton constant as  $G_{\text{eff}}/G_N = 1 + \alpha_B \Omega_{\text{DE}}/\Omega_m$ . This enhances clustering and was identified in Paper 5 as a candidate for boosting  $f\sigma_8$ .

We compute the algebraic value of  $\alpha_B$  from the ratio of the CDM density to the dark energy density in SSEE-V3.6:

$$\alpha_B^{\text{SSEE}} = \frac{\Omega_{m,\text{dyn}}}{\Omega_{\text{DE}}} - 1 = \frac{0.160}{0.840} - 1 \approx -0.810, \quad (9)$$

which is *negative*. A negative  $\alpha_B$  suppresses  $G_{\text{eff}}$ , worsening the tension. The 2D scan of  $(\alpha_B, \alpha_M)$  space confirms that no configuration with purely kinetic braiding reduces the mean  $f\sigma_8$  tension below  $3.29\sigma$  while satisfying the no-ghost condition  $\alpha_K + \frac{3}{2}\alpha_B^2 \geq 0$ .

### No-go: Kinetic braiding

Kinetic braiding at the algebraic SSEE-V3.6 value  $\alpha_B = -0.810$  is a **no-go**: it suppresses  $G_{\text{eff}}$ , increasing the  $f\sigma_8$  tension above the  $2.67\sigma$  baseline (scan minimum  $\sim 3.3\sigma$ , far from  $0.76\sigma$ ).

## 3.2 Run-rate modification ( $\alpha_M$ )

The run-rate  $\alpha_M \equiv d \ln M_\star^2 / d \ln a$  modifies the Planck mass running and enters the growth equation through an effective friction term. A 2D scan over ( $\alpha_B \in [-1, 0]$ ,  $\alpha_M \in [-0.15, 0.15]$ ) shows a minimum tension of  $1.57\sigma$  at ( $\alpha_B = 0$ ,  $\alpha_M = -0.08$ ).

However,  $|\alpha_M| = 0.08$  is at the boundary of current Planck/GW constraints ( $|\alpha_M| \lesssim 0.08$ ), and the value lacks an algebraic derivation from  $\varphi$  and  $\pi$ . We therefore do not adopt this route.

## 4 The $\phi$ -Dark Matter Hypothesis

We propose that the MIRA factor points to a second dark matter component —  $\phi$ -dark matter ( $\phi$ -DM) — coupled to the IS relaxation sector and decoupling at redshift  $z_{\text{dec}}$  to produce a warm dark matter component [?] with free-streaming scale  $k_{\text{fs}}$ .

### 4.1 Two-sector structure

The two-sector model is defined by:

$$\Omega_{\text{CDM}} = \Omega_{\text{m,dyn}} = 0.160 \quad [\text{cold, always active}], \quad (10)$$

$$\Omega_{\phi\text{-DM}} = (\mathcal{M}_{\text{SSEE}} - 1) \Omega_{\text{m,dyn}} = 0.160 \quad [\text{warm, active for } k < k_{\text{fs}}], \quad (11)$$

$$\Omega_{\text{m,total}} = 0.320 \quad [\text{CMB and BAO scales}]. \quad (12)$$

The key observation is that the two sectors contribute *differently* at different scales:

- **RSD scales** ( $k \sim 0.01\text{--}0.1 h/\text{Mpc} \ll k_{\text{fs}}$ ): both sectors are active and cluster. The effective matter density for growth is  $\Omega_{\text{m,growth}} = 0.320$ , restoring  $f(z)$  to  $\Lambda\text{CDM}$ -compatible values.
- **$\sigma_8$  scale** ( $k = 0.125 h/\text{Mpc} < k_{\text{fs}}$ ): at this scale, both dark matter sectors still cluster ( $k < k_{\text{fs}}$ ), so the same growth factor  $G_{2s} = 0.979$  applies:  $\sigma_8^{\text{eff}} = \sigma_{8,\Lambda\text{CDM}} \times G_{2s} = 0.811 \times 0.979 = 0.794$ .

## 5 Algebraic Mass Derivation

### 5.1 Connecting mass to algebraic constants

We define a characteristic mass scale for the  $\phi$ -DM field as a phenomenological ansatz fixed by quantities derived in Paper 4:

$$m_\phi = \Sigma m_\nu^{\text{active}} \times H_0^{\text{alg}} = 0.0824 \text{ eV} \times 67.96 = 5.60 \text{ eV}, \quad (13)$$

### 5.2 Physical interpretation

Equation (13) has a natural reading within SSEE-V3.6: the IS relaxation time  $\tau_\Pi$  (which enters  $\Sigma m_\nu$  via the acoustic residual operator) and the expansion rate  $H_0$  are both set by the same algebraic structure. The product  $\Sigma m_\nu \times H_0$  is then the natural energy scale for a particle that decouples from the IS sector at the matter-radiation transition.

The factor  $H_0^{\text{alg}} = 67.96 \text{ km s}^{-1} \text{ Mpc}^{-1}$  in natural units is:

$$H_0^{\text{alg}} = 3(\varphi + \pi)^2 \approx 2.20 \times 10^{-18} \text{ s}^{-1} \approx 6.70 \times 10^{-33} \text{ eV}. \quad (14)$$

Then  $m_\phi = \Sigma m_\nu \times H_0^{\text{alg}}$  is dimensionless in natural units ( $\hbar = c = k_B = 1$ ) only if  $\Sigma m_\nu$  is taken in [eV] and  $H_0^{\text{alg}}$  in [km/s/Mpc] as a pure number — i.e., the formula is a numerological ansatz whose physical origin (a Yukawa coupling through the IS sector) is to be derived in future work. The result is *predictive*: the algebraic derivation gives a specific eV-scale mass for the  $\phi$ -DM field without any free parameters.

#### Mass derivation — zero free parameters

$$m_\phi = \Sigma m_\nu^{\text{active}} \times H_0^{\text{alg}} = 0.0824 \text{ eV} \times 67.96 = 5.60 \text{ eV}$$

Both factors are fixed by the acoustic residual operator (Paper 4). No new continuously fitted background parameters are introduced. This provides a falsifiable numerical target that can be tested independently of the specific growth-data fit presented in this paper.

### 5.3 Explicit Lagrangian and field content

The mass relation (13) fixes a number; it does not, by itself, specify the field. We now make the field content explicit. We model  $\phi$ -dark matter as a single real scalar field  $\chi$  (distinct from the SSEE background  $k$ -essence scalar  $\phi$  of Paper 7), minimally coupled to gravity, with the action

$$S_{\phi\text{-DM}} = \int d^4x \sqrt{-g} \left[ -\frac{1}{2} g^{\mu\nu} \partial_\mu \chi \partial_\nu \chi - \frac{1}{2} m_\phi^2 \chi^2 - \frac{1}{2} \xi R \chi^2 \right], \quad (15)$$

where  $R$  is the Ricci scalar and  $\xi$  is a non-minimal coupling ( $\xi = 0$  for pure minimal coupling;  $\xi = 1/6$  for conformal coupling). The mass term  $\frac{1}{2} m_\phi^2 \chi^2$  is the standard quadratic



term of a spin-0 boson; the *value*  $m_\phi = 5.60 \text{ eV}$  is the algebraic ansatz of Eq. (13), not a quantity derived from Eq. (15). No renormalisable portal coupling to the Standard Model is included:  $\chi$  interacts only gravitationally. Two features of Eq. (15) are essential below. First,  $\chi$  is a *boson*, not a fermion — which excludes the active–sterile mixing production channel. Second, the action contains *no* free continuous parameter other than  $\xi$ , whose role we now constrain by the production mechanism.

## 5.4 Production mechanism: the Dodelson–Widrow route is excluded

A relic of mass  $m_\phi = 5.60 \text{ eV}$  contributing  $\Omega_{\phi\text{-DM}} h^2 \simeq 0.074$  can in principle be populated by two mechanisms. We tested both against the zero-parameter requirement.

**(a) Dodelson–Widrow (non-resonant mixing) — tested, excluded.** The Dodelson–Widrow mechanism [?] produces the relic through active–sterile oscillations, which requires  $\chi$  to be a fermion and introduces a mixing angle  $\theta$ . Matching the observed abundance with the Boyarsky–Ruchayskiy–Shaposhnikov relation [?] fixes

$$\sin^2(2\theta)_{\text{DW}} \simeq 4.78 \times 10^{-5}. \quad (16)$$

For SSEE to retain zero free parameters,  $\sin^2(2\theta)_{\text{DW}}$  would have to admit a closed algebraic form in  $\varphi, \pi$ . A scan of  $\sim 80$  monomial combinations of  $\varphi, \pi, \Omega, \mathcal{M}_{\text{SSEE}}$  (script `ssee_paperB_DW.py`) returns a best candidate  $\varphi^{-21} = 4.09 \times 10^{-5}$ , which differs from Eq. (16) by 14.6% — beyond the 10% threshold adopted for a “clean” SSEE identity. **This is a negative result:** the DW mixing angle does *not* reduce to the SSEE algebra. Were  $\phi$ -DM a DW sterile neutrino,  $\sin^2(2\theta)$  would be an irreducible free parameter, violating the zero-parameter principle. The DW route is therefore excluded on internal-consistency grounds, independently of any observational bound.

**(b) Gravitational particle production — adopted.** The remaining channel consistent with the action (15) is gravitational particle production [?]: the time-dependent metric during the inflation-to-reheating transition populates the  $\chi$  field directly, with *no* mixing angle. The relic abundance depends only on  $m_\phi$  and the inflationary potential  $V(\phi)$  — and in SSEE both are already fixed ( $m_\phi$  by Eq. (13); the  $\alpha$ -attractor potential with  $\alpha = \varphi^4/3$  by Paper 1). This route introduces no free parameter and selects  $\chi$  as a scalar, consistently with Eq. (15).

**Honest scope statement.** We have established *which* mechanism is compatible with zero parameters (gravitational production) and *which* is excluded (Dodelson–Widrow). We have *not* computed the full gravitational-production integral that would yield  $\Omega_{\phi\text{-DM}} h^2$  from  $m_\phi$  and  $V(\phi)$  *ab initio*; an order-of-magnitude estimate is consistent with the required abundance, but the complete calculation (including the  $\xi$ -dependence) is deferred to a

dedicated study (Paper B). The free-streaming scale  $k_{\text{fs}}$  used in Section 8.3 is therefore retained as an order-of-magnitude estimate, as already caveated there.

## 6 Observational Verification

We solve the linear growth equation for two limits and construct  $f\sigma_8(z)$  for each.

### 6.1 Growth equations

The growth-factor ODE in the conformal Newton gauge is  $\ddot{D} + \mathcal{H}\dot{D} - \frac{3}{2}\mathcal{H}^2\Omega_m(a)D = 0$ , where dots denote derivatives with respect to  $\ln a$ .

For all observationally relevant scales ( $k < k_{\text{fs}}$ ), both dark matter sectors contribute to the growth of structure. The relevant regime is:

**Two-sector regime** ( $k < k_{\text{fs}}$ ):  $\Omega_{\text{m,growth}} = 0.320$  (both CDM and  $\phi$ -DM cluster). The Hubble function is  $E^2(a) = 0.320 a^{-3} + 0.840 f_{\text{DE}}(a)$  with the Chevallier–Polarski–Linder [?] form  $f_{\text{DE}}(a) = a^{-3(1+w_0+w_a)} e^{-3w_a(1-a)}$ . This regime covers both RSD surveys ( $k \sim 0.01\text{--}0.1 h/\text{Mpc}$ ) and the  $\sigma_8$  scale ( $k = 0.125 h/\text{Mpc}$ ).

We integrate the growth ODE with  $\Omega_m = 0.320$  and compare to the  $\Lambda$ CDM solution ( $\Omega_m = 0.315$ ) to obtain the two-sector growth factor:

$$G_{2s} = \frac{D_1^{\text{SSEE}}(\Omega_m = 0.320)}{D_1^{\Lambda\text{CDM}}(\Omega_m = 0.315)} = 0.979, \quad (17)$$

giving  $\sigma_8^{\text{eff}} = \sigma_{8,\Lambda\text{CDM}} \times G_{2s} = 0.811 \times 0.979 = 0.794$ . The growth rate is  $f = d \ln D / d \ln a$  and the redshift-space distortion observable [?] is  $f\sigma_8(z) = f(z) \sigma_8^{\text{eff}} D(z)/D(0)$ .

### 6.2 $f\sigma_8$ results

Table 1 compares the two-sector model against six RSD surveys and against SSEE-V3.6 Paper 5 (single sector,  $\Omega_m = 0.160$ ) and  $\Lambda$ CDM.

The two-sector model reduces the mean  $f\sigma_8$  tension from  $2.67\sigma$  to  $0.76\sigma$ , comparable to and consistent with  $\Lambda$ CDM ( $0.73\sigma$ ) and well within the range of observational systematics in RSD measurements. This result follows self-consistently from the two-sector growth factor  $G_{2s} = 0.979$  with  $\sigma_8^{\text{eff}} = 0.794$ ; no warm dark matter transfer function is required.

### 6.3 $\sigma_8$ and $S_8$

The two-sector growth factor from eq. (17) gives directly:

$$\sigma_8^{\text{eff}} = \sigma_{8,\Lambda\text{CDM}} \times G_{2s} = 0.811 \times 0.979 = 0.794. \quad (18)$$

The derived structure amplitude is  $S_8 \equiv \sigma_8^{\text{eff}} \sqrt{\Omega_{\text{m,total}}/0.3} = 0.794 \times \sqrt{0.320/0.3} = 0.820$ . The KiDS-1000 value  $S_8 = 0.766 \pm 0.020$  [?] gives a tension of  $2.62\sigma$  [=  $(0.820 -$

Table 1:  $f\sigma_8$  predictions vs. observations. Survey values from ?, ?, ?, ? — the same canonical six-point set used in Paper 5.

| Survey                            | $z$   | Obs   | SSEE 2-sector                  |               | SSEE P5                        |               | $\Lambda$ CDM                  |
|-----------------------------------|-------|-------|--------------------------------|---------------|--------------------------------|---------------|--------------------------------|
|                                   |       |       | Pred                           | $\sigma$      | Pred                           | $\sigma$      | $\sigma$                       |
| 6dFGRS                            | 0.067 | 0.423 | 0.402                          | $+0.39\sigma$ | 0.266                          | $+2.85\sigma$ | $-0.38\sigma$                  |
| SDSS MGS                          | 0.150 | 0.490 | 0.417                          | $+0.51\sigma$ | 0.284                          | $+1.42\sigma$ | $+0.21\sigma$                  |
| BOSS DR12                         | 0.380 | 0.497 | 0.443                          | $+1.20\sigma$ | 0.325                          | $+3.81\sigma$ | $+0.46\sigma$                  |
| BOSS DR12                         | 0.510 | 0.458 | 0.449                          | $+0.24\sigma$ | 0.343                          | $+3.03\sigma$ | $-0.43\sigma$                  |
| BOSS DR12                         | 0.610 | 0.436 | 0.449                          | $-0.38\sigma$ | 0.353                          | $+2.44\sigma$ | $-0.97\sigma$                  |
| eBOSS DR16                        | 1.480 | 0.462 | 0.379                          | $+1.84\sigma$ | 0.350                          | $+2.50\sigma$ | $+1.90\sigma$                  |
| <b>Mean <math> \sigma </math></b> |       |       | <b><math>0.76\sigma</math></b> |               | <b><math>2.67\sigma</math></b> |               | <b><math>0.73\sigma</math></b> |

$0.766)/\sqrt{0.020^2 + 0.005^2}]$ , better than  $\Lambda$ CDM ( $3.27\sigma$  vs KiDS). The DES Y3 value  $S_8 = 0.776 \pm 0.017$  [?] gives a tension of  $2.5\sigma [= (0.820 - 0.776)/\sqrt{0.017^2 + 0.005^2}]$ .

**Non-linear and WDM transfer-function robustness.** The  $S_8$  estimate above uses only the two-sector growth factor  $G_{2s}$ , which is conservative: it treats the  $\phi$ -DM as fully clustering on scales  $k < k_{fs}$  and ignores the smooth Viel+2005 WDM transfer function. Including  $T_{WDM}(k) = [1 + (\alpha k)^{2\mu}]^{-5/\mu}$  ( $\mu = 1.12$ , Viel+2005) with the CLASS-calibrated scale  $\alpha_{WDM} = 1.656 h/\text{Mpc}$  and the mixing formula  $P_{\text{mix}} = P_{\text{lin}} [(1 - f_\phi) + f_\phi T_{WDM}]^2$  yields  $\sigma_8^{\text{eff}} = 0.728$  (CAMB; ?) or  $0.737$  (CLASS; ?), hence  $S_8^{\text{WDM}} = 0.752\text{--}0.761$  ( $-0.72\sigma$  to  $-0.25\sigma$  vs KiDS). A Halofit [?] non-linear correction (CAMB Halofit) adds a  $+7.5\%$  boost to  $\sigma_8^{\text{eff}}$ , reduced from the  $\Lambda$ CDM value of  $+11.3\%$  because the WDM suppresses small-scale power at  $k > k_{fs}$ . The resulting non-linear estimate is  $S_8^{\text{nl}} = 0.808$  ( $+2.1\sigma$  KiDS). Noting that KiDS-1000 constrains the *linear*  $\sigma_8$  (HMcode non-linear corrections are folded into the WL analysis pipeline), the primary comparison is with the linear result. The SSEE two-sector model thus brackets the KiDS measurement:  $S_8 \in [0.752, 0.820]$ , encompassing the KiDS value  $0.766 \pm 0.020$ .

**HMcode-2020 baryonic feedback correction.** Applying the CLASS HMcode-2020 baryonic feedback suppression to the WDM baseline ( $S_8^{\text{WDM}} = 0.761$ , CLASS), we obtain a baryonic suppression factor  $B_{\text{eff}} = 0.9447$  (integrated over the CMB lensing  $k$ -range  $0.03\text{--}2.0 h/\text{Mpc}$  with the HMcode `hmcode_version = 2020_baryonic_feedback` parameter), yielding:

$$S_8^{\text{bar}} = S_8^{\text{WDM}} \times B_{\text{eff}}^{1/2} = 0.761 \times 0.9721 = 0.758. \quad (19)$$

This result is  $0.06\sigma$  from DES Y3 ( $S_8 = 0.776 \pm 0.017$ ) and  $0.40\sigma$  from KiDS-1000 ( $S_8 = 0.766 \pm 0.020$ ).

*Calibration caveat.* HMcode-2020 [?] was calibrated against  $\Lambda$ CDM N-body simulations with  $w = -1$  and  $\sigma_8 \approx 0.8$ . Applied to SSEE ( $w_0 = -0.840$ ,  $w_a = -0.670$ ), the suppression factor  $B_{\text{eff}}$  carries an estimated systematic uncertainty of  $\lesssim 5\%$ , propagating to  $\Delta S_8^{\text{bar}} \approx \pm 0.02$ . The nominal result  $S_8^{\text{bar}} = 0.758$  should therefore be interpreted as a

central estimate within the range  $S_8^{\text{bar}} \in [0.74, 0.78]$  pending dedicated dark-energy N-body calibration. The Level 2 step — running dedicated N-body simulations (BAHAMAS, ?; IllustrisTNG, ?; modified for SSEE dark energy) — requires  $\sim 5000$ – $20000$  CPU-hours and is deferred to a future HPC allocation.

## 6.4 Conserved background sectors

The extension operates only at the perturbation level. The background Friedmann equation remains identical to SSEE-V3.6 Papers 1–5 (Table 2), so all background tests pass without modification.

Table 2: Complete observational summary. Results from Papers 2–5 are conserved. New results (this paper) shown in bold.

| Observable                                                            | Result                                                                                  | Paper    |
|-----------------------------------------------------------------------|-----------------------------------------------------------------------------------------|----------|
| $(w_0, w_a)$ vs DESI DR2                                              | $0.05\sigma$                                                                            | 1,2      |
| Cluster $\chi_r^2$                                                    | $0.122$ (7 clusters)                                                                    | 2        |
| CMB $\Delta\text{BIC}$ (plik_lite)                                    | $-31.3$ (SSEE favoured)                                                                 | 3        |
| Algebraic $H_0$                                                       | $67.96 = 3(\varphi + \pi)^2$ (1.1 $\sigma$ Planck)                                      | 4        |
| $n_s$                                                                 | $0.96556 = 1 - \varphi^{-7}$ (0.16 $\sigma$ Planck)                                     | 4        |
| $c_{s,\text{eff}}^2$ (IS)                                             | 0 (exact) — all modes stable                                                            | 5        |
| $S_8$ vs KiDS                                                         | $1.96\sigma$ (vs $3.27\sigma$ $\Lambda\text{CDM}$ )                                     | 5        |
| $f\sigma_8$ <b>mean tension</b>                                       | $0.76\sigma$ ( <b>was</b> $2.67\sigma$ )                                                | <b>6</b> |
| $\sigma_8^{\text{eff}}$ ( $G_{2s} = 0.979$ )                          | $0.794$ ; $S_8 = 0.820$ (2.6 $\sigma$ <b>KiDS</b> )                                     | <b>6</b> |
| $S_8^{\text{WDM}}$ ( <b>CLASS</b> , $\alpha = 1.656$ )                | $0.761$ ( $-0.25\sigma$ <b>KiDS</b> ; $0.09\sigma$ <b>DES</b> )                         | <b>6</b> |
| $S_8^{\text{bar}}$ ( <b>HMcode-2020</b> , $B_{\text{eff}} = 0.9447$ ) | $0.758 \pm 0.02^{\text{sys}}$ (0.06 $\sigma$ <b>DES</b> ; <b>range</b> $[0.74, 0.78]$ ) | <b>6</b> |
| $m_\phi$ (algebraic)                                                  | $5.60 \text{ eV}$ ( <b>zero parameters</b> )                                            | <b>6</b> |
| <b>MCMC</b> $\chi_r^2$ ; $\Delta\text{BIC}$                           | $0.497$ ; $-12.1$ ( <b>SSEE favoured under this test</b> )                              | <b>6</b> |

## 6.5 Figures

Figure 1 shows the  $f\sigma_8$  comparison across the full redshift range and the tension summary for the three models.




Figure 1: **Left:**  $f\sigma_8(z)$  predictions for the two-sector SSEE-V3.6 model (blue solid, mean  $0.76\sigma$ ), SSEE-V3.6 Paper 5 base (red dashed,  $2.67\sigma$ ), and  $\Lambda$ CDM (green dotted,  $0.73\sigma$ ), compared to six RSD surveys (black points). **Right:** Mean  $|f\sigma_8|$  tension bar chart for the three models. Model parameters:  $G_{2s} = 0.979$ ,  $\sigma_8^{\text{eff}} = 0.794$ ,  $m_\phi = 5.60$  eV.

## 7 Bayesian MCMC Validation

To complement the algebraic derivation, we perform an `emcee` [?] MCMC analysis with  $N_w = 100$  walkers and  $N_{\text{steps}} = 25000$  steps (burn-in 6000), sampling three parameters:  $\theta = (\Omega_{\varphi\text{DM}}, k_{\text{fs}}, \sigma_{8,\text{in}})$ . The background ( $H_0, w_0, w_a, \Omega_{m,\text{dyn}}$ ) is held fixed at the algebraic values. Data: the six canonical  $f\sigma_8$  surveys of Paper 5, KiDS-1000  $\sigma_8 = 0.737 \pm 0.020$ , and DES Y3  $S_8 = 0.776 \pm 0.017$ .

### 7.1 Posterior results

Table 3: MCMC posterior summary vs. algebraic predictions. Uncertainties are 68% credible intervals.  $N_{\text{eff}}$  computed from the `emcee` autocorrelation time.

| Parameter                                | Posterior                    | Algebraic     | Tension          | Prior                         |
|------------------------------------------|------------------------------|---------------|------------------|-------------------------------|
| $\Omega_{\varphi\text{DM}}$              | $0.1614^{+0.0106}_{-0.0107}$ | 0.1599        | $0.1\sigma$      | $\mathcal{N}(0.160, 0.015^2)$ |
| $k_{\text{fs}}$ [h/Mpc]                  | $0.491^{+0.089}_{-0.088}$    | 0.493         | $0.0\sigma$      | $\mathcal{U}(0.1, 1.5)$       |
| $\sigma_{8,\text{in}}$                   | $0.8102^{+0.0058}_{-0.0058}$ | Planck: 0.811 | $0.2\sigma$      | $\mathcal{N}(0.811, 0.006^2)$ |
| $\sigma_{8,\text{eff}}$ (derived)        | $0.790^{+0.013}_{-0.013}$    | 0.794         | —                | —                             |
| $S_{8,\text{eff}}$ (derived)             | $0.817^{+0.013}_{-0.013}$    | 0.820         | $2.5\sigma$ KiDS | —                             |
| $\chi_r^2$ (8 data, 3 par.)              | 0.497                        | —             | —                | —                             |
| $\Delta\text{BIC}$ (SSEE– $\Lambda$ CDM) | −12.1                        | —             | SSEE favoured    | —                             |
| Acceptance fraction                      | 0.513                        | —             | —                | —                             |

The posteriors recover the algebraic predictions at  $\lesssim 0.1\sigma$ , confirming that the data are consistent with *all* SSEE parameters being determined purely from  $\varphi$  and  $\pi$ . The  $\Delta\text{BIC} = -12.1$  result (“decisive” on the Jeffreys scale) favours SSEE over  $\Lambda$ CDM despite

having  $k_{\text{SSEE}} = 1$  free sector parameter vs  $k_{\Lambda\text{CDM}} = 0$ , because the lower  $\chi_r^2 = 0.497$  more than compensates the BIC penalty of  $\ln(N_{\text{data}}) \approx 2.1$ .

Figure 2 shows the MCMC corner plot and the posterior  $f\sigma_8(z)$  confidence bands.




Figure 2: **Left:** MCMC posterior corner plot for  $(\Omega_{\phi\text{DM}}, k_{\text{fs}}, \sigma_{8,\text{in}})$ . Vertical dashed lines mark the algebraic predictions; all three lie within  $0.1\sigma$  of the posterior medians. **Right:**  $f\sigma_8(z)$  posterior median (blue solid) with 68% and 95% credible bands (shaded), compared to six RSD surveys (black points) and  $\Lambda\text{CDM}$  (green dotted).  $\chi_r^2 = 0.497$ .

## 8 Discussion

### 8.1 Status of the $S_8$ vs. $f\sigma_8$ split

Paper 5 identified an  $S_8$ – $f\sigma_8$  split: SSEE-V3.6 predicts the amplitude of fluctuations ( $S_8$ ) better than  $\Lambda\text{CDM}$  but predicted the growth rate worse. The two-sector extension resolves the rate tension while adjusting  $\sigma_8$  to the KiDS value:

|                         | Paper 5      | Paper 6 (2-sector)         | $\Lambda\text{CDM}$ |
|-------------------------|--------------|----------------------------|---------------------|
| $\sigma_8^{\text{eff}}$ | 0.702        | 0.794 ( $G_{2s} = 0.979$ ) | 0.811               |
| $S_8$ vs KiDS           | $1.96\sigma$ | $2.62\sigma$               | $3.27\sigma$        |
| $f\sigma_8$ mean        | $2.67\sigma$ | $0.76\sigma$               | $0.73\sigma$        |
| New parameters          | 0            | 0                          | —                   |

At the linear-growth phenomenology level, the two-sector model reduces the mean  $f\sigma_8$  tension to  $0.76\sigma$ , which is comparable to the  $\Lambda\text{CDM}$  reference value of  $0.73\sigma$  under the same survey set and error model. The  $S_8$  tension of  $2.62\sigma$  vs KiDS remains, but improves over  $\Lambda\text{CDM}$  ( $3.27\sigma$ ). The mechanism is purely the two-sector growth factor  $G_{2s} = 0.979$ ; no warm dark matter transfer function is invoked.

## 8.2 KiDS–DES internal tension and its interpretation in SSEE-V3.6

The apparent discrepancy between our  $S_8 = 0.820$  and the KiDS-1000 measurement  $S_8 = 0.766 \pm 0.020$  [?] and the DES Y3 measurement  $S_8 = 0.776 \pm 0.017$  [?] requires careful decomposition.

**Structure of the  $S_8$  statistic.** The quantity  $S_8 \equiv \sigma_8 \sqrt{\Omega_m/0.3}$  is constructed to be approximately orthogonal to the weak-lensing amplitude degeneracy direction. The normalisation  $\Omega_m = 0.3$  is conventional and implicitly assumes the inferred  $\Omega_m$  is close to 0.3. In the SSEE-V3.6 framework the relevant matter density has two distinct values:  $\Omega_{m,\text{dyn}} = 0.160$  (dynamical dark matter sector) and  $\Omega_{m,\text{CMB}} = 0.3199$  (total including the  $\phi$ -DM clustering contribution active at  $k < k_{\text{fs}}$ ). Growth-rate observables probe  $\Omega_{m,\text{dyn}}$ , while the CMB acoustic geometry is sensitive to  $\Omega_{m,\text{CMB}}$ .

**Why KiDS and DES disagree internally.** KiDS-1000 [?] and DES Y3 [?] themselves differ by  $\Delta S_8 \approx 0.4\sigma$ , which is sub-dominant but non-negligible. This residual is driven by: (i) different angular scale cuts ( $\ell_{\text{max}}$ ); (ii) differing baryonic feedback priors; and (iii) partially overlapping shear calibration schemes. These internal systematics shift  $S_8$  at the  $\sim 1\%$  level — comparable to the statistical error — making a direct sub-percent comparison to SSEE-V3.6 premature.

**The correct SSEE comparison.** In the two-sector model the total matter clustering budget uses  $\Omega_m^{\text{tot}} = \Omega_{\text{CDM}} + \Omega_{\phi\text{DM}} = 0.3199$  when  $k < k_{\text{fs}}$ . With this total, the SSEE-V3.6  $S_8$  statistic evaluates to  $S_{8,\text{SSEE}} = \sigma_8^{\text{eff}} \sqrt{0.3199/0.3} = 0.820$ , and the tension with KiDS ( $2.62\sigma$ ) and DES ( $2.5\sigma$ ) are comparable. Neither survey is singled out as anomalous with respect to SSEE-V3.6; both sit  $\sim 2.5\sigma$  below our prediction, consistent with the unresolved  $S_8$  tension that persists even in  $\Lambda\text{CDM}$ .

The improvement over single-sector SSEE-V3.6 (Paper 5:  $S_8 = 0.725$ ,  $1.96\sigma$  KiDS) is driven by the  $\phi$ -DM suppression of  $\sigma_8^{\text{eff}}$  from 0.702 to 0.794, which reduces the KiDS tension from positive to the same sign as  $\Lambda\text{CDM}$ . A joint DES+KiDS likelihood marginalised over  $(\sigma_8, \Omega_m)$  simultaneously — as will be provided by Euclid Wide Survey weak lensing — will provide a definitive test.

## 8.3 Compatibility with Lyman- $\alpha$ and WDM constraints

A potential objection is that  $m_\phi = 5.60$  eV lies far below the standard thermal-WDM Lyman- $\alpha$  bound  $m_{\text{WDM}} \gtrsim 3.3\text{--}3.5$  keV [???]. We address this in three steps.

**Step 1:  $\phi$ -DM is not thermal WDM.** Thermal WDM bounds assume (a) 100% of the dark matter is a single thermally produced species, and (b) the momentum distribution is a Fermi-Dirac or Bose-Einstein function at a single temperature. Neither condition holds here. The  $\phi$ -DM sector accounts for only half of the total matter budget ( $\Omega_{\phi\text{-DM}}/\Omega_m^{\text{tot}} \simeq 0.50$ ).

The remaining sector is cold CDM ( $\Omega_{\text{CDM}} = 0.160$ ), which provides full gravitational clustering at all scales  $k < k_{\text{fs}}$  and is *not* suppressed. For mixed dark matter models with a WDM fraction  $f_{\text{WDM}}$ , the Lyman- $\alpha$  constraint on the WDM component is substantially relaxed (see e.g. ??); for  $f_{\text{WDM}} \approx 0.5$ , the effective bound weakens to  $m_{\text{WDM}}^{\text{eff}} \gtrsim 0.1\text{--}0.5\text{ keV}$ .

**Step 2: The observable is  $k_{\text{fs}}$ , not  $m_\phi$  — and the production mechanism determines the mapping.** The quantity constrained by Lyman- $\alpha$  forest measurements is the matter power spectrum suppression wavenumber  $k_{\text{fs}}$ , not the particle mass directly. For a given mass  $m_\phi$ , the inferred  $k_{\text{fs}}$  depends critically on the production mechanism.

For a *thermally* produced relic at  $m_\phi = 5.60\text{ eV}$  ( $= 5.60 \times 10^{-3}\text{ keV}$ ), the ? formula gives

$$k_{\text{fs}}^{\text{thermal}} = 23.7 \left( \frac{m_\phi}{1\text{ keV}} \right)^{1.11} \left( \frac{\Omega_{\phi\text{-DM}} h^2}{0.12} \right)^{0.15} \approx 0.070\text{ h/Mpc}, \quad \lambda_{\text{fs}}^{\text{thermal}} \approx 90\text{ h}^{-1}\text{ Mpc}. \quad (20)$$

At this scale, the  $\phi$ -DM sector would behave essentially as cold dark matter on all observationally relevant scales ( $k < 0.5\text{ h/Mpc}$ ), and the  $\sigma_8$  tension would not be resolved.

For a *non-resonantly produced* (Dodelson-Widrow-like) component, the free-streaming scale is set not by the thermal velocity but by the typical production momentum; the approximate relation is [?]:

$$k_{\text{fs}}^{\text{DW}} = 7.68 \left( \frac{m_\phi}{1\text{ keV}} \right)^{0.5} \left( \frac{\Omega_{\phi\text{-DM}} h^2}{0.1} \right)^{0.5} \approx 0.487\text{ h/Mpc}, \quad \lambda_{\text{fs}}^{\text{DW}} \approx 12.9\text{ h}^{-1}\text{ Mpc}. \quad (21)$$

The ratio  $k_{\text{fs}}^{\text{DW}}/k_{\text{fs}}^{\text{thermal}} \approx 7.0$  reflects the fact that non-resonant production gives a colder phase-space distribution per unit mass than thermal equilibrium. The SSEE two-sector value  $k_{\text{fs}} = 0.493\text{ h/Mpc}$  is consistent with the DW estimate (Eq. 21) to within 1.2%, well within the calibration uncertainty of both formulas.

**Key point:** the choice of production mechanism is not arbitrary. The algebraic mass  $m_\phi = 5.60\text{ eV}$  (Section 5), combined with the thermal formula, gives  $k_{\text{fs}}^{\text{thermal}} \approx 0.070$  — too small to suppress  $\sigma_8$  at  $k_\sigma \approx 0.125\text{ h/Mpc}$ . The same mass with the DW formula gives  $k_{\text{fs}}^{\text{DW}} \approx 0.487$ , which lies exactly in the window required to resolve the  $f\sigma_8$  tension. This non-trivial coincidence constitutes the primary predictive claim of the two-sector construction: a non-thermal production mechanism is required by the algebraic mass, and the required mechanism predicts a Lyman- $\alpha$  signature within current observational reach.

**Validity caveat.** The DW formula was calibrated for non-resonantly produced sterile neutrinos at  $m_s \sim \mathcal{O}(\text{keV})$ ; applying it at  $m_\phi = 5.60\text{ eV}$  constitutes a three-order-of-magnitude extrapolation in mass. In the absence of a full Boltzmann hierarchy for the  $\phi$ -DM phase-space distribution,  $k_{\text{fs}} = 0.493\text{ h/Mpc}$  should be treated as an *order-of-magnitude estimate* to be refined by future IS-EFT Boltzmann calculations. A direct Lyman- $\alpha$  test should be formulated in terms of the predicted suppression scale  $k_{\text{fs}}$ , not the thermal-WDM mass bound.



**Step 3: What remains observable.** At scales  $k > k_{\text{fs}} = 0.493 h/\text{Mpc}$  (relevant for the Lyman- $\alpha$  forest at  $z \sim 2-4$ ), the CDM sector still contributes ( $\Omega_{\text{CDM}}/\Omega_m^{\text{tot}} \approx 50\%$ ) of the power. Expanding the mixing formula  $P_{\text{mix}}/P_{\Lambda\text{CDM}} = [(1 - f_\phi) + f_\phi T_{\text{WDM}}]^2$  gives

$$\frac{\Delta P(k)}{P_{\Lambda\text{CDM}}} = -2f_\phi(1 - f_\phi)(1 - T_{\text{WDM}}) - f_\phi^2(1 - T_{\text{WDM}})^2, \quad (22)$$

with  $f_\phi \equiv f_{\phi\text{-DM}} \approx 0.5$ . In two asymptotic regimes:

$$k \ll k_{\text{fs}} : \quad T_{\text{WDM}} \rightarrow 1, \quad \frac{\Delta P}{P} \rightarrow 0, \quad (23)$$

$$k \gg k_{\text{fs}} : \quad T_{\text{WDM}} \rightarrow 0, \quad \frac{\Delta P}{P} \rightarrow -f_\phi(2 - f_\phi) \simeq -0.75. \quad (24)$$

At  $k = k_{\text{fs}}$  the CLASS-calibrated value  $T_{\text{WDM}}(k_{\text{fs}}) \approx 0.19$  gives  $\Delta P/P \approx -0.65$ , i.e. 65% suppression. The dominant contribution is the linear term  $-2f_\phi(1 - f_\phi)(1 - T_{\text{WDM}})$ ; the quadratic  $-f_\phi^2(1 - T_{\text{WDM}})^2$  adds only  $\approx 10\%$  of that. A dedicated Lyman- $\alpha$  likelihood analysis (planned with the IS-EFT framework) is required to verify that this suppression profile is consistent with observed forest data. Such analysis constitutes a *falsification test*: if the forest shows no suppression at  $k \approx 0.5 h/\text{Mpc}$ , the two-sector model is excluded.

The observational predictions of this paper — the proposed  $f\sigma_8$  resolution and  $\sigma_8^{\text{eff}} = 0.794$  — depend only on the two-sector growth factor  $G_{2s} = 0.979$  at scales  $k < k_{\text{fs}}$ , and are robust to the Lyman- $\alpha$  issue above.

## 8.4 The $H(z)$ tension

Paper 5 reported  $H(z)$  tension ( $\chi_r^2 = 1.861$  on cosmic chronometers) as a structural consequence of  $\Omega_{\text{m,dyn}} = 0.160$ : the background expansion is driven by the dynamical sector alone, giving a faster expansion at intermediate redshifts. The two-sector extension does not alter the background Friedmann equation and therefore does not affect  $H(z)$ . This tension is the remaining target for future work.

## 8.5 Physical origin of the mass relation

The relation  $m_\phi = \Sigma m_\nu \times H_0^{\text{alg}}$  connects two quantities derived independently in Paper 4 via the acoustic residual operator. We note:

1.  $\Sigma m_\nu = 0.0824 \text{ eV}$  is the sum of active neutrino masses [?] derived from the IS thermal residual. It arises from the coupling of the IS sector to the baryon-photon plasma.
2.  $H_0^{\text{alg}} = 67.96 \text{ km/s/Mpc}$  is the expansion rate set by the IS relaxation horizon.
3. The product  $\Sigma m_\nu \times H_0^{\text{alg}}$  in natural units has dimensions  $[\text{eV}] \times [\text{km/s/Mpc}]$ . The numerical value 5.60 in electron-volts arises when  $H_0^{\text{alg}}$  is taken as a dimensionless number — i.e., as  $H_0$  in units of  $\text{km s}^{-1} \text{ Mpc}^{-1}$ .

The physical interpretation is: the  $\phi$ -DM field acquires a characteristic energy scale set by the same IS relaxation operator that governs active neutrino production. This is structurally analogous to the seesaw mechanism [?], where a sterile-sector mass is related to the active neutrino mass through a common operator — but, as established in Section 5.4, the analogy is dimensional only:  $\phi$ -DM is *not* a sterile neutrino. The field content is fixed by the action (15):  $\phi$ -DM is a real scalar boson, populated by gravitational particle production rather than by active–sterile mixing. The remaining open item is the *ab initio* relic abundance integral, not the field’s identity.

## 8.6 Falsifiable predictions

### Falsifiable predictions — Paper 6

1.  $m_\phi = 5.60 \text{ eV}$ : the characteristic  $\phi$ -DM energy scale, derived from the acoustic residual operator. Because  $\phi$ -DM is a gravitationally produced scalar with no Standard-Model portal (Section 5.4), it is *not* accessible to neutrino-mass experiments; the mass enters observation only through the free-streaming scale  $k_{\text{fs}}(m_\phi)$  imprinted on the matter power spectrum. The falsifiable channel is therefore  $k_{\text{fs}} = 0.493 h/\text{Mpc}$ , probed by the Lyman- $\alpha$  forest and by DESI Y3 / Euclid clustering (2026–2028); see prediction 3.
2.  $\sigma_8^{\text{eff}} = 0.794$  and  $S_8 = 0.820$ : the two-sector growth factor gives a specific prediction for the amplitude of matter fluctuations. If future Euclid weak-lensing constraints confirm  $S_8 < 0.78$ , the two-sector model prediction is validated; if  $S_8 > 0.85$  is observed, the model is falsified.
3.  $f\sigma_8$  at  $0.76\sigma$ : the two-sector  $f\sigma_8$  prediction is testable with DESI Y5 RSD measurements (2026–2027). A confirmed  $f\sigma_8 > 0.47$  at  $z = 0.15$  would tension the model at  $> 1\sigma$ .
4. **Background conserved**:  $H_0 = 67.96$ ,  $(w_0, w_a) = (-0.840, -0.670)$ , and  $\Delta\text{BIC}_{\text{CMB}} = -31.3$  remain as in Papers 1–5. Any experiment detecting  $H_0 > 70$  or  $w_0 > -0.7$  falsifies SSEE-V3.6 at the background level.

## 8.7 Status of the zero-parameter claim

The SSEE-V3.6 philosophy requires that no number be *fitted* to data. We verify:

- $\Omega_{\phi\text{-DM}} = (\mathcal{M}_{\text{SSEE}} - 1) \Omega_{\text{m,dyn}}$ : derived from the MIRA algebraic identity. Not fitted.
- $m_\phi = \Sigma m_\nu \times H_0^{\text{alg}}$ : both factors derive from Paper 4 algebraic operators. Not fitted.
- $G_{2s} = 0.979$ : computed by integrating the growth ODE with  $\Omega_{\text{m,growth}} = 0.320$  (fixed by the two sectors above). Not fitted.
- $\sigma_8^{\text{eff}} = 0.794$ : derived from  $G_{2s}$  and the Planck reference  $\sigma_{8,\Lambda\text{CDM}} = 0.811$ . Not fitted.

The two-sector extension adds zero free parameters to the model. The remaining question is the derivation of the physical coupling and the free-streaming scale  $k_{\text{fs}}$  from first principles within the IS-EFT framework, which we leave for future work.

## 9 Conclusions

We have proposed a phenomenological two-sector resolution for the  $f\sigma_8$  growth-rate tension identified in Paper 5 of the SSEE-V3.6 series. The key results are:

1. **EFT scan:** kinetic braiding ( $\alpha_B$ ) is a no-go — it suppresses growth rather than enhancing it. Run-rate modification ( $\alpha_M = -0.08$ ) reduces the tension to  $1.57\sigma$  but requires tuning to the observational bound.
2. **Two-sector model:** the MIRA algebraic identity  $\mathcal{M}_{\text{SSEE}} \approx 2$  implies a second dark matter sector,  $\Omega_{\phi\text{-DM}} = (\mathcal{M}_{\text{SSEE}} - 1) \Omega_{\text{m,dyn}} = 0.160$ , with a free-streaming cutoff at  $k_{\text{fs}}$ .
3. **Algebraic mass:**  $m_\phi = \Sigma m_\nu^{\text{active}} \times H_0^{\text{alg}} = 0.0824 \times 67.96 = 5.60 \text{ eV}$ , derived from the acoustic residual operator and the algebraic Hubble constant of Paper 4. Deviation from the Dodelson-Widrow target (5.74 eV): 2.4%. Zero new free parameters.
4. **Verification:** two-sector growth factor  $G_{2s} = 0.979$  gives  $\sigma_8^{\text{eff}} = 0.794$  ( $S_8 = 0.820$ ,  $2.62\sigma$  KiDS, better than  $\Lambda\text{CDM}$   $3.27\sigma$ ) and  $f\sigma_8$  mean tension **0.76** $\sigma$  (from  $2.67\sigma$  for the single-sector SSEE-V3.6 baseline,  $\sigma_{8,\text{base}} = 0.702$ ; comparable to  $\Lambda\text{CDM}$   $0.73\sigma$ ). All background tests — CMB ( $\Delta\text{BIC} = -31.3$ ), BAO ( $\chi_r^2 = 0.01$ ), clusters ( $\chi_r^2 = 0.122$ ) — conserved.

The framework SSEE-V3.6-V3.6 now offers a unified linear-level description across six categories of observational tests (background expansion, CMB, BAO, galaxy clusters, weak lensing, and RSD growth rate). The growth tension is reduced phenomenologically via the two-sector growth factor alone, without invoking any warm dark matter transfer function. However, as explicitly discussed, the mass coupling  $m_\phi = \Sigma m_\nu H_0^{\text{alg}}$  remains a numerical ansatz pending derivation from a complete lagrangian, and the free-streaming scale  $k_{\text{fs}}$  requires verification against Lyman- $\alpha$  forest data. The remaining background tension ( $H(z)$ ,  $\chi_r^2 = 1.861$ ) is structural and traceable to  $\Omega_{\text{m,dyn}} = 0.160$ .

The **falsifiable predictions** of this paper are: (1) the  $\phi$ -dark matter free-streaming scale  $k_{\text{fs}} = 0.493 h/\text{Mpc}$ , set by the algebraic mass  $m_\phi = 5.60 \text{ eV}$  and probed by the Lyman- $\alpha$  forest and by DESI Y3 / Euclid galaxy clustering (2026–2028) — the relevant channel because  $\phi$ -DM is a gravitationally produced scalar with no Standard-Model portal (Section 5.4), and so leaves no neutrino-sector signature; and (2)  $\sigma_8^{\text{eff}} = 0.794$  ( $S_8 = 0.820$ ), testable by Euclid weak-lensing (2026–2028).

## Data Availability

All scripts, data files, and compiled PDFs are available at <https://github.com/mikealmeida1721/SSEE>. The exact commit for the original two-sector verification is 9d28620; the self-consistent OptB rewrite (removing WDM transfer function) is committed separately and documented in the repository.

## Author Contributions

Single-author work. M.E.A.V. derived all analytical results, wrote all numerical scripts, and wrote the manuscript.

## Acknowledgements

The author thanks the anonymous referees for constructive comments that improved the manuscript.