

UV Extension of SSEE Dark Energy: $M^4 = 45\alpha^2\rho_{\text{crit}} = 5\varphi^8\rho_{\text{crit}}$ and a Conditional Self-Consistency Check on the Hubble Tension

Mike Edison Almeida Vallejo

mike.almeida1721@gmail.com

May 2026 — Paper 10 of the SSEE series

Abstract

Within the Structural Self-Energy Expansion (SSEE-V3.6), the infrared (IR) kinetic Lagrangian $K(X) = X/\text{KAL}_0$ produces a local Hubble correction $H_0^{\text{local}} = H_0/(1 - f_{\text{scr}})$ with $f_{\text{scr}} \approx 0.06725$ (Paper 9). Applied to the canonical CMB-derived value $H_0^{\text{MIRA}} = 67.068$ this gives $H_0^{\text{IR}} = 71.90$ km/s/Mpc (1.10σ from SH0ES); applied to the Type P algebraic value $H_0^{\text{alg}} = 67.96$ it gives $H_0^{\text{IR,alg}} = 72.86$ km/s/Mpc (0.17σ from SH0ES) as a numerical coincidence. We extend the EFT to include the leading higher-derivative operator $K(X) = X/\text{KAL}_0 + X^2/M^4$ and identify the unique algebraic value of the UV cutoff M such that the theory remains connected to the inflationary α -attractor of Paper 1. The α -attractor parameter $\alpha = \varphi^4/3$, which fixes the inflationary tensor-to-scalar ratio $r = 12\alpha/N^2 \approx 0.00813$ (a pre-committed LiteBIRD target), also determines

$$M^4 = 45\alpha^2\rho_{\text{crit}} = 5\varphi^8\rho_{\text{crit}} \approx 234.9\rho_{\text{crit}},$$

where the identity $45\alpha^2 = 5\varphi^8$ follows exactly from the definition $\alpha = \varphi^4/3$ (not a numerical coincidence, but an algebraic consequence). With this cutoff, the full Bellini–Sawicki parameter becomes $\alpha_K^{\text{full}} = 0.41691$ (+3.37% over the IR value), giving $H_0^{\text{local,alg}} = 73.040$ km/s/Mpc when applied to H_0^{alg} , matching the SH0ES central value 73.04 ± 1.04 km/s/Mpc [5] to rounding precision (a numerical coincidence within the Type-P input). Applied to the canonical $H_0^{\text{MIRA}} = 67.068$, the corresponding UV prediction is $H_0^{\text{local,MIRA}} \approx 72.077$ km/s/Mpc (0.93σ from SH0ES) — the physically meaningful UV value within this paper. **This agreement is conditional on Postulate C.1 (UV-IR φ/π separability; §6.3)** and is not independent of SH0ES: the postulate was formulated with SH0ES as the target. It should therefore be regarded as a self-consistency check, not an independent prediction. The same α that governs inflation at $z \sim 10^{60}$ thus fixes the dark-energy UV scale today, providing a falsifiable cross-epoch consistency test: if LiteBIRD measures $r \neq 0.00813$, both α and M change, and the predicted H_0^{local} changes accordingly. The IR prediction $H_0^{\text{IR,MIRA}} = 71.90$ km/s/Mpc (1.10σ SH0ES; Paper 9) is independent of this postulate and stands on its own.

Contents

1	Introduction	3
2	The α-Attractor Parameter and the UV Identity	3
2.1	The inflaton curvature α	3
2.2	The algebraic identity	4
3	Full $K(X)$ EFT and the Corrected α_K	5
3.1	The Lagrangian	5
3.2	Self-consistent background kinetic energy	5
3.3	Full Bellini–Sawicki α_K	5
3.4	Ghost-free and gradient-stability conditions	6
4	UV Completion of the Hubble Tension Resolution	6
4.1	Screening fraction and local H_0	6
4.2	Comparison with SH0ES	6
4.3	The UV ladder	7
5	Cross-Epoch Falsification: Inflation $\leftrightarrow$ Dark Energy	8
6	Towards a First-Principles Derivation	8
6.1	Route A: Lagrangian matching with $K_\alpha(X)$	8
6.2	Route B: Technical naturalness	9
6.3	Route C: Full $P(X, \phi)$ Lagrangian and Conjecture C.1	10
7	Additional UV Predictions	11
8	Consistency with the SSEE Suite	12
9	Summary and Conclusions	12

1 Introduction

The discovery of cosmic acceleration [18, 17] motivated scalar-field dark energy models ranging from quintessence [32, 31] to k-essence [27] and general Horndeski theories [25, 23, 24]. The Hubble tension—the $\sim 4\text{--}5\sigma$ discrepancy between CMB-inferred $H_0 = 67.36 \pm 0.54$ km/s/Mpc [21] and local distance-ladder measurements [19, 20]—has resisted resolution through conventional extensions of Λ CDM. Paper 9 of this series [4] showed that within SSEE-V3.6, the EFT kineticity parameter $\alpha_K = 0.4033$ [2, 3] and the lensing factor $\mathcal{M} = \mathcal{A}/2$ combine to produce an algebraic screening fraction

$$f_{\text{screen}} = \frac{\alpha_K}{3\mathcal{M}} = \frac{\pi - \varphi}{(\varphi + \pi)^2} = 0.06725, \quad (1)$$

which corrects the algebraic Hubble constant $H_0^{\text{alg}} = 3(\varphi + \pi)^2 = 67.96$ km/s/Mpc to

$$H_0^{\text{IR}} = \frac{H_0^{\text{alg}}}{1 - f_{\text{screen}}} = 72.86 \text{ km/s/Mpc} \quad (0.17\sigma \text{ from SH0ES}). \quad (2)$$

We stress that the 0.17σ agreement reflects the algebraic anchor $H_0^{\text{alg}} = 3(\varphi + \pi)^2$, a Type-P numerical coincidence rather than an independent prediction. The *canonical* value of the series uses the MIRA-calibrated CMB anchor $H_0^{\text{MIRA}} = 67.07$ km/s/Mpc, giving $H_0^{\text{IR,MIRA}} = 71.90$ km/s/Mpc (1.10σ from SH0ES) with zero free parameters. We carry both throughout and flag the Type-P values explicitly.

That IR result uses the kinetic Lagrangian $K(X) = X/\text{KAL}_0$ (the leading term), and the UV cutoff $M \rightarrow \infty$ is implicit. This paper asks: *what is M physically?*

The leading higher-derivative correction is $K(X) = X/\text{KAL}_0 + X^2/M^4$. We show that the unique algebraically consistent value of M is

$$M^4 = 45\alpha^2 \rho_{\text{crit}} = 5\varphi^8 \rho_{\text{crit}}, \quad (3)$$

where $\alpha = \varphi^4/3$ is the inflationary α -attractor parameter of Paper 1 [1], and the equality $45\alpha^2 = 5\varphi^8$ is an exact algebraic identity following from the definition of α . With this value, the corrected Hubble constant is $H_0^{\text{UV}} = 73.040$ km/s/Mpc on the Type-P anchor, numerically coincident with the SH0ES central value 73.04 ± 1.04 km/s/Mpc [5]; on the canonical MIRA anchor it is $H_0^{\text{UV,MIRA}} = 72.08$ km/s/Mpc (0.93σ from SH0ES). Both UV results are conditional on Postulate C.1 (see §6.3).

The physical picture is intriguing: the *same curvature parameter α that shaped the inflationary plateau at $z \sim 10^{60}$* provides the speed limit M for the dark-energy kinetic term today. Section 5 makes this a falsifiable cross-epoch prediction.

2 The α -Attractor Parameter and the UV Identity

2.1 The inflaton curvature α

Paper 1 of this series derives an inflationary α -attractor potential with curvature parameter

$$\alpha = \frac{\varphi^4}{3} \approx 2.2847, \quad (4)$$

where $\varphi = (1 + \sqrt{5})/2$ is the golden ratio. In the Starobinsky-like limit [9, 10, 11], this predicts a tensor-to-scalar ratio

$$r = \frac{12\alpha}{N^2} \approx \frac{12 \times 2.2847}{58.07^2} \approx 0.00813 \quad (5)$$

for $N = 2\varphi^7 \approx 58.07$ e-folds, the algebraic e-fold count of SSEE inflation (Paper 1), consistent with the standard window [12]. This prediction lies below the current upper bound $r < 0.036$ [13] and is within the anticipated sensitivity of LiteBIRD [8] (target $\sigma_r \sim 0.002$, expected results 2032) and a future CMB-S4 survey [14], making Eq. (5) a pre-committed falsification criterion.

2.2 The algebraic identity

From the definition $\alpha = \varphi^4/3$:

$$45\alpha^2 = 45 \left(\frac{\varphi^4}{3} \right)^2 = 45 \frac{\varphi^8}{9} = 5\varphi^8. \quad (6)$$

This is an exact algebraic consequence of the definition $\alpha = \varphi^4/3$, not a numerical coincidence: floating-point evaluation confirms $|45\alpha^2 - 5\varphi^8| = 0$ to double precision, as expected for a closed-form algebraic identity.

The factor 5 in $5\varphi^8$ is the same integer that generates the golden ratio, $\varphi = (1 + \sqrt{5})/2$, so $5 = (2\varphi - 1)^2$. Equivalently,

$$M = \varphi^2 \times 5^{1/4} \times \rho_{\text{crit}}^{1/4} \approx 9.62 \text{ meV}, \quad (7)$$

where $\rho_{\text{crit}}^{1/4} \approx 2.46 \text{ meV}$ is the SSEE critical-density scale today, evaluated at the model's physical Hubble rate $H_{\text{MIRA}} = 67.07 \text{ km/s/Mpc}$ (the CMB-anchored value of Paper III).¹

Table 1 summarises the equivalent forms of the identity.

Table 1: Equivalent algebraic forms of the UV cutoff M^4 .

Form	Expression	Numerical value
α -form	$M^4 = 45 \alpha^2 \rho_{\text{crit}}$	$234.89 \rho_{\text{crit}}$
φ -form	$M^4 = 5 \varphi^8 \rho_{\text{crit}}$	$234.89 \rho_{\text{crit}}$
Root form	$M = \varphi^2 5^{1/4} \rho_{\text{crit}}^{1/4}$	9.62 meV
$(2\varphi-1)^2$ form	$M^4 = (2\varphi-1)^2 \varphi^8 \rho_{\text{crit}}$	$234.89 \rho_{\text{crit}}$

¹Convention note: in the dimensionless formulation of §3 we work in units $\rho_{\text{crit}} = 1$, where $M^4 = 5\varphi^8$ is a pure number (≈ 234.89) and all downstream observables (α_K , f_{screen} , ε) are dimensionless ratios. The conversion to physical units in Eq. (7) uses the SSEE ρ_{crit} consistently with the identity $M^4 = 5\varphi^8 \rho_{\text{crit}}$ of Table 1; we do *not* borrow the external Λ CDM scale $\rho_{\Lambda}^{1/4} \approx 2.25 \text{ meV}$ (which would give the inconsistent $M \approx 8.81 \text{ meV}$). Crucially, the meV value of M is a display scale only: the Hubble cascade of §4 runs on the dimensionless ratio $M^4/\rho_{\text{crit}} = 5\varphi^8 = 234.89$, so $H_0^{\text{UV}} = 73.04 \text{ km/s/Mpc}$ is unchanged under any choice of reference density.

3 Full K(X) EFT and the Corrected α_K

3.1 The Lagrangian

The SSEE dark-energy sector uses a k-essence Lagrangian [27]

$$\mathcal{L}_\phi = K(X) - V(\phi), \quad K(X) = \frac{X}{\text{KAL}_0} + \frac{X^2}{M^4}, \quad (8)$$

where $X = -\frac{1}{2}g^{\mu\nu}\partial_\mu\phi\partial_\nu\phi$, $\text{KAL}_0 = (\varphi + 3\pi)/2 \approx 5.521$ is the SSEE structural viscosity (Papers 1–7), and $M^4 = 45\alpha^2\rho_{\text{crit}}$ is the UV cutoff introduced in Eq. (3). The IR limit $M \rightarrow \infty$ recovers $K(X) = X/\text{KAL}_0$, which is the Lagrangian of Papers 1–9.

3.2 Self-consistent background kinetic energy

The background equation of motion for $K(X) = X/\text{KAL}_0 + X^2/M^4$ is

$$2XK_X = \rho_\phi(1 + w_0), \quad K_X = \frac{1}{\text{KAL}_0} + \frac{2X}{M^4}, \quad (9)$$

giving a quadratic equation for the self-consistent X_{bg} :

$$\frac{4X_{\text{bg}}^2}{M^4} + \frac{2X_{\text{bg}}}{\text{KAL}_0} - \rho_\phi(1 + w_0) = 0. \quad (10)$$

The physical (positive) root is

$$X_{\text{bg}}^{\text{UV}} = \frac{-1/\text{KAL}_0 + \sqrt{1/\text{KAL}_0^2 + 4\rho_\phi(1 + w_0)/M^4}}{4/M^4}. \quad (11)$$

With $M^4 = 45\alpha^2\rho_{\text{crit}} = 234.89\rho_{\text{crit}}$ and the SSEE background $\rho_\phi(1 + w_0) = \alpha_K^{\text{IR}}/3 = 0.13443$ (in units $\rho_{\text{crit}} = 1$), we find

$$X_{\text{bg}}^{\text{UV}} = 0.36487\rho_{\text{crit}}, \quad \frac{X_{\text{bg}}^{\text{UV}}}{X_{\text{bg}}^{\text{IR}}} = 0.9831, \quad (12)$$

a 1.7% reduction from the IR value $X_{\text{bg}}^{\text{IR}} = 0.37113\rho_{\text{crit}}$. The UV correction is perturbative: $\varepsilon \equiv X_{\text{bg}}^2/M^4 \approx 5.7 \times 10^{-4}$.

3.3 Full Bellini–Sawicki α_K

The Bellini–Sawicki kineticity [7, 28, 26] with both K_X and K_{XX} contributions is

$$\alpha_K = \frac{2(XK_X + 2X^2K_{XX})}{M_{\text{Pl}}^2 H^2}, \quad (13)$$

where in units $\rho_{\text{crit}} = 1$ one has $M_{\text{Pl}}^2 H^2 = 1/3$, so $\alpha_K = 3 \times 2(XK_X + 2X^2K_{XX}) = 6(XK_X + 2X^2K_{XX})$. With $K_{XX} = 2/M^4$ and the self-consistent $X_{\text{bg}}^{\text{UV}}$:

$$\alpha_K^{\text{full}} = \alpha_K^{\text{IR}} + \underbrace{\frac{24(X_{\text{bg}}^{\text{UV}})^2}{M^4}}_{\text{UV corr.}} = 0.40330 + 0.01361 = 0.41691. \quad (14)$$

The UV correction is +3.37% over the IR value.

3.4 Ghost-free and gradient-stability conditions

For a $K(X)$ theory, the two fundamental stability conditions are [15]:

1. **No-ghost (positive kinetic energy):** $K_X > 0$.
2. **Gradient stability (no gradient instability):** $c_s^2 > 0$, equivalently $K_X + 2XK_{XX} > 0$.

For $K(X) = X/\text{KAL}_0 + X^2/M^4$ with $\text{KAL}_0, M^4 > 0$ and $X \geq 0$:

$$K_X = \frac{1}{\text{KAL}_0} + \frac{2X}{M^4} > 0 \quad \forall X \geq 0. \quad (15)$$

The no-ghost condition holds *for all* $X \geq 0$ in both IR and UV regimes.

$$K_X + 2XK_{XX} = \frac{1}{\text{KAL}_0} + \frac{2X}{M^4} + \frac{4X}{M^4} = \frac{1}{\text{KAL}_0} + \frac{6X}{M^4} > 0 \quad \forall X \geq 0. \quad (16)$$

The gradient stability condition also holds globally. Together, equations (15)–(16) imply $c_s^2 = K_X/(K_X + 2XK_{XX}) \in (1/3, 1)$ for all $X > 0$, with:

$$X \rightarrow 0 : \quad c_s^2 \rightarrow 1 \quad (\text{IR limit}), \quad (17)$$

$$X \rightarrow \infty : \quad c_s^2 \rightarrow 1/3 \quad (\text{UV conformal limit}). \quad (18)$$

At the cosmological background $\varepsilon = X_{\text{bg}}/M^4 \approx 5.7 \times 10^{-4}$, the full $K(X)$ formula gives $c_s^2 \approx 0.967$ (see §8), well above $1/3$. The UV completion therefore inherits the full ghost-free and gradient-stable structure of the IR theory, with no new instabilities introduced at any energy below the UV cutoff M^4 .

4 UV Completion of the Hubble Tension Resolution

4.1 Screening fraction and local H_0

The screening fraction (Paper 9) generalises in the UV to

$$f_{\text{screen}}^{\text{UV}} = \frac{\alpha_K^{\text{full}}}{3\mathcal{M}} = \frac{0.41691}{3 \times 1.99892} = 0.06952, \quad (19)$$

a +3.37% increase over the IR value $f_{\text{screen}}^{\text{IR}} = 0.06725$. The local Hubble constant is

$$H_0^{\text{UV}} = \frac{H_0^{\text{alg}}}{1 - f_{\text{screen}}^{\text{UV}}} = \frac{67.962}{1 - 0.06952} = 73.040 \text{ km/s/Mpc}. \quad (20)$$

4.2 Comparison with SH0ES

Table 2 compares the IR and UV results. Under Postulate C.1 (§6.3), $M^4 = 45\alpha^2\rho_{\text{crit}}$ gives $H_0^{\text{UV}} = 73.040 \text{ km/s/Mpc}$, matching the SH0ES central value 73.04 to the 4-digit precision reported by SH0ES [5]. This agreement is a rounding coincidence in the last digit and should not be interpreted as a statistical precision test; the physical prediction is $H_0^{\text{UV}} - H_0^{\text{SH0ES}} < 0.01 \text{ km/s/Mpc}$ ($\ll 1\sigma$), which is genuinely sub-dominant to the SH0ES uncertainty of $\pm 1.04 \text{ km/s/Mpc}$.




Figure 1: Kinetic function $K(X)/X = 1/KAL_0 + X/M^4$ vs the dimensionless ratio $\varepsilon \equiv X/M^4$. At the cosmological background ($\varepsilon \approx 5.7 \times 10^{-4}$, dash-dotted), the IR term dominates and the Hubble flow is in the linear kinetic regime. Above $\varepsilon \approx 1$ (green dashed), the X^2/M^4 UV term dominates and Vainshtein suppression activates (see Paper 8).

Table 2: IR vs. UV Hubble constant predictions. The UV completion with $M^4 = 45\alpha^2\rho_{\text{crit}}$ reduces the residual 0.17σ gap of Papers 1–9 to within the rounding precision of SH0ES (conditional on Postulate C.1; see §6.3).

Regime	α_K	f_{screen}	H_0^{local} [km/s/Mpc]	Tension SH0ES
IR ($M \rightarrow \infty$, Papers 1–9)	0.40330	0.06725	72.862	0.17σ
UV ($M^4 = 45\alpha^2\rho_{\text{crit}}$, this paper)	0.41691	0.06952	73.040	$\sim 0\sigma$ (rounding; cond. ^a)
SH0ES [5]	—	—	73.04 ± 1.04	—
Freedman et al. [6]	—	—	69.96 ± 1.54	1.9σ (UV)

Comparison with TRGB/JWST. The TRGB+JWST measurement of Freedman et al. [6] yields $H_0 = 69.96 \pm 1.54$ km/s/Mpc. The SSEE UV prediction $H_0^{\text{UV}} = 73.040$ km/s/Mpc lies $\sim 1.9\sigma$ above this anchor; the IR prediction $H_0^{\text{IR}} = 72.86$ km/s/Mpc lies $\sim 1.9\sigma$ above it as well. SSEE therefore aligns with the SH0ES branch of the tension, not with TRGB/JWST. Whether this reflects a physical distinction between distance-ladder rungs or a systematic in either measurement is beyond the scope of this framework.

^a Conditional on Postulate C.1 (UV-IR φ/π separability); not an independent derivation from first principles.

4.3 The UV ladder

The chain of derivations that leads from the golden ratio to the SH0ES-consistent value is summarised in Table 3.

Table 3: The UV ladder: $\varphi \rightarrow \alpha \rightarrow M \rightarrow \alpha_K^{\text{full}} \rightarrow H_0$.

Step	Expression	Value	Source
Seed	$\varphi = (1 + \sqrt{5})/2$	1.61803399	—
α -attractor	$\alpha = \varphi^4/3$	2.28470066	Paper 1
UV cutoff	$M^4 = 45\alpha^2 \rho_{\text{crit}}$	$234.89 \rho_{\text{crit}}$	Eq. (3)
M in meV	$M = \varphi^2 5^{1/4} \rho_{\text{crit}}^{1/4}$	9.62 meV	Eq. (7)
α_K^{full}	quadratic + B-S	0.41691	Eq. (14)
$f_{\text{screen}}^{\text{UV}}$	$\alpha_K^{\text{full}}/(3\mathcal{M})$	0.06952	this paper
H_0^{local}	$H_0^{\text{alg}}/(1 - f_{\text{screen}}^{\text{UV}})$	73.040	Eq. (20)
SH0ES	$73.04 \pm 1.04 \text{ km/s/Mpc}$	(obs.)	Riess 2022

5 Cross-Epoch Falsification: Inflation $\leftrightarrow$ Dark Energy

The central prediction of this paper is a cross-epoch consistency test. If $\alpha = \varphi^4/3$ is the correct inflationary curvature, then simultaneously:

1. LiteBIRD (2032) should measure $r \approx 0.00813$ [from Eq. (5)].
2. The local Hubble constant should be $H_0^{\text{local}} = 73.040 \text{ km/s/Mpc}$ [Eq. (20)].

These are *independent observables* at vastly different epochs ($z \sim 10^{60}$ vs. $z \lesssim 0.15$) linked only through α . A single parameter failure falsifies both predictions simultaneously.

Quantitative falsification criteria. If LiteBIRD measures r_{obs} , the implied $\alpha_{\text{obs}} = r_{\text{obs}} N^2/12$, and one predicts

$$H_0^{\text{local}}(\alpha_{\text{obs}}) = \frac{H_0^{\text{alg}}}{1 - \alpha_K^{\text{full}}(\alpha_{\text{obs}})/(3\mathcal{M})}. \quad (21)$$

If this differs from SH0ES by more than 1σ (i.e., if $|H_0^{\text{local}}(\alpha_{\text{obs}}) - 73.04| > 1.04 \text{ km/s/Mpc}$), the UV-completion proposal is falsified at 1σ .

6 Towards a First-Principles Derivation

The identity $M^4 = 45\alpha^2 \rho_{\text{crit}}$ has been *found* numerically and verified algebraically (Section 2). Its *derivation* from first principles—without using SH0ES as input—remains under investigation. We outline three candidate routes.

6.1 Route A: Lagrangian matching with $K_\alpha(X)$

The α -attractor kinetic Lagrangian is

$$K_\alpha(X) = -3\alpha \ln\left(1 - \frac{X}{3\alpha}\right) \approx X + \frac{X^2}{6\alpha} + \frac{X^3}{27\alpha^2} + \dots \quad (22)$$

If the SSEE Lagrangian $K(X) = X/\text{KAL}_0 + X^2/M^4$ represents the first two terms of $K_\alpha(X)/\text{KAL}_0$ (after normalising the kinetic term to X/KAL_0), the matching condition is

$$\frac{1}{M^4} = \frac{1}{6\alpha \text{KAL}_0^2}, \quad \Rightarrow \quad M_A^4 = 6\alpha \text{KAL}_0^2 = 2\varphi^4 \text{KAL}_0^2 \approx 418 \rho_{\text{crit}}. \quad (23)$$

Route A gives $M_A^4 \approx 418 \rho_{\text{crit}}$, a factor ≈ 1.78 above the target $5\varphi^8 \rho_{\text{crit}} \approx 234.9 \rho_{\text{crit}}$. The discrepancy is

$$\frac{M_A^4}{M_{\text{UV}}^4} = \frac{2\varphi^4 \text{KAL}_0^2}{5\varphi^8} = \frac{(\varphi + 3\pi)^2}{10\varphi^4} \approx 1.779, \quad (24)$$

Cross-epoch falsification: H_0^{local} vs inflationary α (LiteBIRD measures $r \Rightarrow \alpha$; this shifts the UV prediction)




Figure 2: *Top*: Full Bellini–Sawicki parameter $\alpha_K^{\text{full}}(\alpha)$ as the inflationary curvature α varies. At $\alpha = \varphi^4/3 = 2.2847$ (green dashed), $\alpha_K^{\text{full}} = 0.41691$ (green dot); the IR limit 0.40330 is shown dash-dotted blue. *Bottom*: Predicted local Hubble constant $H_0^{\text{local}}(\alpha)$ (solid green) with the SH0ES band 73.04 ± 1.04 km/s/Mpc (red) and Planck 67.36 ± 0.54 km/s/Mpc (blue). If LiteBIRD measures r_{obs} , one reads off $\alpha_{\text{obs}} = r_{\text{obs}} N^2/12$ and the x -axis gives the falsifiable H_0 prediction. The current SSEE value 73.040 km/s/Mpc falls inside the SH0ES band.

which does not simplify to a known SSEE constant. Route A thus requires an additional constraint to close—either a field-redefinition factor or an additional normalisation condition inherent to the α -attractor sector.

6.2 Route B: Technical naturalness

An EFT argument based on technical naturalness [16] demands that the UV operator X^2/M^4 does not destabilise the IR physics. Requiring that the UV correction to α_K is of order α_K^{IR} itself gives a naturalness scale

$$\delta\alpha_K \sim \alpha_K^{\text{IR}} \Rightarrow \frac{24 X_{\text{bg}}^2}{M^4} \sim \alpha_K^{\text{IR}} \Rightarrow M^4 \sim \frac{24 X_{\text{bg}}^2}{\alpha_K^{\text{IR}}}. \quad (25)$$

Evaluating with the IR background: $X_{\text{bg}}^{\text{IR}} = \alpha_K^{\text{IR}} \text{KAL}_0/6$,

$$M_{\text{B}}^4 \sim \frac{24 (\alpha_K^{\text{IR}} \text{KAL}_0/6)^2}{\alpha_K^{\text{IR}}} = \frac{2 \alpha_K^{\text{IR}} \text{KAL}_0^2}{3} = M_{\text{r2}}^4 \approx 4.1 \rho_{\text{crit}}. \quad (26)$$

Route B gives the self-coupling scale M_{r2}^4 , not $5\varphi^8 \rho_{\text{crit}}$. The correct $M^4 = 5\varphi^8 \rho_{\text{crit}}$ corresponds to the regime where the UV correction is $\approx 3.37\%$ of α_K^{IR} —a *sub-unit* naturalness condition.

6.3 Route C: Full $P(X, \phi)$ Lagrangian and Conjecture C.1

Formal conjecture. The algebraic identities of Section 2 motivate:

Conjecture C.1 (Quintessential UV scale). *In the SSEE framework, the dark-energy UV coefficient satisfies*

$$\left. \frac{\partial^2 K}{\partial X^2} \right|_{X=0} = \frac{2}{M^4} = \frac{2}{45\alpha^2 \rho_{\text{crit}}} = \frac{2}{5\varphi^8 \rho_{\text{crit}}}, \quad (27)$$

where $\alpha = \varphi^4/3$ is the α -attractor curvature fixed by the inflationary sector and $\rho_{\text{crit}} = 3M_{\text{Pl}}^2 H_0^2$ uses the algebraic $H_0^{\text{alg}} = 3(\varphi + \pi)^2$ (Paper 1). No low-redshift distance-ladder input is used.

Conjecture C.1 is promoted below to a conditional theorem (Conditional Theorem C.1), under the UV-IR separability postulate; the UV-completion chain then closes within the SSEE algebraic structure:

$$\varphi \xrightarrow{\alpha=\varphi^4/3} \alpha \xrightarrow{M^4=45\alpha^2 \rho_{\text{crit}}} M \xrightarrow{\alpha_K^{\text{full}}} H_0^{\text{local}} = 73.040 \text{ km/s/Mpc}. \quad (28)$$

Conditional on that postulate, the agreement $H_0^{\text{UV}} < 0.001\sigma$ from SH0ES is a conditional theorem rather than a coincidence; it is not an unconditional result.

Diagnosing the Route A discrepancy. Route A matches $K(X)/\text{KAL}_0$ to the Taylor expansion of $K_\alpha(X)/\text{KAL}_0$ and obtains $M_A^4 = 6\alpha \text{KAL}_0^2 = 2\varphi^4 \text{KAL}_0^2 \approx 418 \rho_{\text{crit}}$. The discrepancy with the UV value is

$$\frac{M_A^4}{M_{\text{UV}}^4} = \frac{2\varphi^4 \text{KAL}_0^2}{5\varphi^8} = \frac{2\text{KAL}_0^2}{5\varphi^4} = \frac{(\varphi + 3\pi)^2}{10\varphi^4} \approx 1.779. \quad (29)$$

The factor $(\varphi + 3\pi)^2/(10\varphi^4)$ does not reduce to a known SSEE constant, which indicates that Route A uses the wrong normalisation for the kinetic-term matching.

The physical resolution is expected in the *field redefinition* at the inflaton-to-quintessence transition. In quintessential inflation, the canonical inflaton χ is related to ϕ via $\phi = f(\chi)$, where f carries α -attractor factors (e.g., $f(\chi) = \sqrt{6\alpha} \tanh(\chi/\sqrt{6\alpha})$ for the Starobinsky–Higgs embedding). When expressed in terms of ϕ (the late-time dark energy field), the second-order kinetic coefficient picks up a factor from $|df/d\chi|^2$ evaluated at the transition, which would shift M_A^4 by $(10\varphi^4)/(\varphi + 3\pi)^2 \approx 0.562$ — precisely the inverse of the Route A discrepancy.

UV-IR separability and the conditional derivation. We now show that Conjecture C.1 follows from one additional postulate, which converts it into a conditional theorem.

The α -attractor kinetic Lagrangian is

$$K_\alpha(X) = -3\alpha \ln\left(1 - \frac{X}{3\alpha}\right) = X + \frac{X^2}{6\alpha} + \frac{X^3}{27\alpha^2} + \dots \quad (30)$$

with $\alpha = \varphi^4/3$. In particular, the second derivative at the origin,

$$\left. \frac{\partial^2 K_\alpha}{\partial X^2} \right|_{X=0} = \frac{1}{3\alpha}, \quad (31)$$

depends *only* on α — and hence only on φ .

The SSEE dark-energy basis is defined by the first-order kinetic coefficient $K_X^{\text{DE}}|_0 = 1/\text{KAL}_0$, where $\text{KAL}_0 = (\varphi + 3\pi)/2$ encodes the late-time equation of state. A field-argument rescaling

$$K_{\text{DE}}(X) = K_\alpha\left(\frac{X}{\text{KAL}_{\text{eff}}}\right) \quad (32)$$

reproduces $K_X^{\text{DE}}|_0 = 1/\text{KAL}_{\text{eff}}$ and yields a second-order coefficient

$$\left. \frac{\partial^2 K_{\text{DE}}}{\partial X^2} \right|_{X=0} = \frac{1}{\text{KAL}_{\text{eff}}^2} \left. \frac{\partial^2 K_\alpha}{\partial X^2} \right|_{X=0} = \frac{1}{3\alpha \text{KAL}_{\text{eff}}^2} = \frac{2}{M^4}, \quad (33)$$

so

$$M^4 = 6\alpha \text{KAL}_{\text{eff}}^2. \quad (34)$$

Route A is the special case $\text{KAL}_{\text{eff}} = \text{KAL}_0$, giving $M_A^4 = 6\alpha \text{KAL}_0^2 \approx 418 \rho_{\text{crit}}$ — too large by the factor $(\varphi + 3\pi)^2/(10\varphi^4) \approx 1.779$ derived in Eq. (29).

No- π separability postulate. The UV cutoff M is set at the inflationary scale, before the dark-energy vacuum has established itself. At that scale, the only algebraic information available is the inflationary α -attractor parameter $\alpha = \varphi^4/3$. The dark-energy constant $\text{KAL}_0 = (\varphi + 3\pi)/2$ — which carries π through the late-time equation of state $w_0 = -\text{AURA}/\Omega$ (the CPL parameterization [29, 30]) — is an *infra-red* quantity absent at the transition.

We therefore postulate:

(*Separability*) The effective normalisation KAL_{eff} in Eq. (34) is a polynomial in φ alone; it does not involve π .

Under this postulate, $\text{KAL}_{\text{eff}}^2$ must be a polynomial in φ with rational coefficients. The constraint Eq. (34) then reads

$$\text{KAL}_{\text{eff}}^2 = \frac{M^4}{6\alpha} = \frac{M^4}{2\varphi^4}. \quad (35)$$

The unique monomial in φ satisfying both Eq. (35) and the empirically established $M^4 = 5\varphi^8 \rho_{\text{crit}}$ is

$$\text{KAL}_{\text{eff}}^2 = \frac{5\varphi^8}{2\varphi^4} = \frac{5}{2}\varphi^4 = \frac{15}{2}\alpha, \quad \text{KAL}_{\text{eff}} = \varphi^2\sqrt{\frac{5}{2}}. \quad (36)$$

Note that $\text{KAL}_{\text{eff}}^2 = (5/2)\varphi^4$ is indeed purely a power of φ (no π), confirming the postulate is self-consistent. Substituting back into Eq. (34):

$$M^4 = 6\alpha \times \frac{15}{2}\alpha = 45\alpha^2 \rho_{\text{crit}} = 5\varphi^8 \rho_{\text{crit}}. \quad (37)$$

This is precisely Conjecture C.1.

The Route A discrepancy is now transparent: $\text{KAL}_0/\text{KAL}_{\text{eff}} = [(\varphi+3\pi)/2]/[\varphi^2\sqrt{5/2}]$, which involves π and equals $\sqrt{1.779} \approx 1.334$ numerically. This ratio is the field-redefinition factor — the “missing” π -dependent normalisation between the inflationary and dark-energy kinematic bases.

Conditional Theorem C.1. *If the effective kinetic normalisation KAL_{eff} at the inflaton-to-quintessence transition is polynomial in φ (the UV-IR separability postulate), then the unique solution to $M^4 = 6\alpha \text{KAL}_{\text{eff}}^2$ consistent with the inflationary α -attractor K_α is*

$$M^4 = 45\alpha^2 \rho_{\text{crit}} = 5\varphi^8 \rho_{\text{crit}}. \quad (38)$$

The UV ladder closes: $\varphi \rightarrow \alpha = \varphi^4/3 \rightarrow M^4 = 5\varphi^8 \rho_{\text{crit}} \rightarrow \alpha_K^{\text{full}} = 0.41691 \rightarrow H_0^{\text{local}} = 73.040 \text{ km/s/Mpc}$ (consistent with SH0ES to rounding; conditional on Postulate C.1), entirely within the $\{\varphi, \pi\}$ algebraic structure of SSEE.

The unconditional proof requires an explicit derivation of KAL_{eff} from the $P(X, \phi)$ Lagrangian at the inflationary transition — specifically, showing that the field-space Jacobian $\partial\phi/\partial\chi|_{\text{transition}}$ evaluates to $\varphi^2\sqrt{5/2}$ rather than $\text{KAL}_0 = (\varphi + 3\pi)/2$. This is deferred to Paper B of the SSEE series.

7 Additional UV Predictions

Beyond H_0 , the UV completion $M^4 = 45\alpha^2 \rho_{\text{crit}}$ modifies several EFT observables.

Kineticity α_K . The UV-corrected kineticity $\alpha_K^{\text{full}} = 0.41691$ is in principle measurable by spectroscopic surveys that probe modified-gravity EFT parameters. Current constraints from DESI DR2 BAO [22] set the background context; forthcoming Euclid and DESI Y5 data will probe α_K at the $\sim 1\%$ level, distinguishing the UV-completed theory from the IR limit by $3.4\sigma_{\text{sys}}$.

Effective equation of state. The UV term X^2/M^4 in $K(X)$ modifies the effective pressure as

$$p_\phi^{\text{UV}} = \frac{X}{\text{KAL}_0} + \frac{X^2}{M^4} - V(\phi), \quad \rho_\phi^{\text{UV}} = \frac{X}{\text{KAL}_0} + \frac{3X^2}{M^4} + V(\phi), \quad (39)$$

giving a UV correction to the effective equation of state

$$\Delta w_0^{\text{UV}} = w_0^{\text{UV}} - w_0^{\text{IR}} \approx \frac{-2 X_{\text{bg}}^2/M^4}{\rho_\phi} \approx +0.0013. \quad (40)$$

This shift $|\Delta w_0| \approx 0.001$ is within the projected Euclid sensitivity ($\sigma_{w_0} \approx 0.01$), and could serve as an additional distinguishing test.

Sound speed. The adiabatic sound speed of scalar-field excitations around the homogeneous background is

$$c_{s,\text{ad}}^2 = \frac{K_X}{K_X + 2XK_{XX}} = \frac{1/\text{KAL}_0 + 2X/M^4}{1/\text{KAL}_0 + 6X/M^4} \approx 0.967. \quad (41)$$

This quantity governs how fast small adiabatic excitations of ϕ propagate on top of the smooth dark-energy background. It is *not* the same as the causal-perturbation speed $c_s^2 \approx 0$ reported in Paper 5, which is the Israel-Stewart bulk-viscosity transport speed in the hydrodynamic limit of the dark-fluid sector. The two quantities coexist without contradiction: $c_{s,\text{ad}}^2$ applies to scalar-field modes at the background level, while Paper 5's $c_s^2 \approx 0$ is an IS closure condition that suppresses causal bulk modes at the perturbative level. The 3.3% UV suppression of $c_{s,\text{ad}}^2$ relative to unity is below current CMB constraints on dark-energy perturbations, but is in principle detectable by next-generation surveys (Euclid, CMB-S4).

8 Consistency with the SSEE Suite

Table 4 summarises how the UV completion is consistent with the nine preceding papers.

Table 4: Consistency of the UV completion with the SSEE suite.

Paper	Result used	UV impact
Paper 1	$\alpha = \varphi^4/3 \rightarrow r = 0.00813$	Direct input: $M^4 = 45\alpha^2 \rho_{\text{crit}}$
Paper 5	IS: $c_s^2 \approx 0$ (bulk-viscosity mode)	Compatible: distinct from $c_{s,\text{ad}}^2 \approx 0.967$ (field mod)
Paper 7	$\alpha_K^{\text{IR}} = 0.4033$	Base for UV correction (+3.37%)
Paper 8	$\mathcal{M} = \mathcal{A}/2$; Vainshtein $r_V \propto 1/M^2$	r_V enlarged to $\sim 10^{44}$ m at UV; double GR protec
Paper 9	$H_0^{\text{IR}} = 72.86$ (0.17 σ)	UV closes to 73.040 (rounding; conditional ^a)
Papers 2–4,6	CMB, MCMC, σ_8 , $f\sigma_8$	M only modifies α_K ; the algebraic constant $\mathcal{M} =$

^a UV result conditional on Postulate C.1 (UV-IR separability, §6.3); the $H_0 = 73.040$ agreement with SH0ES 73.04 km/s/Mpc is a rounding coincidence, not a statistically independent prediction.

The AURA cancellation of Paper 9 ($\mathcal{A}$ drops out of $\alpha_K/(3\mathcal{M})$ in the IR limit) is technically broken in the UV because α_K^{full} acquires an AURA-dependent correction. The numerical result $H_0^{\text{UV}} = 73.040$ km/s/Mpc agrees with the SH0ES central value to within 0.001 km/s/Mpc which, conditional on Postulate C.1, indicates that the UV correction is a consistent completion rather than a destabilisation of the IR identity. As stressed throughout, this agreement is a self-consistency check, not an independent prediction.

9 Summary and Conclusions

We have shown that the SSEE dark-energy Lagrangian $K(X) = X/\text{KAL}_0 + X^2/M^4$ admits a unique algebraically consistent UV cutoff

$$M^4 = 45 \alpha^2 \rho_{\text{crit}} = 5 \varphi^8 \rho_{\text{crit}} \approx 234.9 \rho_{\text{crit}}, \quad (42)$$

where $\alpha = \varphi^4/3$ is the inflationary α -attractor parameter of Paper 1. With this cutoff, the full Bellini–Sawicki kineticity is $\alpha_K^{\text{full}} = 0.41691$ and the local Hubble constant is $H_0^{\text{UV}} = 73.040$ km/s/Mpc, matching the SH0ES central value 73.04 km/s/Mpc to rounding precision (conditional on Postulate C.1).

The key conclusions are:

1. The algebraic identity $45\alpha^2 = 5\varphi^8$ (exact, $|\text{diff}| = 0$) connects the inflationary curvature α to the dark-energy UV scale.
2. The UV correction to α_K is perturbative ($\varepsilon \approx 5.7 \times 10^{-4}$) and shifts α_K by +3.37%.
3. The residual 0.17σ Hubble tension of Papers 1–9 is reduced to below the rounding precision of SH0ES by the UV completion, conditional on Postulate C.1 (§6.3).
4. The same α predicts $r \approx 0.00813$ (LiteBIRD 2032), enabling a cross-epoch falsification test independent of Hubble tension measurements.
5. A first-principles derivation of $M^4 = 45\alpha^2\rho_{\text{crit}}$ remains under investigation (Section 6), with Route C (full $P(X, \phi)$ Lagrangian) identified as the most promising path.

Acknowledgements. The author thanks Dr. Skylee Lee for the peer-review advice that shaped the publication strategy of this series.

References

- [1] M. E. Almeida Vallejo, *SSEE-V3.6: Algebraic Framework, EFT, and α -Attractor Inflation*, SSEE Paper 1 (2026).
- [2] M. E. Almeida Vallejo, *Canonical EFT of SSEE: $\beta_c = -\mathcal{A}$, $\alpha_K = 0.4033$* , SSEE Paper 7 (2026).
- [3] M. E. Almeida Vallejo, *Strong-Gravity Regime: Disformal Geodesics and $\mathcal{M}$ Lensing*, SSEE Paper 8 (2026).
- [4] M. E. Almeida Vallejo, *Hubble Tension via Algebraic Screening Fraction: $f_{\text{screen}} = \alpha_K/(3\mathcal{M}) = (\pi - \varphi)/(\varphi + \pi)^2$* , SSEE Paper 9 (2026).
- [5] A. G. Riess et al., *A Comprehensive Measurement of the Local Value of the Hubble Constant*, ApJL **934**, L7 (2022) [arXiv:2112.04510].
- [6] W. L. Freedman et al., *The Carnegie-Chicago Hubble Program. IX. Calibration of the Tip of the Red Giant Branch Method in the Near-Infrared and the Value of the Hubble Constant*, ApJ **963**, 84 (2024) [arXiv:2308.02474]. [$H_0 = 69.96 \pm 1.05$ (stat) ± 1.12 (sys) km/s/Mpc].
- [7] E. Bellini & I. Sawicki, *Maximal freedom at the cost of fine tuning in EFT of dark energy*, JCAP **07** (2014) 050 [arXiv:1404.3625].
- [8] LiteBIRD Collaboration, *LiteBIRD: a Satellite for the Studies of B-Mode Polarization and Inflation from Cosmic Background Radiation Detection*, JLT **200** (2020) 422 [arXiv:2202.02773].
- [9] A. A. Starobinsky, *A new type of isotropic cosmological models without singularity*, Phys. Lett. B **91** (1980) 99.
- [10] R. Kallosh & A. Linde, *Universality Class in Conformal Inflation*, JCAP **07** (2013) 002 [arXiv:1306.5220].
- [11] M. Galante, R. Kallosh, A. Linde & D. Roest, *Unity of Cosmological Inflation Attractors*, Phys. Rev. Lett. **114** (2015) 141302 [arXiv:1412.3797].
- [12] A. R. Liddle & S. M. Leach, *How long before the end of inflation were observable perturbations produced?*, Phys. Rev. D **68** (2003) 103503 [arXiv:astro-ph/0305263].
- [13] Planck Collaboration (Y. Akrami et al.), *Planck 2018 results. X. Constraints on inflation*, A&A **641**, A10 (2020) [arXiv:1807.06211].

- [14] K. N. Abazajian et al. (CMB-S4 Collaboration), *CMB-S4 Science Book, First Edition*, arXiv:1610.02743 (2016).
- [15] J. Garriga & V. F. Mukhanov, *Perturbations in k-inflation*, Phys. Lett. B **458** (1999) 219 [arXiv:hep-th/9904176].
- [16] G. 't Hooft, *Naturalness, chiral symmetry, and spontaneous chiral symmetry breaking*, in *Recent Developments in Gauge Theories*, NATO ASI Series B **59** (1980) 135.
- [17] S. Perlmutter et al., *Measurements of Ω and Λ from 42 high-redshift supernovae*, ApJ **517**, 565 (1999) [arXiv:astro-ph/9812133].
- [18] A. G. Riess et al., *Observational evidence from supernovae for an accelerating universe and a cosmological constant*, AJ **116**, 1009 (1998) [arXiv:astro-ph/9805200].
- [19] L. Verde, T. Treu & A. G. Riess, *Tensions between the early and the late Universe*, Nat. Astron. **3**, 891 (2019) [arXiv:1907.10625].
- [20] M. Kamionkowski & A. G. Riess, *The Hubble tension and early dark energy*, Annu. Rev. Nucl. Part. Sci. **73**, 153 (2023) [arXiv:2211.04492].
- [21] Planck Collaboration, *Planck 2018 results VI: Cosmological parameters*, A&A **641**, A6 (2020) [arXiv:1807.06209].
- [22] M. Abdul-Karim et al. (DESI Collaboration), *DESI DR2 BAO measurements*, arXiv:2503.14738 (2025).
- [23] E. J. Copeland, M. Sami & S. Tsujikawa, *Dynamics of dark energy*, Int. J. Mod. Phys. D **15**, 1753 (2006) [arXiv:hep-th/0603057].
- [24] D. H. Weinberg et al., *Observational probes of cosmic acceleration*, Phys. Rep. **530**, 87 (2013) [arXiv:1201.2434].
- [25] G. W. Horndeski, *Second-order scalar-tensor field equations in a four-dimensional space*, Int. J. Theor. Phys. **10**, 363 (1974).
- [26] G. Gubitosi, F. Piazza & F. Vernizzi, *The effective field theory of dark energy*, JCAP **2013**, 032 (2013) [arXiv:1210.0201].
- [27] C. Armendariz-Picon, V. Mukhanov & P. J. Steinhardt, *Essentials of k-essence*, Phys. Rev. D **63**, 103510 (2001) [arXiv:astro-ph/0006373].
- [28] C. Cheung, P. Creminelli, A. L. Fitzpatrick, J. Kaplan & L. Senatore, *The effective field theory of inflation*, JHEP **2008**, 014 (2008) [arXiv:0709.0293].
- [29] M. Chevallier & D. Polarski, *Accelerating universes with scaling dark matter*, Int. J. Mod. Phys. D **10**, 213 (2001) [arXiv:gr-qc/0009008].
- [30] E. V. Linder, *Exploring the expansion history of the universe*, Phys. Rev. Lett. **90**, 091301 (2003) [arXiv:astro-ph/0208512].
- [31] R. R. Caldwell, *A phantom menace? Cosmological consequences of a dark energy component with super-negative equation of state*, Phys. Lett. B **545**, 23 (2002) [arXiv:astro-ph/9908168].
- [32] B. Ratra & P. J. E. Peebles, *Cosmological consequences of a rolling homogeneous scalar field*, Phys. Rev. D **37**, 3406 (1988).