

SSEE-V3.6 and the Hubble Tension: An Algebraic Screening Fraction from First Principles

Mike Edison Almeida Vallejo

mike.almeida1721@gmail.com

May 2026 — Paper 9 of the SSEE series

Abstract

The Hubble tension between the CMB-inferred value $H_0^{\text{CMB}} = 67.4 \pm 0.5$ km/s/Mpc and the local distance-ladder measurement $H_0^{\text{SH0ES}} = 73.04 \pm 1.04$ km/s/Mpc remains one of the sharpest anomalies in modern cosmology. Within the Structural Self-Energy Expansion (SSEE-V3.6) framework, the algebraic constants φ (golden ratio) and π determine all parameters with zero free parameters. We derive algebraically that the dark-energy EFT of Paper 7 and the disformal geodesic of Paper 8 together imply a *local screening fraction*

$$f_{\text{screen}} = \frac{\alpha_K}{3\mathcal{M}} = \frac{\pi - \varphi}{(\varphi + \pi)^2} \approx 0.0673,$$

where $\alpha_K = 3\Omega_{\text{DE}}(1 + w_0)$ is the EFT kineticity derived from the background equation of state, and $\mathcal{M} = \beta_c/2 = (3\varphi + \pi)/4 \approx 1.9989$ is the effective optical coupling derived from the disformal null geodesic. The two structural constants $\mathcal{A} = 2\mathcal{M}$ cancel exactly in the ratio, leaving a pure function of φ and π . Applied as a local correction to the cosmological value $H_0^{\text{alg}} = 3(\varphi + \pi)^2$,

$$H_0^{\text{local}} = \frac{H_0^{\text{alg}}}{1 - f_{\text{screen}}} = \frac{67.96}{1 - 0.0673} \approx 72.86 \text{ km/s/Mpc},$$

which lies 0.17σ from SH0ES (Riess et al. 2022) with *zero free parameters*. The correction operates at late times ($z < 0.15$) through the scalar kinetic energy contribution to the local expansion rate, and does not modify the sound horizon r_d , the CMB peak positions, or the growth factor σ_8 derived in Papers 3 and 5.

Contents

1	Introduction	3
2	Background: EFT Kineticity and Disformal Coupling	3
2.1	The SSEE action (Paper 7)	3
2.2	The disformal coupling and MIRA (Paper 8)	4

3	Algebraic Derivation: $f_{\text{screen}} = \alpha_K/(3\mathcal{M})$	4
3.1	Step 1: Exact expression for α_K	5
3.2	Step 2: Exact expression for $3\mathcal{M}$	5
3.3	Step 3: The cancellation and the identity	5
3.4	Numerical verification	6
4	Physical Interpretation	6
4.1	Two-epoch structure of H_0 measurements	6
4.2	Local Hubble correction	7
4.3	Why the CMB is unaffected	7
5	Consistency with the SSEE Series	8
6	Pre-committed Falsifiable Prediction	8
7	Limitations and Future Work	9
8	Conclusions	9

1 Introduction

The Hubble tension [Verde et al., 2019, Riess et al., 2022, Kamionkowski & Riess, 2023] is the 4–6 σ discrepancy between measurements of the present expansion rate obtained by two independent methods:

- **Early-universe inference:** CMB anisotropies (Planck PR4, Planck Collaboration 2020) combined with acoustic oscillation physics yield $H_0 = 67.4 \pm 0.5$ km/s/Mpc under the assumption of Λ CDM.
- **Local distance-ladder:** Type Ia supernovae calibrated with Cepheids in the SH0ES programme [Riess et al., 2022] give $H_0 = 73.04 \pm 1.04$ km/s/Mpc, probing predominantly $z < 0.15$.

The SSEE-V3.6 framework [Almeida, Paper1,P,P,P] derives the cosmological expansion rate algebraically:

$$H_0^{\text{alg}} = 3(\varphi + \pi)^2 \approx 67.96 \text{ km/s/Mpc}, \quad (1)$$

which agrees with Planck at the 0.81 σ level (MCMC Paper 2, Almeida Paper2) but does not, at first sight, resolve the local tension.

In this paper we show that the same algebraic structure that produces H_0^{alg} also produces a local kinematic correction $f_{\text{screen}} \approx 0.0673$ from first principles, predicting $H_0^{\text{local}} = 72.86$ km/s/Mpc. The derivation combines two results already established in this series:

1. From Paper 7 [Almeida, Paper7]: the EFT kineticity $\alpha_K = 3 \Omega_{\text{DE}}(1 + w_0)$, which follows from the background Friedmann equations and the k -essence action $K(X) = X/\text{KAL}_0$.
2. From Paper 8 [Almeida, Paper8]: the effective optical coupling $\mathcal{M} = \beta_c/2 = (3\varphi + \pi)/4$, which emerges from the disformal null geodesic of dark matter.

Together these give the exact algebraic identity $f_{\text{screen}} = \alpha_K/(3\mathcal{M})$.

Organisation. Section 2 recalls the key results from Papers 7 and 8. Section 3 derives the algebraic identity in three steps. Section 4 interprets the result physically. Section 5 verifies consistency with the rest of the SSEE series. Section 6 states the pre-committed falsifiable prediction and its current observational status. Section 8 summarises.

2 Background: EFT Kineticity and Disformal Coupling

2.1 The SSEE action (Paper 7)

The fundamental action of SSEE-V3.6 is a k -essence scalar minimally coupled to gravity and conformally coupled to dark matter:

$$S = \int d^4x \sqrt{-g} \left[\frac{M_{\text{Pl}}^2}{2} R + K(X, \phi) + \mathcal{L}_m \right], \quad (2)$$

with kinetic kernel

$$K(X, \phi) = \frac{X}{\text{KAL}_0} + \frac{X^2}{M^4} - V_0 e^{-\lambda\phi/M_{\text{Pl}}}, \quad (3)$$

where $X = -\frac{1}{2}g^{\mu\nu}\partial_\mu\phi\partial_\nu\phi$ and

$$\text{KAL}_0 = \frac{\varphi + \pi}{2} + \pi \approx 5.521, \quad (4)$$

$$\beta_c \equiv \mathcal{A} = \frac{3\varphi + \pi}{2} \approx 3.998. \quad (5)$$

All parameters are algebraically fixed; no fitting is performed.

The dark-energy equation-of-state parameters derived in Paper 1 are

$$w_0 = -\frac{\mathcal{A}}{\varphi + \pi} \approx -0.840, \quad (6)$$

$$\Omega_{\text{DE}} = -w_0 = \frac{\mathcal{A}}{\varphi + \pi} \approx 0.840. \quad (7)$$

The EFT kineticity in the Bellini–Sawicki classification [Bellini & Sawicki, 2015] is defined as

$$\alpha_K \equiv \frac{2X K_X}{M_{\text{Pl}}^2 H^2}, \quad (8)$$

where $K_X = \partial K / \partial X$. Paper 7 derives $\alpha_K = 0.4033$ numerically from CLASS/EFTCAMB; in Section 3 we provide its exact algebraic expression.

2.2 The disformal coupling and MIRA (Paper 8)

Paper 8 extends the action to the strong-gravity regime by coupling dark matter to the disformal metric

$$\tilde{g}_{\mu\nu} = g_{\mu\nu} + \frac{2}{M^4} \partial_\mu\phi\partial_\nu\phi. \quad (9)$$

The disformal null geodesic for a photon passing through a dark-matter halo yields a total deflection angle

$$\hat{\alpha}_{\text{total}} \approx \beta_c \left(\frac{4GM}{c^2 b} \right), \quad (10)$$

where b is the impact parameter and $\beta_c = \mathcal{A}$. The effective Einstein ring radius is therefore amplified by

$$\frac{\theta_E^{\text{SSEE}}}{\theta_E^{\text{GR}}} \approx \sqrt{\beta_c} \approx \mathcal{M}, \quad \mathcal{M} \equiv \frac{\beta_c}{2} = \frac{3\varphi + \pi}{4} \approx 1.9989. \quad (11)$$

The constant $\mathcal{M}$ encodes the effective optical coupling of the disformal sector; it is not a fitted parameter but derived from $\beta_c = \mathcal{A}$. Paper 8 further derives $\mathcal{M} = \Omega_{\text{DE}}^{\text{CMB}} / \Omega_{\text{DE}}^{\text{dyn}}$, confirming its role as a structural ratio between the CMB-era and dynamical matter fractions.

3 Algebraic Derivation: $f_{\text{screen}} = \alpha_K / (3\mathcal{M})$

We derive the screening fraction in three steps using only established SSEE relations.

3.1 Step 1: Exact expression for α_K

The background Friedmann equations give

$$\rho_\phi = \frac{X}{\text{KAL}_0} + V = 3M_{\text{Pl}}^2 H^2 \Omega_{\text{DE}}, \quad (12)$$

$$p_\phi = \frac{X}{\text{KAL}_0} - V = w_0 \rho_\phi. \quad (13)$$

Subtracting: $V = \rho_\phi(1 - w_0)/2$; adding: $X/\text{KAL}_0 = \rho_\phi(1 + w_0)/2 = 3M_{\text{Pl}}^2 H^2 \Omega_{\text{DE}}(1 + w_0)/2$.

For the IR kinetic term $K(X) = X/\text{KAL}_0$ (EFT regime, $X \ll M^4/\text{KAL}_0$): $K_X = 1/\text{KAL}_0$. Substituting into equation (8):

$$\alpha_K = \frac{2X}{\text{KAL}_0 M_{\text{Pl}}^2 H^2} = \frac{2 \cdot \text{KAL}_0 \rho_\phi(1 + w_0)/2}{\text{KAL}_0 M_{\text{Pl}}^2 H^2} = \frac{\rho_\phi(1 + w_0)}{M_{\text{Pl}}^2 H^2} = 3 \Omega_{\text{DE}}(1 + w_0). \quad (14)$$

Using the SSEE algebraic values (6)–(7),

$$\Omega_{\text{DE}}(1 + w_0) = \frac{\mathcal{A}}{\varphi + \pi} \cdot \frac{\varphi + \pi - \mathcal{A}}{\varphi + \pi} = \frac{\mathcal{A}(\pi - \varphi)/2}{(\varphi + \pi)^2}, \quad (15)$$

where we used $(\varphi + \pi) - \mathcal{A} = (\pi - \varphi)/2$ (exact identity from the definitions (5) and $\varphi + \pi = \Omega$):

$$\Omega - \mathcal{A} = (\varphi + \pi) - \frac{3\varphi + \pi}{2} = \frac{2\varphi + 2\pi - 3\varphi - \pi}{2} = \frac{\pi - \varphi}{2}. \quad (16)$$

Therefore,

$$\boxed{\alpha_K = \frac{3\mathcal{A}(\pi - \varphi)}{2(\varphi + \pi)^2}}. \quad (17)$$

Numerically: $\alpha_K = 3 \times 3.998 \times 1.524/(2 \times 22.65) = 0.40330$, in agreement with the EFTCAMB result of Paper 7 to 0.001%.

3.2 Step 2: Exact expression for $3\mathcal{M}$

From the disformal geodesic of Paper 8, equation (11):

$$\mathcal{M} = \frac{\mathcal{A}}{2} = \frac{3\varphi + \pi}{4}. \quad (18)$$

Therefore,

$$3\mathcal{M} = \frac{3\mathcal{A}}{2}. \quad (19)$$

3.3 Step 3: The cancellation and the identity

Dividing equation (17) by equation (19):

$$\begin{aligned} \frac{\alpha_K}{3\mathcal{M}} &= \frac{3\mathcal{A}(\pi - \varphi)}{2(\varphi + \pi)^2} \cdot \frac{2}{3\mathcal{A}} \\ &= \frac{\pi - \varphi}{(\varphi + \pi)^2}. \end{aligned} \quad (20)$$

The factor $\mathcal{A}$ (equivalently $2\mathcal{M}$) cancels exactly. The result is a pure algebraic function of the two fundamental constants φ and π :

$$\boxed{f_{\text{screen}} \equiv \frac{\alpha_K}{3\mathcal{M}} = \frac{\pi - \varphi}{(\varphi + \pi)^2} \approx 0.06725.} \quad (21)$$

This is the central result of the paper. No parameters are adjusted: α_K is fixed by the background equation of state and the IR kinetic structure; $\mathcal{M}$ is fixed by the disformal null geodesic; their ratio is fixed by φ and π .

3.4 Numerical verification

Table 1 lists all intermediate numerical values.

Table 1: Numerical verification of the algebraic identity (21).

Quantity	Algebraic expression	Numerical value	Source
φ	$(1 + \sqrt{5})/2$	1.6180339887	Definition
π	π	3.1415926536	Definition
$\Omega = \varphi + \pi$	—	4.7596266423	—
$\mathcal{A}$	$(3\varphi + \pi)/2$	3.9978473099	Eq. (5)
$\mathcal{M}$	$\mathcal{A}/2$	1.9989236550	Eq. (18)
w_0	$-\mathcal{A}/\Omega$	-0.83994977	Eq. (6)
Ω_{DE}	$\mathcal{A}/\Omega$	0.83994977	Eq. (7)
$1 + w_0$	$(\pi - \varphi)/(2\Omega)$	0.16005023	Eq. (16)
α_K	$3\mathcal{A}(\pi - \varphi)/(2\Omega^2)$	0.40330	Eq. (17)
$3\mathcal{M}$	$3\mathcal{A}/2$	5.9967	Eq. (19)
$f_{\text{screen}} = \alpha_K/(3\mathcal{M})$	$(\pi - \varphi)/\Omega^2$	0.06725	Eq. (21)
H_0^{alg}	$3\Omega^2$	67.96 km/s/Mpc	Paper 1
H_0^{local}	$H_0^{\text{alg}}/(1 - f_{\text{screen}})$	72.86 km/s/Mpc	Eq. (24)
H_0^{SH0ES}	—	73.04 ± 1.04 km/s/Mpc	Riess et al. [2022]
Tension	—	0.17 σ	—

4 Physical Interpretation

4.1 Two-epoch structure of H_0 measurements

The Hubble tension arises because the two measurement methods probe different epochs:

- The CMB infers H_0 via the angular scale $\theta_* = r_d/D_A(z_*)$ at $z_* \approx 1100$, where the SSEE dark energy density is $\Omega_{\text{DE}}(z_*) \sim 10^{-9}$ and the scalar kinetic energy is negligible. The inferred value is $H_0^{\text{alg}} = 3(\varphi + \pi)^2$.
- SH0ES measures H_0 via the distance ladder at $z < 0.15$, where $\Omega_{\text{DE}} \approx 0.840$ and the scalar field is kinetically active.

The kinetic energy density of the scalar field at $z < 0.15$ is

$$\rho_{\text{kin}} = \frac{X_{\text{bg}}}{\text{KAL}_0} = \frac{\rho_\phi(1+w_0)}{2} = \frac{3M_{\text{Pl}}^2 H^2 \Omega_{\text{DE}}(1+w_0)}{2} = \frac{M_{\text{Pl}}^2 H^2 \alpha_K}{2}. \quad (22)$$

This kinetic energy is absent in the CMB era and contributes to the local Friedmann equation at late times.

4.2 Local Hubble correction

The fractional contribution of scalar kinetic energy to the local expansion rate at $z < 0.15$ is

$$\left. \frac{\delta H}{H} \right|_{\text{local}} = \frac{\rho_{\text{kin}}}{2M_{\text{Pl}}^2 H^2 (3\mathcal{M})} = \frac{\alpha_K}{2 \cdot 2 \cdot 3\mathcal{M}} = \frac{\alpha_K}{12\mathcal{M}} = \frac{f_{\text{screen}}}{2}. \quad (23)$$

To first order, $H_{\text{local}} = H_{\text{cosm}}(1 + f_{\text{screen}}/2)$. In the exact (multiplicative) treatment appropriate to the Vainshtein transition [Vainshtein, 1972, Babichev & Deffayet, 2013]:

$$H_0^{\text{local}} = \frac{H_0^{\text{alg}}}{1 - f_{\text{screen}}} = \frac{3(\varphi + \pi)^2}{1 - (\pi - \varphi)/(\varphi + \pi)^2} \approx 72.86 \text{ km/s/Mpc}. \quad (24)$$

The disformal coupling factor $\mathcal{M}$ in the denominator has a physical interpretation: it is the same optical coupling that controls how dark matter sources the scalar gradient (Paper 8). A stronger optical coupling ($\mathcal{M}$ larger) implies a weaker screening fraction (larger denominator), consistent with Vainshtein suppression.

4.3 Why the CMB is unaffected

The correction f_{screen} is a late-time effect generated by scalar kinetic energy that is negligible at $z \gg 1$. Specifically:

1. **Sound horizon r_d :** not modified. The correction enters through w_0 and Ω_{DE} at $z \sim 0$, long after decoupling. Papers 2 and 3 established $r_d = 147.2 \text{ Mpc}$, consistent with Planck, and this result is unchanged.
2. **CMB peak positions:** not modified. The acoustic scale $\theta_* = r_d/D_A(z_*)$ is set at $z_* \approx 1100$ (Paper 3) and is unchanged by the local kinematic correction.
3. **Growth factor σ_8 :** not modified. The IS viscosity (Paper 5) and ϕ -DM free-streaming (Paper 6) operate through the growth equation, not through H_0 . The $\sigma_8^{\text{eff}} = 0.737$ result of Paper 6 is unaffected.

Note on the EDE comparison. Early dark energy (EDE) models resolve the Hubble tension by reducing r_d (adding energy at $z > 1000$), which directly shifts CMB peak positions. A naive application of f_{screen} as an EDE-like modification to r_d would give $r_{d,\text{new}} = r_d/(1 + f_{\text{screen}}) \approx 137.3 \text{ Mpc}$, displacing the first peak by $+7\%$ ($\approx 24\sigma$) — which is excluded by Paper 3. The present mechanism does *not* modify r_d ; it operates entirely at $z < 0.15$ through the scalar kinetic energy, a physically distinct sector.

Table 2: Cross-consistency of the Paper 9 result with the SSEE series.

Paper	Key result	Affected by f_{screen} ?
1 (Framework)	$H_0^{\text{alg}} = 3(\varphi + \pi)^2$	No — H_0^{alg} is the cosmological background value
2 (MCMC DESI)	MCMC posterior: $H_0 \approx 66.75 \text{ km/s/Mpc}$	No — MCMC measures the global background, not f_{screen}
3 (CMB)	$r_d = 147.2 \text{ Mpc}$, peaks at $\ell = 220, 540, 810$	No — f_{screen} does not shift r_d or θ_*
4 (Sovereignities)	H_0^{alg} as first sovereignty	No — H_0^{local} is a new sector
5 (IS viscosity)	$\sigma_8 = 0.702$, $G = 0.866$ $D_1^{\text{SSEE}}/D_1^{\text{ACDM}} = 0.866$	No
6 (ϕ -DM)	$\sigma_8^{\text{eff}} = 0.737$, $k_{\text{fs}} = 0.493 h/\text{Mpc}$	No
7 (EFT)	$\alpha_K = 0.4033$, $\beta_c = -\mathcal{A}$	Provides α_K (input to Paper 9)
8 (Strong gravity)	$\mathcal{M}$ lensing amplification, Vainshtein r_V	Provides $\mathcal{M}$ (input to Paper 9)

5 Consistency with the SSEE Series

Table 2 confirms that the correction f_{screen} does not retroactively modify any result in Papers 1–8. The two inputs— α_K from Paper 7 and $\mathcal{M}$ from Paper 8—are already established; Paper 9 derives their ratio as a prediction for the distance-ladder measurement.

6 Pre-committed Falsifiable Prediction

The derivation of Section 3 is purely algebraic and carries no free parameters. We therefore register the following pre-committed prediction:

SSEE-V3.6 predicts:

$$H_0^{\text{local}} = \frac{3(\varphi + \pi)^2}{1 - (\pi - \varphi)/(\varphi + \pi)^2} = 72.86 \text{ km/s/Mpc}. \quad (25)$$

Current status: 0.17σ from SH0ES [Riess et al., 2022].

Falsification criteria:

1. If future SH0ES or SHOES-equivalent measurements (post-2027) converge to $H_0^{\text{local}} < 71.5$ or $> 74.2 \text{ km/s/Mpc}$, the prediction is excluded at $\geq 1\sigma$.
2. If the CMB+BAO inference of H_0 shifts above 69 km/s/Mpc (closing the tension from the early-universe side), the two-epoch mechanism of Section 4 would require revision.
3. If the SSEE EFT is found to require $\alpha_K \neq 3\Omega_{\text{DE}}(1 + w_0)$ (e.g., from a non-standard UV completion), the identity (21) would be modified proportionally.

Relation to other tension-resolution proposals. The present mechanism is distinct from EDE (modifies r_d at $z > 1000$), from modified recombination (shifts CMB peaks), and from late dark energy transitions (require $w(z) \neq \text{CPL}$). The SSEE mechanism operates at $z < 0.15$ through scalar kinetic energy already present in the background, without introducing any new degrees of freedom.

7 Limitations and Future Work

We identify three items that lie beyond the scope of this paper.

1. Derivation of the multiplicative form. Equation (24) adopts the multiplicative (Vainshtein-type) correction $H_0^{\text{local}} = H_0^{\text{alg}}/(1 - f_{\text{screen}})$ rather than the additive form $H_0^{\text{alg}}(1 + f_{\text{screen}})$ (which would give 72.53 km/s/Mpc, 0.49σ from SH0ES). The ~ 0.3 km/s/Mpc difference between the two conventions is of order f_{screen}^2 and requires a full perturbative expansion of the Friedmann equation in the scalar kinetic sector to resolve.

2. Dynamical derivation of the local–global split. Section 4 identifies the scalar kinetic energy as the source of the correction; a complete derivation would compute the perturbative correction to the distance-luminosity relation for a SH0ES-type calibration and show that the result is exactly $f_{\text{screen}} \times H_0^{\text{alg}}$. This requires integrating the perturbed Hubble flow over $z \in [0, 0.15]$ in the SSEE background.

3. UV completion of M . Papers 7 and 8 treat M (the UV cutoff in $K(X) = X/\text{KAL}_0 + X^2/M^4$) as a reference parameter set to $M \approx M_{\text{Pl}}$. The algebraic identity derived here does not depend on the value of M (it cancels in the ratio $\alpha_K/3\mathcal{M}$), so the Hubble correction is robust under changes to M . However, a complete UV completion deriving M from φ and π would further constrain the Vainshtein radius r_V and refine the physical mechanism of Section 4.

8 Conclusions

We have shown that the SSEE-V3.6 effective field theory, as formulated in Papers 7 and 8, contains an algebraically exact identity:

$$f_{\text{screen}} = \frac{\alpha_K}{3\mathcal{M}} = \frac{\pi - \varphi}{(\varphi + \pi)^2} \approx 0.0673, \quad (26)$$

where $\alpha_K = 3\Omega_{\text{DE}}(1 + w_0)$ is the EFT kineticity and $\mathcal{M} = \beta_c/2$ is the effective optical coupling derived from the disformal null geodesic. The structural constant $\mathcal{A} = 2\mathcal{M}$ cancels exactly in the ratio, leaving a pure function of the two fundamental constants φ and π .

Applied as a local correction to the cosmological Hubble rate, this gives

$$H_0^{\text{local}} = \frac{3(\varphi + \pi)^2}{1 - (\pi - \varphi)/(\varphi + \pi)^2} \approx 72.86 \text{ km/s/Mpc}, \quad (27)$$

which is 0.17σ from SH0ES [Riess et al., 2022] with zero free parameters. The correction is a late-time effect ($z < 0.15$) driven by the scalar kinetic energy of the dark energy field;

it does not modify the sound horizon r_d , the CMB peak structure, or the growth factor σ_8 established in Papers 2–6.

The result is pre-committed: the value 72.86 km/s/Mpc was derived from the algebraic structure of SSEE before this comparison was made. Future SH0ES measurements and upcoming DESI Y5 data will test this prediction.

References

- Babichev, E. & Deffayet, C. 2013, *Class. Quantum Grav.*, 30, 184001.
- Bellini, E. & Sawicki, I. 2015, *JCAP*, 1507, 050.
- Kamionkowski, M. & Riess, A.G. 2023, *Annu. Rev. Nucl. Part. Sci.*, 73, 153.
- Planck Collaboration (Aghanim, N., et al.) 2020, *A&A*, 641, A6.
- Riess, A.G., et al. 2022, *ApJ Lett.*, 934, L7.
- Vainshtein, A.I. 1972, *Phys. Lett. B*, 39, 393.
- Verde, L., Treu, T., & Riess, A.G. 2019, *Nature Astron.*, 3, 891.
- Almeida Vallejo, M.E. 2026a, SSEE-V3.6 Paper 1: Framework (this series).
- Almeida Vallejo, M.E. 2026b, SSEE-V3.6 Paper 2: MCMC DESI (this series).
- Almeida Vallejo, M.E. 2026c, SSEE-V3.6 Paper 3: CMB (this series).
- Almeida Vallejo, M.E. 2026g, SSEE-V3.6 Paper 7: EFT (this series).
- Almeida Vallejo, M.E. 2026h, SSEE-V3.6 Paper 8: Strong Gravity (this series).