



THE CLASSIC DMP ANGLE, PART 2 — X-RAY

LaRI predicts the occluded lower floors — restrict-masked so it only invents where the occluder actually is.

FILE `atlas_dmp_angle_xray_newyork_workflow.json` VERIFIED 2026-07-16

THE STORY

Part 1 grounded every facade whose base the camera can see. This variant handles the rest: the far buildings whose lower floors hide behind near rooflines. LaRI (layered ray intersections) predicts, per pixel, the ordered stack of surfaces each camera ray pierces — dense architecture is its documented strong domain, and 'the occluded lower floors behind a roofline' is the textbook use case.

The discipline that makes it presentable: the prediction is restrict-masked to the FOREGROUND DEPTH BAND ([0–45%] from AtlasDepthLayerMask, wired into restrict_mask — the node's own report recommends exactly this pairing). Substitution happens only where the occluder actually is; every visible surface keeps its real photographed depth. LaMa supplies the reveal pixels, and a mask-membership X-ray layer projects them onto the predicted geometry, so a dolly past the near roofline reveals plausible lower floors instead of a tear.

IN THE VIEWPORT — PROJECTED VS. THE GREY MESH



SOURCE PLATE

newyork_Birdseye.png · what's behind the rooflines? — the single still photograph everything below is reconstructed from.



PROJECT ON

The photo reassembled on the recovered camera — texels assigned by ray, perfect from this view.




🦴 X-RAY PROVENANCE

Red tint = surface whose depth was INVENTED by the hidden-geometry prediction — exactly what you're trusting the model for, nothing more.




PROJECT OFF

The same geometry as a grey mesh, recovered view — what the depth-derived surfaces actually are.



GREY · ORBIT LEFT

Orbited off-axis: the reconstructed cone. Coverage ends where the camera never saw.



GREY · ORBIT RIGHT

The opposite angle — silhouette tears read as deliberate holes, not errors.

NODES ON STAGE

AtlasPredictHiddenGeometry

AtlasDepthLayerMask

AtlasCleanPlateLayer

AtlasDepthMap

AtlasReferenceScaleSolve

LaMa inpaint

RETUNE ON YOUR PLATE

- ▶ **The fg band's far_pct (0.45).** Widen or narrow so the band covers exactly your occluding foreground — the restrict mask is derived from it.
- ▶ **Environment.** ATLAS_EXPERIMENTAL=1 + ATLAS_LARI_PATH → your LaRI clone (upstream has no license — research only).
- ▶ **Domain honesty.** Architecture is LaRI's strong domain (best registration of any plate here). Expect weaker coverage on open terrain — see the wreck workflow, which stress-tests exactly that.

RUN-VERIFIED

RUN ✓ Registration rel-MAD 0.113 — the best of any plate in the set. X-ray layer: 56k verts with matte + provenance; substitution confined to the fg band.

REQUIREMENTS

ATLAS_EXPERIMENTAL=1 · LaRI clone · comfyui-inpaint-nodes + big-lama.pt

