

Hartshorne Solutions

mlwells

2025

Feel free to add your Discord name to the author list. Let's add our usernames to our solutions, too, so that people get proper credit.

Contents

1.1 Affine Varieties	2
1.8	2
2.1 Sheaves	3
1.2	3

1.1 Affine Varieties

1.8

Let Y be an affine variety of dimension r in \mathbb{A}^n . Let H be a hypersurface in \mathbb{A}^n , and assume that $Y \not\subseteq H$. Then every irreducible component of $Y \cap H$ has dimension $r - 1$.

1.8 mlwells

Suppose that $H = Z(f)$ with f irreducible in $k[x_1, \dots, x_n]$. Then the projection of f in $A(Y) := k[x_1, \dots, x_n]/I(Y)$ is not equal to $\bar{0}$ since by assumption $(f) \not\subseteq I(Y)$. Since $A(Y)$ is a domain (due to the irreducibility of Y), the element \bar{f} is not a zero divisor. Assuming that $Y \cap H \neq \emptyset$, we have that \bar{f} is not a unit in $A(Y)$. To see this, let $P \in Y \cap H$. Then $I(Y), (f) \subset \mathfrak{m}_P$, the maximal ideal of $k[x_1, \dots, x_n]$ corresponding to P . This implies $(\bar{f}) \subset \mathfrak{m}_P/I(Y)$, the latter being a maximal ideal in $A(Y)$. Thus, \bar{f} is not a unit.

We apply Theorem 1.11A to get that every minimal prime ideal \mathfrak{p} in $A(Y)$ containing \bar{f} has height 1. The irreducible components of $Y \cap H$ and the minimal prime ideals containing (\bar{f}) correspond, and since each of these prime ideals has height 1, the corresponding varieties have dimension $r - 1$ by Theorem 1.8A.

2.1 Sheaves

1.2

- For any morphism of sheaves $\phi : \mathcal{F} \rightarrow \mathcal{G}$, show that for each point P , $(\ker \phi)_P = \ker(\phi_P)$ and $(\operatorname{im} \phi)_P = \operatorname{im}(\phi_P)$.
- Show that ϕ is injective (surjective) iff induced map on stalks ϕ_P is injective (surjective) for all P .
- Show that the sequence of sheaves $\cdots \rightarrow \mathcal{F}^{i-1} \rightarrow \mathcal{F}^i \rightarrow \mathcal{F}^{i+1} \rightarrow \cdots$ is exact iff corresponding sequences of stalks are exact for all P .

1.2 yakimk

- Since stalks are defined as a filtered colimit and kernels are a particular type of limit (pullback) and [filtered colimits commute with finite limits](#) isomorphism $(\ker \phi)_P = \ker(\phi_P)$ is automatic.

To show a similar thing for the image we first note that $(\operatorname{im} \phi)_P \simeq ((\operatorname{im}_{pre} \phi)^+)_P \simeq (\operatorname{im}_{pre} \phi)_P$, where the first isomorphism is by definition and the second one using the fact that sheafification has the same stalks as original presheaf. Now we conclude noticing that $\operatorname{im}_{pre} \phi \simeq \ker(\operatorname{coker} \phi)$, i.e. it is a finite limit of finite colimits and since again stalks are filtered colimits they commute with finite limits and colimits (in for instance abelian category) we conclude that $(\operatorname{im} \phi)_P \simeq \operatorname{im}(\phi_P)$.

- We show that a map of sheaves $\phi : \mathcal{F} \rightarrow \mathcal{G}$ is injective iff it induced maps on stalks are injective (proof for surjectivity is essentially identical).

Consider the following commutative diagram

$$\begin{array}{ccc} \mathcal{F}(U) & \xrightarrow{\phi_U} & \mathcal{G}(U) \\ \iota_1 \downarrow & & \downarrow \iota_2 \\ \prod_{P \in U} \mathcal{F}_P & \xrightarrow{\prod \tilde{\phi}_P} & \prod_{P \in U} \mathcal{G}_P \end{array}$$

Note that since section of a sheaf on an open set is determined by its stalks, ι_1 and ι_2 are both injective.

Suppose $\tilde{\phi}_P$ is injective for all P . Take two sections $f, g \in \mathcal{F}(U)$, since $\prod \tilde{\phi}_P$ and ι_1 are injective (former by the hypothesis that $\tilde{\phi}_P$ are injective), their composition is injective. If ϕ_U were not injective, we could find two different sections that go to the same class in $\prod \mathcal{G}_P$, which would contradict injectivity of $\prod \tilde{\phi}_P \circ \iota_1$.

Similarly if ϕ_U is injective then if some of $\tilde{\phi}_P$ were not injective, its composition with ι_1 would not be injective contradicting commutativity of the diagram above.

- We have to show that $\cdots \rightarrow \mathcal{F}^{i-1} \rightarrow \mathcal{F}^i \rightarrow \mathcal{F}^{i+1} \rightarrow \cdots$ is exact iff induced sequence of stalks are exact for all P .

Assuming that sequence of sheaves is exact we show induced sequences are exact at all stalks. By the previous section $\operatorname{im} \phi_P^i \simeq (\operatorname{im} \phi^i)_P \simeq (\ker \phi^{i+1})_P \simeq \ker \phi_P^{i+1}$.

Now assume that all sequences of stalks are exact. By the same reasoning as before we see that $\operatorname{im} \phi^i \simeq \ker \phi^{i+1}$ (since they are isomorphic on each stalk and hence are equal).