

Deriving the Luminosity of the Photosphere

MICHAEL MOSS^{1,2}

¹*The Department of Physics, The George Washington University, 725 21st NW, Washington, DC 20052, USA*

²*Astrophysics Science Division, NASA Goddard Space Flight Center, Greenbelt, MD 20771, USA*

Keywords: Gamma-Ray Bursts (629)

1. DERIVATION

This follows from the work of Hascoët et al. (2013).

The thermal energy injected into the outflow at the origin of the jet can be obtained from

$$\dot{E}_{\text{th}} = \epsilon_{\text{th}} \dot{E} = \epsilon_{\text{th}} \frac{\Omega}{4\pi} \dot{E}_{\text{iso}} = a T_0^4 c S_0 \quad (1)$$

where \dot{E} is the total power injected into the outflow, \dot{E}_{iso} is the isotropic equivalent total injected power, ϵ_{th} is the fraction of energy stored in thermal form, T_0 is the temperature at the origin of the flow, $S_0 = \pi \ell^2$ is the cross section of the outflow at the origin, ℓ is the initial jet radius, $a = 4\sigma/c$ is the radiation constant, σ is the Stephan-Boltzmann constant, c is the speed of light, and fraction of solid angle is $\frac{\Omega}{4\pi} \simeq \frac{\theta^2}{4}$, where θ is the jet opening angle.

The radius at which the material becomes transparent to photons and releases thermal radiation is known as the photosphere,

$$\begin{aligned} R_{\text{ph}} &\simeq \frac{\kappa \dot{M}}{8\pi c \Gamma^2} \\ &\simeq \frac{\kappa \frac{\dot{E}}{c^2 \Gamma}}{8\pi c \Gamma^2} = \frac{\kappa \dot{E}}{8\pi c^3 \Gamma^3} \\ &\simeq 2.9 \times 10^3 \frac{\kappa_{0.2} \dot{E}_{\text{iso},53}}{(1 + \sigma) \Gamma_2^3} \text{ cm} \end{aligned} \quad (2)$$

where κ ($\kappa_{0.2}$ in units of $0.2 \text{ cm}^2 \text{ g}^{-1}$) is the material opacity, Γ (Γ_2 in units of 100) the Lorentz factor of the shell, and σ is the magnetization of the material.

Assuming no dissipation takes place below the photosphere, mass and entropy conservation lead to the expressions

$$\beta \Gamma \rho S(R) = \text{Const.} \quad (3)$$

$$\frac{T}{\rho^{1/3}} = \text{Const.} \quad (4)$$

where ρ is the comoving density, $S(R) = \pi \theta^2 R^2$ is the surface perpendicular to the flow at a radius R , $\beta = v/c \sim 1$ where v is the velocity of the flow, and $\Gamma = (1 - \beta^2)^{-1/2}$. Equating the two above expressions leads to

$$\beta \Gamma T^3 S(R) = \text{const.} \quad (5)$$

allowing us to define the temperature at a radius R to be

$$T(R) \simeq T_0 \times (\theta^{-2/3} R^{-2/3} \ell^{2/3} \Gamma^{-1/3}) \quad (6)$$

Using Equations 1, 2, and 6, the luminosity of the thermal radiation released at the photosphere can be expressed as

$$\begin{aligned} L_{\text{th}} &= \Gamma^2 a T^4(R_{\text{ph}}) c S(R_{\text{ph}}) \\ &= \dot{E}_{\text{th}} \times (\theta^{-2/3} R_{\text{ph}}^{-2/3} \ell^{2/3} \Gamma^{2/3}) \\ &= \epsilon_{\text{th}} \dot{E} \theta^{-2/3} \ell^{2/3} \left(\frac{\kappa \dot{E}}{8\pi c^3 \Gamma^3} \right)^{-2/3} \Gamma^{2/3} \\ &= \epsilon_{\text{th}} c^2 \dot{E}^{1/3} \left(\frac{8\pi \ell}{\kappa \theta} \right)^{2/3} \Gamma^{8/3} \end{aligned} \quad (7)$$

So, we can see that the photospheric luminosity is strongly dependent on only the Lorentz factor of the material in the outflow.

Questions:

1. Why is there a factor of Γ^2 in Equation 7?

⁶¹ 2. In the equation for R_{ph} (Eq 9 in [Hascoët et al.](#) ⁶³ equivalent energy and not the beaming corrected?
⁶² [2013](#)), why is \dot{M} associated with the isotropic ⁶⁴ (Note in Eq 2 here I write the beaming correct,
⁶⁵ but this leads to incorrect results.)

REFERENCES

⁶⁶ Hascoët, R., Daigne, F., & Mochkovitch, R. 2013, A&A,
⁶⁷ 551, A124, doi: [10.1051/0004-6361/201220023](https://doi.org/10.1051/0004-6361/201220023)