

# tParton: A Python package for next-to-leading order evolution of transversity parton distribution functions

Congzhou M Sha<sup>1</sup> and Bailing Ma<sup>2</sup>

<sup>1</sup> Penn State College of Medicine, Hershey, PA 17033, USA <sup>2</sup> Wake Forest University School of Medicine, Winston-Salem, NC 27101, USA ¶ Corresponding author

DOI: [10.xxxxxx/draft](https://doi.org/10.xxxxxx/draft)

## Software

- [Review](#) ↗
- [Repository](#) ↗
- [Archive](#) ↗

Editor: [Open Journals](#) ↗

## Reviewers:

- [@openjournals](#)

Submitted: 01 January 1970

Published: unpublished

## License

Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License ([CC BY 4.0](#)).

## Summary

Parton distribution functions (PDFs) describe the probability of finding quarks and gluons (collectively called partons) within hadrons such as protons and neutrons. These functions are fundamental to our understanding of quantum chromodynamics (QCD) and are essential for interpreting high-energy physics experiments. The transversity PDF, which encodes information about the transverse spin structure of hadrons, is particularly challenging to measure experimentally and has been less studied computationally compared to unpolarized and helicity PDFs.

tParton is a Python package that implements two distinct methods for solving the Dokshitzer–Gribov–Lipatov–Altarelli–Parisi (DGLAP) evolution equations for transversity PDFs at leading order (LO) and next-to-leading order (NLO) in perturbative QCD. The package provides both a command-line interface and a Python API, making it accessible for both quick calculations and integration into larger analysis workflows.

## Statement of need

PDFs must be evolved from one energy scale to another to enable comparisons between different experiments and theoretical predictions. While numerous codes exist for evolving unpolarized and helicity PDFs (such as QCDNUM (Botje, 2011), EKO (Candido et al., 2022), HOPPET (Salam & Rojo, 2009), and APFEL++ (Bertone et al., 2014, 2017)), options for transversity PDF evolution are limited. The original Fortran implementation by Hirai et al. (Hirai et al., 1998) is nearly 30 years old and no longer easily accessible. APFEL++ (Bertone et al., 2017) provides an implementation using direct numerical integration, but no publicly available code has implemented the alternative Mellin moment method proposed by Vogelsang (Vogelsang, 1998).

tParton fills this gap by providing:

1. **Two complementary methods:** A direct integration method (following Hirai et al.) and a Mellin moment method (following Vogelsang), allowing users to choose based on their accuracy and computational needs.
2. **Modern Python implementation:** Built on NumPy (Harris et al., 2020) and SciPy (Virtanen et al., 2020), tParton is easy to install via pip and integrates seamlessly with the Python scientific computing ecosystem.
3. **Comprehensive validation:** The package includes extensive examples and validation against both Mathematica implementations and APFEL++ results, with detailed discussion of discretization effects and method comparisons.

39 **4. Dual interface:** Both command-line tools for standalone use and importable modules for  
40 integration into larger projects.

41 The package is aimed at researchers in hadronic physics, particularly those analyzing semi-  
42 inclusive deep inelastic scattering experiments and studying nucleon spin structure. It has been  
43 validated and documented in a detailed preprint (Sha & Ma, 2025).

## 44 Implementation

45 tParton implements the DGLAP evolution equation for the transversity PDF:

$$\frac{\partial}{\partial t} \Delta_T q^\pm(x, t) = \frac{\alpha_s(t)}{2\pi} \Delta_T P_{q^\pm}(x) \otimes \Delta_T q^\pm(x, t)$$

46 where  $t = \ln Q^2$ ,  $Q^2$  is the energy scale,  $\Delta_T P_{q^\pm}$  is the transversity splitting function, and  $\otimes$   
47 denotes Mellin convolution defined by:

$$f(x) \otimes g(x) := \int_x^1 \frac{dy}{y} f\left(\frac{x}{y}\right) g(y)$$

### 48 Method 1: Direct integration (Hirai method)

49 The first method discretizes both the momentum fraction  $x$  and energy scale  $Q^2$  into grids and  
50 solves the integro-differential equation using the Euler method for  $Q^2$  evolution and Simpson's  
51 rule for  $x$  integration. This approach is straightforward but can be computationally expensive  
52 for fine grids.

### 53 Method 2: Mellin moment method (Vogelsang method)

54 The second method exploits the convolution theorem for Mellin transforms. The solution is  
55 expressed in terms of Mellin moments:

$$\mathcal{M}[\Delta_T q^\pm](Q^2; s) = K(s, Q^2, Q_0^2) \mathcal{M}[\Delta_T q^\pm](Q_0^2; s)$$

56 where  $K$  contains the evolution kernel depending on the splitting function moments. Since  
57 analytic expressions for the evolution kernel are available, this method obviates the need to  
58 solve the ODE. The evolved PDF is reconstructed via inverse Mellin transform using the Cohen  
59 contour method. This approach is typically faster for evaluation of the transversity PDF at  
60 single points, and less sensitive to discretization for smooth PDFs.

61 Both methods support LO and NLO evolution with numerically exact or analytical forms of  
62 the running coupling constant  $\alpha_s(Q^2)$ . See our arXiv preprint for detailed computational  
63 complexity analysis (Sha & Ma, 2025).


64 Our primary future goal is to implement NNLO evolution, for which analytic Mellin moment  
65 expressions have been recently calculated (Blümlein et al., 2021).

## 66 Examples and validation


67 The package includes extensive Jupyter notebooks in the examples/ directory that:

- 68 ■ Generate initial transversity distributions based on literature models (Hirai et al., 1998)
- 69 ■ Compare both evolution methods against each other and against APFEL++
- 70 ■ Demonstrate sensitivity to numerical parameters (grid resolution, timesteps)

- 71     ■ Reproduce figures from the associated preprint (Sha & Ma, 2025)
- 72     A separate Mathematica notebook validates the analytical expressions for the Mellin moments
- 73     of the splitting functions, providing an independent check of the theoretical framework. The
- 74     key validation figures from our preprint are shown below:



**Figure 1:** Evolution of the initial GS-A type PDF with numerically evolved  $\alpha_s(Q^2)$ , comparing the Hirai method (Python), the Vogelsang method (Python), Mathematica validation, and APFEL++ evolution. APFEL++ evolution data was provided by V Bertone.



**Figure 2:** Relative error of the various methods, with the Hirai method (Python) as control.

- 75     Users can evolve a PDF with a single command:
- ```
python -m tparton m input.dat 3.1 10.6 --morp plus -o output.dat
```
- 76     Or import and use the package programmatically:
- ```
from tparton.m_evolution import evolve
result = evolve(input_pdf, Q0_squared=3.1, Q_squared=10.6,
               morp='plus', order='NLO')
```

Complete online documentation of the API and detailed examples are available on the [GitHub Pages](#) associated with the repository.

## Acknowledgements

We acknowledge helpful discussions with colleagues in the hadronic physics community and thank the maintainers of APFEL++ for providing comparison benchmarks.

## References

- Bertone, V., Carrazza, S., & Nocera, E. R. (2017). APFEL++: A new PDF evolution library in C++14. *European Physical Journal C*, 77(8), 516. <https://doi.org/10.1140/epjc/s10052-017-5088-y>
- Bertone, V., Carrazza, S., & Rojo, J. (2014). APFEL: A PDF evolution library with QED corrections. *Computer Physics Communications*, 185, 1647–1668. <https://doi.org/10.1016/j.cpc.2014.03.007>
- Blümlein, J., Marquard, P., Schneider, C., & Schönwald, K. (2021). The three-loop unpolarized and polarized non-singlet anomalous dimensions from off shell operator matrix elements. *Nuclear Physics B*, 971, 115542. <https://doi.org/10.1016/j.nuclphysb.2021.115542>
- Botje, M. (2011). QCDNUM: Fast QCD evolution and convolution. *Computer Physics Communications*, 182(2), 490–532. <https://doi.org/10.1016/j.cpc.2010.10.020>
- Candido, A., Hekhorn, F., & Magni, G. (2022). EKO: Evolution kernel operators. *European Physical Journal C*, 82(10), 976. <https://doi.org/10.1140/epjc/s10052-022-10878-w>
- Harris, C. R., Millman, K. J., Walt, S. J. van der, Gommers, R., Virtanen, P., Cournapeau, D., Wieser, E., Taylor, J., Berg, S., Smith, N. J., Kern, R., Picus, M., Hoyer, S., Kerkwijk, M. H. van, Brett, M., Haldane, A., Río, J. F. del, Wiebe, M., Peterson, P., ... Oliphant, T. E. (2020). Array programming with NumPy. *Nature*, 585(7825), 357–362. <https://doi.org/10.1038/s41586-020-2649-2>
- Hirai, M., Kumano, S., & Miyama, M. (1998). Numerical solution of Q2 evolution equation for the transversity distribution  $\Delta_T q$ . *Computer Physics Communications*, 111(1), 150–166. [https://doi.org/10.1016/S0010-4655\(98\)00028-9](https://doi.org/10.1016/S0010-4655(98)00028-9)
- Salam, G. P., & Rojo, J. (2009). A higher order perturbative parton evolution toolkit (HOPPET). *Computer Physics Communications*, 180, 120–156. <https://doi.org/10.1016/j.cpc.2008.08.010>
- Sha, C. M., & Ma, B. (2025). *tParton: Implementation of next-to-leading order evolution of transversity parton distribution functions*. <https://arxiv.org/abs/2409.00221>
- Virtanen, P., Gommers, R., Oliphant, T. E., Haberland, M., Reddy, T., Cournapeau, D., Burovski, E., Peterson, P., Weckesser, W., Bright, J., van der Walt, S. J., Brett, M., Wilson, J., Millman, K. J., Mayorov, N., Nelson, A. R. J., Jones, E., Kern, R., Larson, E., ... SciPy 1.0 Contributors. (2020). SciPy 1.0: Fundamental algorithms for scientific computing in Python. *Nature Methods*, 17, 261–272. <https://doi.org/10.1038/s41592-019-0686-2>
- Vogelsang, W. (1998). Next-to-leading order evolution of transversity distributions and Soffer's inequality. *Physical Review D*, 57, 1886–1894. <https://doi.org/10.1103/PhysRevD.57.1886>