

# tParton: A Python package for next-to-leading order evolution of transversity parton distribution functions

Congzhou M Sha<sup>1</sup> and Bailing Ma<sup>2</sup>

<sup>1</sup> Penn State College of Medicine, Hershey, PA 17033, USA <sup>2</sup> Wake Forest University School of Medicine, Winston-Salem, NC 27101, USA ¶ Corresponding author

DOI: [10.xxxxxx/draft](https://doi.org/10.xxxxxx/draft)

## Software

- [Review](#)
- [Repository](#)
- [Archive](#)

Editor: [Open Journals](#)

## Reviewers:

- [@openjournals](#)

Submitted: 01 January 1970

Published: unpublished

## License

Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License ([CC BY 4.0](#)).

## Summary

Parton distribution functions (PDFs) describe the probability of finding quarks and gluons (collectively called partons) within hadrons such as protons and neutrons. These functions are fundamental to our understanding of quantum chromodynamics (QCD) and are essential for interpreting high-energy physics experiments. The transversity PDF, which encodes information about the transverse spin structure of hadrons, is particularly challenging to measure experimentally and has been less studied computationally compared to unpolarized and helicity PDFs.

tParton is a Python package that implements two distinct methods for solving the Dokshitzer–Gribov–Lipatov–Altarelli–Parisi (DGLAP) evolution equations for transversity PDFs at leading order (LO) and next-to-leading order (NLO) in perturbative QCD. The package provides both a command-line interface and a Python API, making it accessible for both quick calculations and integration into larger analysis workflows.

## Statement of need

PDFs must be evolved from one energy scale to another to enable comparisons between different experiments and theoretical predictions. While numerous codes exist for evolving unpolarized and helicity PDFs (such as QCDNUM (Botje, 2011), EKO (Candido et al., 2022), HOPPET (Salam & Rojo, 2009), and APFEL++ (Bertone et al., 2014, 2017)), options for transversity PDF evolution are limited. The original Fortran implementation by Hirai et al. (Hirai et al., 1998) is nearly 30 years old and no longer accessible. APFEL++ (Bertone et al., 2017) provides an implementation using direct numerical integration, but no publicly available code has implemented the alternative Mellin moment method proposed by Vogelsang (Vogelsang, 1998).

tParton fills this gap by providing:

1. **Two complementary methods:** A direct integration method (following Hirai et al.) and a Mellin moment method (following Vogelsang), allowing users to choose based on their accuracy and computational needs.
2. **Modern Python implementation:** Built on NumPy (Harris et al., 2020) and SciPy (Virtanen et al., 2020), tParton is easy to install via pip and integrates seamlessly with the Python scientific computing ecosystem.
3. **Comprehensive validation:** The package includes extensive examples and validation against both Mathematica implementations and APFEL++ results, with detailed discussion of discretization effects and method comparisons.

39     **4. Dual interface:** Both command-line tools for standalone use and importable modules for  
40     integration into larger projects.

41     The package is aimed at researchers in hadronic physics, particularly those analyzing semi-  
42     inclusive deep inelastic scattering experiments and studying nucleon spin structure. It has been  
43     validated and documented in a detailed preprint (Sha & Ma, 2024).

## 44     **Implementation**

45     tParton implements the DGLAP evolution equation for the transversity PDF:

$$\frac{\partial}{\partial t} \Delta_T q^\pm(x, t) = \frac{\alpha_s(t)}{2\pi} \Delta_T P_{q^\pm}(x) \otimes \Delta_T q^\pm(x, t)$$

46     where  $t = \ln Q^2$ ,  $Q^2$  is the energy scale,  $\Delta_T P_{q^\pm}$  is the transversity splitting function, and  $\otimes$   
47     denotes Mellin convolution.

### 48     **Method 1: Direct integration (Hirai method)**

49     The first method discretizes both the momentum fraction  $x$  and energy scale  $Q^2$  into grids and  
50     solves the integro-differential equation using the Euler method for  $Q^2$  evolution and Simpson's  
51     rule for  $x$  integration. This approach is straightforward but can be computationally expensive  
52     for fine grids.

### 53     **Method 2: Mellin moment method (Vogelsang method)**

54     The second method exploits the convolution theorem for Mellin transforms. The solution is  
55     expressed in terms of Mellin moments:

$$\mathcal{M}[\Delta_T q^\pm](Q^2; s) = K(s, Q^2, Q_0^2) \mathcal{M}[\Delta_T q^\pm](Q_0^2; s)$$

56     where  $K$  contains the evolution kernel depending on the splitting function moments. The  
57     evolved PDF is reconstructed via inverse Mellin transform using the Talbot contour integration  
58     method. This approach is typically faster and less sensitive to discretization for smooth PDFs.

59     Both methods support LO and NLO evolution with exact or analytical forms of the running  
60     coupling constant  $\alpha_s(Q^2)$ .

## 61     **Examples and validation**

62     The package includes extensive Jupyter notebooks in the `examples/` directory that:

- 63     ▪ Generate initial transversity distributions based on literature models (Hirai et al., 1998)
- 64     ▪ Compare both evolution methods against each other and against APFEL++
- 65     ▪ Demonstrate sensitivity to numerical parameters (grid resolution, timesteps)
- 66     ▪ Reproduce figures from the associated preprint

67     A separate Mathematica notebook validates the analytical expressions for the Mellin moments  
68     of the splitting functions, providing an independent check of the theoretical framework.

69     Users can evolve a PDF with a single command:

```
python -m tparton m input.dat 3.1 10.6 --morp plus -o output.dat
```

70     Or import and use the package programmatically:

```
from tparton.m_evolution import evolve
result = evolve(input_pdf, Q0_squared=3.1, Q_squared=10.6,
               morp='plus', order='NLO')
```

## Acknowledgements

We acknowledge helpful discussions with colleagues in the hadronic physics community and thank the maintainers of APFEL++ for providing comparison benchmarks.

## References

- Bertone, V., Carrazza, S., & Nocera, E. R. (2017). APFEL++: A new PDF evolution library in C++14. *European Physical Journal C*, 77(8), 516. <https://doi.org/10.1140/epjc/s10052-017-5088-y>
- Bertone, V., Carrazza, S., & Rojo, J. (2014). APFEL: A PDF evolution library with QED corrections. *Computer Physics Communications*, 185, 1647–1668. <https://doi.org/10.1016/j.cpc.2014.03.007>
- Botje, M. (2011). QCDNUM: Fast QCD evolution and convolution. *Computer Physics Communications*, 182(2), 490–532. <https://doi.org/10.1016/j.cpc.2010.10.020>
- Candido, A., Hekhorn, F., & Magni, G. (2022). EKO: Evolution kernel operators. *European Physical Journal C*, 82(10), 976. <https://doi.org/10.1140/epjc/s10052-022-10878-w>
- Harris, C. R., Millman, K. J., Walt, S. J. van der, Gommers, R., Virtanen, P., Cournapeau, D., Wieser, E., Taylor, J., Berg, S., Smith, N. J., Kern, R., Picus, M., Hoyer, S., Kerkwijk, M. H. van, Brett, M., Haldane, A., Río, J. F. del, Wiebe, M., Peterson, P., ... Oliphant, T. E. (2020). Array programming with NumPy. *Nature*, 585(7825), 357–362. <https://doi.org/10.1038/s41586-020-2649-2>
- Hirai, M., Kumano, S., & Miyama, M. (1998). Numerical solution of Q2 evolution equation for the transversity distribution  $\Delta_T q$ . *Computer Physics Communications*, 111(1), 150–166. [https://doi.org/10.1016/S0010-4655\(98\)00028-9](https://doi.org/10.1016/S0010-4655(98)00028-9)
- Salam, G. P., & Rojo, J. (2009). A higher order perturbative parton evolution toolkit (HOPPET). *Computer Physics Communications*, 180, 120–156. <https://doi.org/10.1016/j.cpc.2008.08.010>
- Sha, C. M., & Ma, B. (2024). *tParton: Implementation of next-to-leading order evolution of transversity parton distribution functions*. <https://arxiv.org/abs/2409.00221>
- Virtanen, P., Gommers, R., Oliphant, T. E., Haberland, M., Reddy, T., Cournapeau, D., Burovski, E., Peterson, P., Weckesser, W., Bright, J., van der Walt, S. J., Brett, M., Wilson, J., Millman, K. J., Mayorov, N., Nelson, A. R. J., Jones, E., Kern, R., Larson, E., ... SciPy 1.0 Contributors. (2020). SciPy 1.0: Fundamental algorithms for scientific computing in Python. *Nature Methods*, 17, 261–272. <https://doi.org/10.1038/s41592-019-0686-2>
- Vogelsang, W. (1998). Next-to-leading order evolution of transversity distributions and Soffer's inequality. *Physical Review D*, 57, 1886–1894. <https://doi.org/10.1103/PhysRevD.57.1886>