

# THE BORSUK-ULAM THEOREM IN REAL-COHESIVE HOMOTOPY TYPE THEORY

DANIEL CICALA, CHANDRIKA SADANAND, MICHAEL SHULMAN,  
AND AMELIA TEBBE

ABSTRACT. Borsuk-Ulam!

## WRITING NOTES

Writing assignments:

- Amelia—section 5
- Chandrika—section 4
- Daniel—sections 2 and 3

Formalizing the cohomology proofs will be determined later.

## 1. INTRODUCTION

## 2. OVERVIEW OF REAL-COHESIVE HOMOTOPY TYPE THEORY

OUTLINE:

- HoTT as foundations
- Interpreting AlgTop theorems in HoTT is obstructed by discontinuous functions
- Relating continuous and discontinuous with flat and sharp, which are borrowed from cohesive topoi
- Formalizing flat and sharp in HoTT + axioms needed, e.g. Rflat
- Connecting sets used in AlgTop with HITs used in HoTT via shape

## 3. TRANSLATING BORSUK-ULAM TO HOMOTOPY TYPE THEORY

OUTLINE:

- **Subsection 1.** Give statements for BU-classic, BU-odd, BU-retract) a la wikipedia. The proof strategy: show BU-retract implies BU-odd which is equivalent to BU-classic, then prove BU-retract. Give the proof for BU-retract.
- **Subsection 2.** Translate the classical statement into propositions as types. We want to model classical proof. The failure of contrapositive rule in constructive logic—(not  $q$  implies not  $p$ ) is ( $p$  implies not not  $q$ )—means our proof strategy is BU-retract implies not not BU-odd which is equivalent to not not BU-classic. But not not BU-classic is sharp BU-classic. Prove BU-retract.
- To close out the section, list the ingredients we need to prove BU-retract.

## 4. TOPOLOGICAL AND HOMOTOPICAL REAL PROJECTIVE SPACES

## OUTLINE:

- Define  $n$ -disks as both sets and types, the latter which is simply 1, since they're contractible. Show that  $\int \mathbb{D} = \mathbb{D}$
- Define  $n$ -spheres as sets. Use pushouts to glue disks together. Explain why we need to glue with a collar—i.e. the “topology” (as encoded by continuous paths  $\mathbb{R} \rightarrow \mathbf{X}$  of a type  $\mathbf{X}$ . Show, via Shulman, that  $\int \mathbb{S}^n = \mathbb{S}^n$
- Define  $\mathbb{R}P^n$  as sets using pushouts and collaring. Recall Bulcholtz and Egbert's definition of HIT  $\mathbb{R}P^n$ . Prove that  $\int RRP^n = \mathbb{R}P^n$

## 5. COHOMOLOGY

## OUTLINE:

- Subsection 1. Define cohomology for  $\mathbb{Z}/2\mathbb{Z}$  coefficients and the EM-spaces for  $\mathbb{R}P^n$
- **Subsection 2.** Show that we get a commutative graded ring structure for cohomology of any type  $\mathbf{X}$  with  $\mathbb{Z}/2\mathbb{Z}$ -coefficients. Follow Brunerie's thesis.
- **Subsection 3.** Compute  $\mathbb{Z}/2\mathbb{Z}$ -cohomology ring for  $\mathbb{R}P^n$  using Mayer-Vietoris. This needs us to first compute cohomology for disks and spheres.

## 6. THE BORSUK-ULAM THEOREM

## OUTLINE:

- The proof is done by this point. Just put it all together and reconnect the dots for the reader.

## REFERENCES

- [BR17] Ulrik Buchholtz and Egbert Rijke. The real projective spaces in homotopy type theory. To appear in LICS'17. arXiv:1704.05770, 2017.
- [Bru16] Guillaume Brunerie. *On the homotopy groups of spheres in homotopy type theory*. PhD thesis, Université de Nice, 2016. arXiv:1606.05916.
- [Shu17] Michael Shulman. Brouwer's fixed-point theorem in real-cohesive homotopy type theory. To appear in MSCS. arXiv:1509.07584, 2017.