
10/28/2026: Big O, Big Omega, and Big Theta Notations

CSCI 246: Discrete Structures

Textbook reference: Ch 11.2, Epp

Today's Agenda

- Mini-lecture ($\approx$ 30 mins)
- Group exercises ($\approx$ 20 mins)


Quiz 7 Scores
median=16.7
Frequency
Values (as percentage)
0
1
2
3
4
5
6
7
8
0
20
40
60
80
100
120
140

Study Guide For Quiz 8

Group exercises. Know how to solve the group exercises from the course meetings on Induction (10/14) and Recurrence (10/16).

Readings.

Sec. 22 (Induction)

- Know how to use Theorem 22.2 / Proof Template 17 to prove Props. 22.3 and 22.5.
- Understand the difference between Proof Template 17 and 18.

Sec. 23 (Recurrence)

- Know how to solve first-order recurrences from Props 23.1 and 23.3. For example, see Example 23.2.
- Know how to solve second-order recurrences from Theorems 23.5 and 23.9. For example, see Example 23.6.
- Know how to use Theorem 23.17 to find a polynomial expression to generate the first few terms of a sequence, as in Example 23.18.

Important Note. The max of your quiz 7 and quiz 8 scores will be used for both.

Mini-lecture:
Big O, Big Omega, Big Theta Notations

Which would you rather have?

Option A

$1,000,000 on Nov 1

Option B

$1 on Nov 1,
doubled by the
end of every day

Which would you rather have?

Option A

$1,000,000 on Nov 1



Hopper chooses Option A.

Option B

$1 on Nov 1,
doubled by the
end of every day



Bob chooses Option B.

Which would you rather have?

Option A

$1,000,000 on Nov 1



Hopper chooses Option A.

On Nov 30:

$1,000,000

Option B

$1 on Nov 1,
doubled by the
end of every day



Bob chooses Option B.

On Nov 30:

$1,073,741,824

Why?

Why?

Let n be the date in November, $f(n)$ be the value of your bank account on date n.

Why?


Let n be the date in November, $f(n)$ be the value of your bank account on date n.

Option A

$$f(n) = 1,000,000$$

Option B

$$g(n) = 2^n$$



$g(n) = 2^n$ is eventually greater than $f(n) = 1,000,000$.

Why?


Let n be the date in November, $f(n)$ be the value of your bank account on date n.

Option A

$$f(n) = 1,000,000$$

Option B

$$g(n) = 2^n$$



$\underbrace{g(n) = C2^n}_{\text{exponential}}$ is eventually greater than $\underbrace{f(n) = D}_{\text{constant}}$.

Which algorithm would you rather apply to a large dataset?

Option A

Passes through each sample 10,000 times.

Option B

Passes through each pair of samples once.

Which algorithm would you rather apply to a large dataset?

Option A

Passes through each sample 10,000 times.



Bob chooses Option A.

Option B

Passes through each pair of samples once.



Hopper chooses Option B.

Which algorithm would you rather apply to a large dataset?

Option A

Passes through each sample 10,000 times.



Bob chooses Option A.

10 billion

Option B

Passes through each pair of samples once.



Hopper chooses Option B.

Computations on the Million Song Dataset:

$\approx \frac{1}{2}$ **trillion**

Why?

Why?

Let n be the number of data points, $f(n)$ be the number of computations.

Why?


Let n be the number of data points, $f(n)$ be the number of computations.

Option A

$$f(n) = 10,000n$$

Option B

$$g(n) = n^2$$



$g(n) = n^2$ is eventually greater than $f(n) = 10,000n$.

Why?


Let n be the number of data points, $f(n)$ be the number of computations.

Option A

$$f(n) = 10,000n$$

Option B

$$g(n) = n^2$$



$\underbrace{g(n) = Cn^2}_{\text{quadratic}}$ is eventually greater than $\underbrace{f(n) = Dn}_{\text{linear}}$.

How to compare the efficiency of algorithms?



Figure 1: German mathematician Paul Bachmann

How to compare the efficiency of algorithms?





Figure 1: German mathematician Paul Bachmann

Definition

Let f and g be real-valued non-negative functions defined on the same set of nonnegative integers.

Then f is of order at most g, written $\boldsymbol{f(n)}$ **is** $\boldsymbol{O(g(n))}$ (f of n is big-O of g of n) if and only if there exist positive real number B and integer b such that

$$|f(n)| \leq B|g(n)| \quad \text{for every integer } n \geq b$$




Definition

Let f and g be real-valued non-negative functions defined on the same set of nonnegative integers.

Then f is of order at most g, written $\boldsymbol{f(n)}$ **is** $\boldsymbol{O(g(n))}$ (f of n is big-O of g of n) if and only if there exist positive real number B and integer b such that

$$|f(n)| \leq B|g(n)| \quad \text{for every integer } n \geq b$$



Poll. How would you summarize this definition in words?

Poll. How would you summarize this definition in words?

Poll. How would you summarize this definition in words?

Solution. The function f is eventually smaller in size than some positive multiple of g.

Poll. How would you summarize this definition in words?

Solution. The function f is eventually smaller in size than some positive multiple of g.


We say that "f is of order at most g".

Poll. How would you summarize this definition in words?

Solution. The function f is eventually smaller in size than some positive multiple of g.


We say that "f is of order at most g".

Classification Theorem



Poll. How should we interpret this graph?


Classification Theorem



Poll. How should we interpret this graph?

- A taxonomy of growth rates (or scaling behavior): Any function is big O of everything equal to or above it, but is **not** big O of anything below it.


Classification Theorem



Poll. How should we interpret this graph?

- A taxonomy of growth rates (or scaling behavior): Any function is big O of everything equal to or above it, but is **not** big O of anything below it.
- **Example**: n is big O of n, n^2, 2^n, $n!$, and n^n. Even if we multiply n by a large constant, those functions will always eventually get bigger!

Classification Theorem



Poll. How should we interpret this graph?

- A taxonomy of growth rates (or scaling behavior): Any function is big O of everything equal to or above it, but is **not** big O of anything below it.
- **Example**: n is big O of n, n^2, 2^n, $n!$, and n^n. Even if we multiply n by a large constant, those functions will always eventually get bigger!
- **Anti-Example**: n is not big O of $n^{\frac{1}{2}}$. No matter how much you scale up $n^{\frac{1}{2}}$ by a large constant, n will eventually get bigger!

How to check Big O membership?

Example: $f(n) = 3n + 100$ is $O(n)$

Proof. Take $n \geq 1$. Then

$$|3n + 100| \leq |3n + 100n| = 103|n|.$$

(To satisfy the definition of big O, we take $b = 1, B = 103$.)

How to check Big O membership?

Example: $f(n) = 3n + 100$ is $O(n)$

Proof. Take $n \geq 1$. Then

$$|3n + 100| \leq |3n + 100n| = 103|n|.$$

(To satisfy the definition of big O, we take $b = 1, B = 103$.)

Example: $f(n) = 9n^2 + 3n - 2$ is $O(n^2)$.

Proof. Take $n \geq 1$. Then

$$|9n^2 + 3n - 2| \leq |9n^2 + 3n^2 + 2n^2| = 14|n^2|.$$

(To satisfy the definition of big O, we take $b = 1, B = 14$.)

How to check Big O membership?

Example: $f(n) = 3n + 100$ is $O(n)$

Proof. Take $n \geq 1$. Then

$$|3n + 100| \leq |3n + 100n| = 103|n|.$$

(To satisfy the definition of big O, we take $b = 1, B = 103$.)

Example: $f(n) = 9n^2 + 3n - 2$ is $O(n^2)$.

Proof. Take $n \geq 1$. Then

$$|9n^2 + 3n - 2| \leq |9n^2 + 3n^2 + 2n^2| = 14|n^2|.$$

(To satisfy the definition of big O, we take $b = 1, B = 14$.)

Remark. Big O notation hides irrelevant details.

How to check Big O membership?

Example: $f(n) = 3n + 100$ is $O(n)$

Proof. Take $n \geq 1$. Then

$$|3n + 100| \leq |3n + 100n| = 103|n|.$$

(To satisfy the definition of big O, we take $b = 1, B = 103$.)

Example: $f(n) = 9n^2 + 3n - 2$ is $O(n^2)$.

Proof. Take $n \geq 1$. Then

$$|9n^2 + 3n - 2| \leq |9n^2 + 3n^2 + 2n^2| = 14|n^2|.$$

(To satisfy the definition of big O, we take $b = 1, B = 14$.)

Remark. Big O notation hides irrelevant details.

Poll. Do you see a pattern?

How to check Big O membership?

Example: $f(n) = 3n + 100$ is $O(n)$

Proof. Take $n \geq 1$. Then

$$|3n + 100| \leq |3n + 100n| = 103|n|.$$

(To satisfy the definition of big O, we take $b = 1, B = 103$.)

Example: $f(n) = 9n^2 + 3n - 2$ is $O(n^2)$.

Proof. Take $n \geq 1$. Then

$$|9n^2 + 3n - 2| \leq |9n^2 + 3n^2 + 2n^2| = 14|n^2|.$$

(To satisfy the definition of big O, we take $b = 1, B = 14$.)

Remark. Big O notation hides irrelevant details.

Poll. Do you see a pattern?

Solution. Polynomials are big O of their highest-order term.

How to check Big O membership?

Example: $f(n) = 3n + 100$ is $O(n)$

Proof. Take $n \geq 1$. Then

$$|3n + 100| \leq |3n + 100n| = 103|n|.$$

(To satisfy the definition of big O, we take $b = 1, B = 103$.)

Example: $f(n) = 9n^2 + 3n - 2$ is $O(n^2)$.

Proof. Take $n \geq 1$. Then

$$|9n^2 + 3n - 2| \leq |9n^2 + 3n^2 + 2n^2| = 14|n^2|.$$

(To satisfy the definition of big O, we take $b = 1, B = 14$.)

Remark. Big O notation hides irrelevant details.


Poll. Do you see a pattern?

Solution. Polynomials are big O of their highest-order term.


Remark. ...and also big O of still higher-order terms. E.g. for $n \geq 1$,

$$14|n^2| \leq 14|n^3| \leq 14|n^4| \leq \dots$$

Classification Theorem: Zooming In On Polynomials




Classification Theorem: Zooming In On Polynomials




A taxonomy of growth rates: As you go down the legend, each function is big O of everything above it, but **not** big O of everything below it.

Classification Theorem: Zooming In On Polynomials



A taxonomy of growth rates: As you go down the legend, each function is big O of everything above it, but **not** big O of everything below it.

Classification Theorem: Zooming In On Polynomials



A taxonomy of growth rates: As you go down the legend, each function is big O of everything above it, but **not** big O of everything below it.

Theorem

For any function f and positive real numbers u and v with $u < v$,

$$f \text{ is } O(x^u) \implies f \text{ is } O(x^v)$$

A major problem with Big O statements

Poll. Suppose a person is analyzing the efficiency of algorithms, and finds that f is $O(x^5)$ and g is $O(x^4)$. Because $4 < 5$, this person concludes that g has a better algorithmic efficiency than f. Is this person's conclusion correct?

A major problem with Big O statements

Poll. Suppose a person is analyzing the efficiency of algorithms, and finds that f is $O(x^5)$ and g is $O(x^4)$. Because $4 < 5$, this person concludes that g has a better algorithmic efficiency than f. Is this person's conclusion correct?

Solution. No. For instance, take $f(x) = x^2$ and $g(x) = x^3$. Then

f is $O(x^2), O(x^3), O(x^4), O(x^5), \ldots$

and

g is $O(x^3), O(x^4), O(x^5), \ldots$

A major problem with Big O statements

Poll. Suppose a person is analyzing the efficiency of algorithms, and finds that f is $O(x^5)$ and g is $O(x^4)$. Because $4 < 5$, this person concludes that g has a better algorithmic efficiency than f. Is this person's conclusion correct?

Solution. No. For instance, take $f(x) = x^2$ and $g(x) = x^3$. Then

f is $O(x^2), O(x^3), O(x^4), O(x^5), \ldots$

and

g is $O(x^3), O(x^4), O(x^5), \ldots$

A major problem with Big O statements

Poll. Suppose a person is analyzing the efficiency of algorithms, and finds that f is $O(x^5)$ and g is $O(x^4)$. Because $4 < 5$, this person concludes that g has a better algorithmic efficiency than f. Is this person's conclusion correct?

Solution. No. For instance, take $f(x) = x^2$ and $g(x) = x^3$. Then

f is $O(x^2), O(x^3), O(x^4), O(x^5), \ldots$

and

g is $O(x^3), O(x^4), O(x^5), \ldots$

Remark. The problem is that the upper bounds for Big O can be needlessly large!

A major problem with Big O statements

Poll. Suppose a person is analyzing the efficiency of algorithms, and finds that f is $O(x^5)$ and g is $O(x^4)$. Because $4 < 5$, this person concludes that g has a better algorithmic efficiency than f. Is this person's conclusion correct?

Solution. No. For instance, take $f(x) = x^2$ and $g(x) = x^3$. Then

f is $O(x^2), O(x^3), O(x^4), O(x^5), \ldots$

and

g is $O(x^3), O(x^4), O(x^5), \ldots$

Remark. The problem is that the upper bounds for Big O can be needlessly large!

We should interpret "g is $O(x^4)$" as: "g is of order **at most** x^4".

How to compare the efficiency of algorithms?



Figure 2: American computer scientist and mathematician Donald Knuth

How to compare the efficiency of algorithms?







Figure 2: American computer scientist and mathematician Donald Knuth

Overview: Big Theta combines Big O and Big Omega

Big O	f is $O(g)$	f is of order at most g	The values of f are eventually **less** than those of a positive multiple of g.
Big Omega	f is $\Omega(g)$	f is of order at least g	The values of f are eventually **greater** than those of a positive multiple of g.
Big Theta	f is $\Theta(g)$	f is of order g	The values of f are eventually **less** than those of a positive multiple of g and **greater** than those of a positive multiple of g.


$f(n)$ is $\Omega(g(n))$
Graph of f
$(n, f(n))$
Graph of Ag
$(n, Ag(n))$
a
n


$f(n)$ is $O(g(n))$
Graph of Bg
$(n, Bg(n))$
Graph of f
$(n, f(n))$
b
n



$f(n)$ is $\Theta(g(n))$
Graph of Bg
$(n, Bg(n))$
Graph of f
$(n, f(n))$
Graph of Ag
$(n, Ag(n))$
k
n

Big Omega: Definition

Definition

Let f and g be real-valued non-negative functions defined on the same set of nonnegative integers.
Then f is of order at least g, written $\boldsymbol{f(n)}$ **is** $\boldsymbol{\Omega(g(n))}$ (f of n is big-Omega of g of n) if and only if there exist positive real number A and integer a such that


$$|f(n)| \geq A|g(n)| \quad \text{for every integer } n \geq a$$



Big Theta: Definition

Definition

Let f and g be real-valued non-negative functions defined on the same set of nonnegative integers.
Then f is of order g, written $\boldsymbol{f(n)}$ **is** $\boldsymbol{\Theta(g(n))}$ (f of n is big-Theta of g of n) if and only if $\boldsymbol{f(n)}$ is $\boldsymbol{O(g(n))}$ and $\boldsymbol{f(n)}$ is $\boldsymbol{\Omega(g(n))}$.



A useful theorem on Big Theta

Theorem: On Polynomial Orders

If m is any integer with $m \geq 0$ and $c_0, c_1, c_2, \ldots c_m$ are real numbers with $c_m \geq 0$, then

$$c_m x^m + c_{m-1} x^{m-1} + \ldots + c_1 x + c_0 \text{ is } \Theta(x^m).$$

Group exercises

Random group assignments

Ashley, Isabelle: 7
Bailey, Tanna: 3
Blaisdell, Olyver: 7
Brown, Ashton: 10
Cardenas, Dominic: 9
Cutler, George: 12
Fabik, Roland: 6
Fellows, Audrey: 9
Fitzgerald, Ryan: 11
Fuller, Ryan: 3
Griffen, Clara: 1
Grutsch, Tanner: 2
Hager, Nathan: 8
Horner, Jayden: 10
Johnson, Jaret: 5
Kintzel, Samantha: 6
Lameres, Alexis: 2
Lee, Holden: 4
Mayala, Reece: 1
McLean, Connor: 5
Mcbride, Daniel: 1
Miller, Maverick: 9
Moss, Emily: 8
Mousseau, Matthew: 4
Neuman, Evan: 4
Putnam, John: 5
Ramos, Vydel: 12
Rathbun, Jedidiah: 6
Roberts, Matthew: 11
Roller, Nathan: 7
Ryan, Otis: 8
Shevtsov, Timothy: 2
Stensland, Emma: 11
Thiessen, Logan: 3
Thompson, Jaiden: 12
Vath, Ethan: 10

Group exercises

Exercises.

1. Let $a_n = 1,000,000n + 3,000,000$. Prove that $a_n = O(n)$.
2. Let $a_n = 5 + \frac{1}{n} + \frac{1}{n+1}$. Prove that $a_n = O(1)$.
3. Let $a_n = n^2 + n + 1 + \frac{1}{n} + \sin n$. Prove that $a_n = O(n^2)$.
4. Let $a_n = 3n^2 + 7$. Prove that $a_n = \Theta(n^2)$.
5. Prove Theorem 7.9 from the Ponomarenko textbook.

Note. Please solve all the problems above using the definitions of Big-O and Big-Θ.

Theorem 7.9 (Ponomarenko). Let a_n be a sequence, and b_n, c_n be test sequences. If $a_n = O(b_n)$ and $b_n = O(c_n)$, then $a_n = O(c_n)$.

Solution to Group Exercise #1

Problem. Let $a_n = 1,000,000n + 3,000,000$. Prove that $a_n = O(n)$.

Solution. For $n \geq 1$,

$$|1,000,000n + 3,000,000| \leq |1,000,000n + 3,000,000n| = 4,000,000|n|$$

This satisfies the definition that a_n is $O(n)$ by setting $b = 1$ and $B = 4,000,000$.

Remark. The notation (b and B) comes from the definition of Big O given in this slide deck.

Solution to Group Exercise #2

Problem. Let $a_n = 5 + \frac{1}{n} + \frac{1}{n+1}$. Prove that $a_n = O(1)$.

Solution. For $n \geq 1$,

$$|5 + \frac{1}{n} + \frac{1}{n+1}| \leq |5 + 1 + 1| = 7 \cdot |1|$$

This satisfies the definition that a_n is $O(1)$ by setting $b = 1$ and $B = 7$.

Remark. The notation (b and B) comes from the definition of Big O given in this slide deck.

Solution to Group Exercise #3

Problem. Let $a_n = n^2 + n + 1 + \frac{1}{n} + \sin n$. Prove that $a_n = O(n^2)$.

Solution. For $n \geq 1$,

$$\begin{aligned} |n^2 + n + 1 + \frac{1}{n} + \sin n| &\leq |n^2 + n + 1 + 1 + 1| \\ &\leq |n^2 + n^2 + n^2 + n^2 + n^2| \\ &= 5|n^2| \end{aligned}$$

This satisfies the definition that a_n is $O(n^2)$ by setting $b = 1$ and $B = 5$.

Remark. The notation (b and B) comes from the definition of Big O given in this slide deck.

Solution to Group Exercise #4

Problem. Let $a_n = 3n^2 + 7$. Prove that $a_n = \Theta(n^2)$.

Solution. For $n \geq 1$,

$$|3n^2 + 7| \leq |3n^2 + 7n^2| = 10|n^2|$$

This satisfies the definition that a_n is $O(n^2)$ by setting $b = 1$ and $B = 10$.

Again for $n \geq 1$,

$$|3n^2 + 7| \geq |3n^2| = 3|n^2|$$

This satisfies the definition that a_n is $\Omega(n^2)$ by setting $a = 1$ and $A = 3$.

Since a_n is both $O(n^2)$ and $\Omega(n^2)$, we can conclude that a_n is $\Theta(n^2)$.

Remark. The notation (a, b, A, B) comes from the definitions of Big O and Big Theta given in this slide deck.

Solution to Group Exercise #5

Problem. Prove Theorem 7.9 from the Ponomarenko textbook.

Theorem 7.9 (Ponomarenko). Let a_n be a sequence, and b_n, c_n be test sequences. If $a_n = O(b_n)$ and $b_n = O(c_n)$, then $a_n = O(c_n)$.

Solution. Since $a_n = O(b_n)$, we have that for $n \geq b_1$,

$$|a_n| \leq B_1|b_n| \qquad \text{for some constant } B_1. \tag{1}$$

Since $b_n = O(c_n)$, we have that for $n \geq b_2$,

$$|b_n| \leq B_2|c_n| \qquad \text{for some constant } B_2. \tag{2}$$

Hence for $n \geq \max\{b_1, b_2\}$,

$$|a_n| \overset{\text{Eq. (1)}}{\leq} B_1|b_n| \overset{\text{Eq. (2)}}{\leq} B_1 \cdot B_2|c_n|$$

This satisfies the definition that a_n is $O(c_n)$ by setting $b = \max\{b_1, b_2\}$ and $B = B_1 \cdot B_2$.