10/12/2026: Smallest Counterexample

CSCI 246: Discrete Structures

Textbook reference: Sec 21, Scheinerman

Today's Agenda

- ($\approx$ 30 min) Mini-lecture
- ($\approx$ 20 min) Group exercises

The Well-Ordering Principle

In today's class, we'll use the following **axiom**.

The Well-Ordering Principle

In today's class, we'll use the following **axiom**.

Axiom: The Well Ordering Principle

Every non-empty set of natural numbers contains a least element.

The Well-Ordering Principle

In today's class, we'll use the following **axiom**.

Axiom: The Well Ordering Principle

Every non-empty set of natural numbers contains a least element.

Seems pretty intuitive, right!?

The Well-Ordering Principle

In today's class, we'll use the following **axiom**.

Axiom: The Well Ordering Principle

Every non-empty set of natural numbers contains a least element.

Seems pretty intuitive, right!?

Example 1

Let

$$S = \{n \in \mathbb{N} : n^2 > 50\}.$$

This set is a nonempty subset of the natural numbers. By the well-ordering principle, S contains a least element: ???

The Well-Ordering Principle

In today's class, we'll use the following **axiom**.

Axiom: The Well Ordering Principle

Every non-empty set of natural numbers contains a least element.

Seems pretty intuitive, right!?

Example 1

Let

$$S = \{n \in \mathbb{N} : n^2 > 50\}.$$

This set is a nonempty subset of the natural numbers. By the well-ordering principle, S contains a least element: **8.**

The Well-Ordering Principle

In today's class, we'll use the following **axiom**.

Axiom: The Well Ordering Principle

Every non-empty set of natural numbers contains a least element.

Seems pretty intuitive, right!?

Example 2

Let

$$P = \{x \in \mathbb{N} : x \text{ is prime}\}.$$

This set is a nonempty subset of the natural numbers. By the well-ordering principle, P contains a least element: ???.

The Well-Ordering Principle

In today's class, we'll use the following **axiom**.

Axiom: The Well Ordering Principle

Every non-empty set of natural numbers contains a least element.

Seems pretty intuitive, right!?

Example 2

Let

$$P = \{x \in \mathbb{N} : x \text{ is prime}\}.$$

This set is a nonempty subset of the natural numbers. By the well-ordering principle, P contains a least element: **2.**

Smallest counterexample: the idea

Smallest counterexample is new proof technique.

Smallest counterexample: the idea

Smallest counterexample is new proof technique. Here we prove a claim using proof by contradiction.

Smallest counterexample: the idea

Smallest counterexample is new proof technique. Here we prove a claim using proof by contradiction.

At a high level, the argument goes like this:

- Notice: If there exists a counterexample, there must be a minimal counterexample.

Smallest counterexample: the idea

Smallest counterexample is new proof technique. Here we prove a claim using proof by contradiction.

At a high level, the argument goes like this:

- Notice: If there exists a counterexample, there must be a minimal counterexample.
- Show: The minimal counterexample produces a contradiction.

Smallest counterexample: the idea

Smallest counterexample is new proof technique. Here we prove a claim using proof by contradiction.

At a high level, the argument goes like this:

- Notice: If there exists a counterexample, there must be a minimal counterexample.
- Show: The minimal counterexample produces a contradiction.
- Conclude: The claim holds!

Proof by smallest counterexample: A data science example

Setting. Data often arrives as a **stream** (sensor readings, web logs, a file too big for memory). We can't store everything and then average, so we keep a **running mean**, updating it as each point x_n arrives:

$$\mu_1 = x_1, \qquad \mu_n = \mu_{n-1} + \frac{x_n - \mu_{n-1}}{n} \quad \text{for } n \geq 2.$$

Proof by smallest counterexample: A data science example

Setting. Data often arrives as a **stream** (sensor readings, web logs, a file too big for memory). We can't store everything and then average, so we keep a **running mean**, updating it as each point x_n arrives:

$$\mu_1 = x_1, \qquad \mu_n = \mu_{n-1} + \frac{x_n - \mu_{n-1}}{n} \quad \text{for } n \geq 2.$$

Claim.

For every positive integer n, the running mean equals the true mean:

$$\mu_n = \frac{x_1 + x_2 + \cdots + x_n}{n}.$$

Proof by smallest counterexample: A data science example

Setting. Data often arrives as a **stream** (sensor readings, web logs, a file too big for memory). We can't store everything and then average, so we keep a **running mean**, updating it as each point x_n arrives:

$$\mu_1 = x_1, \qquad \mu_n = \mu_{n-1} + \frac{x_n - \mu_{n-1}}{n} \quad \text{for } n \geq 2.$$

Claim.

For every positive integer n, the running mean equals the true mean:

$$\mu_n = \frac{x_1 + x_2 + \cdots + x_n}{n}.$$

Proof. Suppose not. Then the set of n for which the claim fails is a nonempty set of positive integers, so by the Well-Ordering Principle it has a **smallest element** s.

Proof by smallest counterexample: A data science example

Setting. Data often arrives as a **stream** (sensor readings, web logs, a file too big for memory). We can't store everything and then average, so we keep a **running mean**, updating it as each point x_n arrives:

$$\mu_1 = x_1, \qquad \mu_n = \mu_{n-1} + \frac{x_n - \mu_{n-1}}{n} \quad \text{for } n \geq 2.$$

Claim.

For every positive integer n, the running mean equals the true mean:

$$\mu_n = \frac{x_1 + x_2 + \cdots + x_n}{n}.$$

Proof. Suppose not. Then the set of n for which the claim fails is a nonempty set of positive integers, so by the Well-Ordering Principle it has a **smallest element** s.

- $s \neq 1$, since $\mu_1 = x_1 = \frac{x_1}{1}$.

Proof by smallest counterexample: A data science example

Setting. Data often arrives as a **stream** (sensor readings, web logs, a file too big for memory). We can't store everything and then average, so we keep a **running mean**, updating it as each point x_n arrives:

$$\mu_1 = x_1, \qquad \mu_n = \mu_{n-1} + \frac{x_n - \mu_{n-1}}{n} \quad \text{for } n \geq 2.$$

Claim.

For every positive integer n, the running mean equals the true mean:

$$\mu_n = \frac{x_1 + x_2 + \cdots + x_n}{n}.$$

Proof. Suppose not. Then the set of n for which the claim fails is a nonempty set of positive integers, so by the Well-Ordering Principle it has a **smallest element** s.

- $s \neq 1$, since $\mu_1 = x_1 = \frac{x_1}{1}$.
- Since $1 \leq s - 1 < s$, the claim **holds** for $s - 1$: $\quad (s-1)\,\mu_{s-1} = x_1 + \cdots + x_{s-1}$.

Proof by smallest counterexample: A data science example

Setting. Data often arrives as a **stream** (sensor readings, web logs, a file too big for memory). We can't store everything and then average, so we keep a **running mean**, updating it as each point x_n arrives:

$$\mu_1 = x_1, \qquad \mu_n = \mu_{n-1} + \frac{x_n - \mu_{n-1}}{n} \quad \text{for } n \geq 2.$$

Claim.

For every positive integer n, the running mean equals the true mean:

$$\mu_n = \frac{x_1 + x_2 + \cdots + x_n}{n}.$$

Proof. Suppose not. Then the set of n for which the claim fails is a nonempty set of positive integers, so by the Well-Ordering Principle it has a **smallest element** s.

- $s \neq 1$, since $\mu_1 = x_1 = \frac{x_1}{1}$.
- Since $1 \leq s - 1 < s$, the claim **holds** for $s - 1$: $\quad (s-1)\,\mu_{s-1} = x_1 + \cdots + x_{s-1}$.
- So

$$\mu_s = \mu_{s-1} + \frac{x_s - \mu_{s-1}}{s} = \frac{(s-1)\,\mu_{s-1} + x_s}{s} = \frac{x_1 + \cdots + x_{s-1} + x_s}{s},$$

so the claim holds for s, which contradicts s being a counterexample. ■

Proof by smallest counterexample: A data science example

Setting. Data often arrives as a **stream** (sensor readings, web logs, a file too big for memory). We can't store everything and then average, so we keep a **running mean**, updating it as each point x_n arrives:

$$\mu_1 = x_1, \qquad \mu_n = \mu_{n-1} + \frac{x_n - \mu_{n-1}}{n} \quad \text{for } n \geq 2.$$

Claim.

For every positive integer n, the running mean equals the true mean:

$$\mu_n = \frac{x_1 + x_2 + \cdots + x_n}{n}.$$

Proof. Suppose not. Then the set of n for which the claim fails is a nonempty set of positive integers, so by the Well-Ordering Principle it has a **smallest element** s.

- $s \neq 1$, since $\mu_1 = x_1 = \frac{x_1}{1}$.
- Since $1 \leq s - 1 < s$, the claim **holds** for $s - 1$: $\quad (s-1)\,\mu_{s-1} = x_1 + \cdots + x_{s-1}$.
- So

$$\mu_s = \mu_{s-1} + \frac{x_s - \mu_{s-1}}{s} = \frac{(s-1)\,\mu_{s-1} + x_s}{s} = \frac{x_1 + \cdots + x_{s-1} + x_s}{s},$$

  so the claim holds for s, which contradicts s being a counterexample. ■

Why care? We've proven a one-pass streaming algorithm correct for *every* input length. ✓

The most famous proof by smallest counterexample

Prime Factorization Theorem (existence part of the Fundamental Theorem of Arithmetic)

Every natural number $n \geq 2$ can be written as a product of primes.

The most famous proof by smallest counterexample

Prime Factorization Theorem (existence part of the Fundamental Theorem of Arithmetic)

Every natural number $n \geq 2$ can be written as a product of primes.

Examples. $21 = 3 \times 7$, $100 = 2 \times 2 \times 5 \times 5$, $13 = 13$ (a product of *one* prime).

The most famous proof by smallest counterexample

Prime Factorization Theorem (existence part of the Fundamental Theorem of Arithmetic)

Every natural number $n \geq 2$ can be written as a product of primes.

Examples. $21 = 3 \times 7$, $100 = 2 \times 2 \times 5 \times 5$, $13 = 13$ (a product of *one* prime).

Why should we care?

The most famous proof by smallest counterexample

Prime Factorization Theorem (existence part of the Fundamental Theorem of Arithmetic)

Every natural number $n \geq 2$ can be written as a product of primes.

Examples. $21 = 3 \times 7$, $100 = 2 \times 2 \times 5 \times 5$, $13 = 13$ (a product of *one* prime).

Why should we care? Internet Security! Your bank account is safe because every large number has a prime factorization, but nobody can find it.

The most famous proof by smallest counterexample

Prime Factorization Theorem (existence part of the Fundamental Theorem of Arithmetic)

Every natural number $n \geq 2$ can be written as a product of primes.

Examples. $21 = 3 \times 7$, $100 = 2 \times 2 \times 5 \times 5$, $13 = 13$ (a product of *one* prime).

Proof. Suppose not. Then the set of exceptions

$$X = \{n \in \mathbb{N} : n \geq 2 \text{ and } n \text{ is } \textbf{not} \text{ a product of primes}\}$$

is nonempty, so by the Well-Ordering Principle it has a **smallest element** s.

The most famous proof by smallest counterexample

Prime Factorization Theorem (existence part of the Fundamental Theorem of Arithmetic)

Every natural number $n \geq 2$ can be written as a product of primes.

Examples. $21 = 3 \times 7$, $100 = 2 \times 2 \times 5 \times 5$, $13 = 13$ (a product of *one* prime).

Proof. Suppose not. Then the set of exceptions

$$X = \{n \in \mathbb{N} : n \geq 2 \text{ and } n \text{ is } \textbf{not} \text{ a product of primes}\}$$

is nonempty, so by the Well-Ordering Principle it has a **smallest element** s.

- Why can't s be prime?

The most famous proof by smallest counterexample

Prime Factorization Theorem (existence part of the Fundamental Theorem of Arithmetic)

Every natural number $n \geq 2$ can be written as a product of primes.

Examples. $21 = 3 \times 7$, $100 = 2 \times 2 \times 5 \times 5$, $13 = 13$ (a product of *one* prime).

Proof. Suppose not. Then the set of exceptions

$$X = \{n \in \mathbb{N} : n \geq 2 \text{ and } n \text{ is } \textbf{not} \text{ a product of primes}\}$$

is nonempty, so by the Well-Ordering Principle it has a **smallest element** s.

- Why can't s be prime? A prime is a product of one prime. So s is not prime.

The most famous proof by smallest counterexample

Prime Factorization Theorem (existence part of the Fundamental Theorem of Arithmetic)

Every natural number $n \geq 2$ can be written as a product of primes.

Examples. $21 = 3 \times 7$, $100 = 2 \times 2 \times 5 \times 5$, $13 = 13$ (a product of *one* prime).

Proof. Suppose not. Then the set of exceptions

$$X = \{n \in \mathbb{N} : n \geq 2 \text{ and } n \text{ is } \textbf{not} \text{ a product of primes}\}$$

is nonempty, so by the Well-Ordering Principle it has a **smallest element** s.

- Why can't s be prime? A prime is a product of one prime. So s is not prime. Thus s is **composite** (recalling that $s \geq 2$).

The most famous proof by smallest counterexample

Prime Factorization Theorem (existence part of the Fundamental Theorem of Arithmetic)

Every natural number $n \geq 2$ can be written as a product of primes.

Examples. $21 = 3 \times 7$, $100 = 2 \times 2 \times 5 \times 5$, $13 = 13$ (a product of *one* prime).

Proof. Suppose not. Then the set of exceptions

$$X = \{n \in \mathbb{N} : n \geq 2 \text{ and } n \text{ is } \textbf{not} \text{ a product of primes}\}$$

is nonempty, so by the Well-Ordering Principle it has a **smallest element** s.

- Why can't s be prime? A prime is a product of one prime. So s is not prime. Thus s is **composite** (recalling that $s \geq 2$).
- What does that mean?

The most famous proof by smallest counterexample

Prime Factorization Theorem (existence part of the Fundamental Theorem of Arithmetic)

Every natural number $n \geq 2$ can be written as a product of primes.

Examples. $21 = 3 \times 7$, $100 = 2 \times 2 \times 5 \times 5$, $13 = 13$ (a product of *one* prime).

Proof. Suppose not. Then the set of exceptions

$$X = \{n \in \mathbb{N} : n \geq 2 \text{ and } n \text{ is } \textbf{not} \text{ a product of primes}\}$$

is nonempty, so by the Well-Ordering Principle it has a **smallest element** s.

- Why can't s be prime? A prime is a product of one prime. So s is not prime. Thus s is **composite** (recalling that $s \geq 2$).
- Hence $s = ab$ for some integers a, b with $1 < a, b < s$.

The most famous proof by smallest counterexample

Prime Factorization Theorem (existence part of the Fundamental Theorem of Arithmetic)

Every natural number $n \geq 2$ can be written as a product of primes.

Examples. $21 = 3 \times 7$, $100 = 2 \times 2 \times 5 \times 5$, $13 = 13$ (a product of *one* prime).

Proof. Suppose not. Then the set of exceptions

$$X = \{n \in \mathbb{N} : n \geq 2 \text{ and } n \text{ is } \textbf{not} \text{ a product of primes}\}$$

is nonempty, so by the Well-Ordering Principle it has a **smallest element** s.

- Why can't s be prime? A prime is a product of one prime. So s is not prime. Thus s is **composite** (recalling that $s \geq 2$).
- Hence $s = ab$ for some integers a, b with $1 < a, b < s$.
- What do we know about a, b?

The most famous proof by smallest counterexample

Prime Factorization Theorem (existence part of the Fundamental Theorem of Arithmetic)

Every natural number $n \geq 2$ can be written as a product of primes.

Examples. $21 = 3 \times 7$, $100 = 2 \times 2 \times 5 \times 5$, $13 = 13$ (a product of *one* prime).

Proof. Suppose not. Then the set of exceptions

$$X = \{n \in \mathbb{N} : n \geq 2 \text{ and } n \text{ is } \textbf{not} \text{ a product of primes}\}$$

is nonempty, so by the Well-Ordering Principle it has a **smallest element** s.

- Why can't s be prime? A prime is a product of one prime. So s is not prime. Thus s is **composite** (recalling that $s \geq 2$).
- Hence $s = ab$ for some integers a, b with $1 < a, b < s$.
- Since s is the smallest element of X, neither a nor b is in X. So each is a product of primes.

The most famous proof by smallest counterexample

Prime Factorization Theorem (existence part of the Fundamental Theorem of Arithmetic)

Every natural number $n \geq 2$ can be written as a product of primes.

Examples. $21 = 3 \times 7$, $100 = 2 \times 2 \times 5 \times 5$, $13 = 13$ (a product of *one* prime).

Proof. Suppose not. Then the set of exceptions

$$X = \{n \in \mathbb{N} : n \geq 2 \text{ and } n \text{ is } \textbf{not} \text{ a product of primes}\}$$

is nonempty, so by the Well-Ordering Principle it has a **smallest element** s.

- Why can't s be prime? A prime is a product of one prime. So s is not prime. Thus s is **composite** (recalling that $s \geq 2$).
- Hence $s = ab$ for some integers a, b with $1 < a, b < s$.
- Since s is the smallest element of X, neither a nor b is in X. So each is a product of primes.
- So what?

The most famous proof by smallest counterexample

Prime Factorization Theorem (existence part of the Fundamental Theorem of Arithmetic)

Every natural number $n \geq 2$ can be written as a product of primes.

Examples. $21 = 3 \times 7$, $100 = 2 \times 2 \times 5 \times 5$, $13 = 13$ (a product of *one* prime).

Proof. Suppose not. Then the set of exceptions

$$X = \{n \in \mathbb{N} : n \geq 2 \text{ and } n \text{ is } \textbf{not} \text{ a product of primes}\}$$

is nonempty, so by the Well-Ordering Principle it has a **smallest element** s.

- Why can't s be prime? A prime is a product of one prime. So s is not prime. Thus s is **composite** (recalling that $s \geq 2$).
- Hence $s = ab$ for some integers a, b with $1 < a, b < s$.
- Since s is the smallest element of X, neither a nor b is in X. So each is a product of primes.
- Then $s = ab$ is a product of primes, contradicting $s \in X$. ■

Proof by smallest counterexample: Binary representation

Statement. *Every positive integer can be written as a sum of* **distinct** *powers of 2.*

Proof by smallest counterexample: Binary representation

Statement. *Every positive integer can be written as a sum of* **distinct** *powers of 2.*
That is, for any positive integer n,

$$n = 2^{a_1} + 2^{a_2} + \cdots + 2^{a_k}$$

for some nonnegative integers $a_1 > a_2 > \cdots > a_k$.

Proof by smallest counterexample: Binary representation

Statement. *Every positive integer can be written as a sum of* **distinct** *powers of 2.* That is, for any positive integer n,

$$n = 2^{a_1} + 2^{a_2} + \cdots + 2^{a_k}$$

for some nonnegative integers $a_1 > a_2 > \cdots > a_k$.

Remark. This is the statement that "every positive integer has a binary representation." For example,

$$13 = 8 + 4 + 1 = 2^3 + 2^2 + 2^0 = 1101_{\text{two}}.$$

Proof by smallest counterexample: Binary representation

Statement. *Every positive integer can be written as a sum of* **distinct** *powers of 2.* That is, for any positive integer n,

$$n = 2^{a_1} + 2^{a_2} + \cdots + 2^{a_k}$$

for some nonnegative integers $a_1 > a_2 > \cdots > a_k$.

Proof (using the Well-Ordering Principle). Suppose, for contradiction, that the statement is **false**. Then the set X of positive integers that **cannot** be written as a sum of distinct powers of 2 is nonempty. By the Well-Ordering Principle, X has a **smallest element**; call it s.

Proof by smallest counterexample: Binary representation

Statement. *Every positive integer can be written as a sum of* **distinct** *powers of 2.*
That is, for any positive integer n,

$$n = 2^{a_1} + 2^{a_2} + \cdots + 2^{a_k}$$

for some nonnegative integers $a_1 > a_2 > \cdots > a_k$.

Proof (using the Well-Ordering Principle). Suppose, for contradiction, that the statement is **false**. Then the set X of positive integers that **cannot** be written as a sum of distinct powers of 2 is nonempty. By the Well-Ordering Principle, X has a **smallest element**; call it s.

So every positive integer less than s **can** be written this way, but s itself **cannot**.

Proof by smallest counterexample: Binary representation

Statement. *Every positive integer can be written as a sum of* **distinct** *powers of 2.*
That is, for any positive integer n,

$$n = 2^{a_1} + 2^{a_2} + \cdots + 2^{a_k}$$

for some nonnegative integers $a_1 > a_2 > \cdots > a_k$.

Proof (using the Well-Ordering Principle). Suppose, for contradiction, that the statement is **false**. Then the set X of positive integers that **cannot** be written as a sum of distinct powers of 2 is nonempty. By the Well-Ordering Principle, X has a **smallest element**; call it s.

So every positive integer less than s **can** be written this way, but s itself **cannot**.

Note that s is not itself a power of 2 (a single power of 2 is such a sum), so $2^k < s < 2^{k+1}$ for some k.

Proof by smallest counterexample: Binary representation

Statement. *Every positive integer can be written as a sum of* **distinct** *powers of 2.* That is, for any positive integer n,

$$n = 2^{a_1} + 2^{a_2} + \cdots + 2^{a_k}$$

for some nonnegative integers $a_1 > a_2 > \cdots > a_k$.

Proof (using the Well-Ordering Principle). Suppose, for contradiction, that the statement is **false**. Then the set X of positive integers that **cannot** be written as a sum of distinct powers of 2 is nonempty. By the Well-Ordering Principle, X has a **smallest element**; call it s.

So every positive integer less than s **can** be written this way, but s itself **cannot**.

Note that s is not itself a power of 2 (a single power of 2 is such a sum), so $2^k < s < 2^{k+1}$ for some k. Now define $r = s - 2^k$. Then $1 \leq r < 2^k$, since $s < 2^{k+1} = 2^k + 2^k$.

Proof by smallest counterexample: Binary representation

Statement. *Every positive integer can be written as a sum of* **distinct** *powers of 2.* That is, for any positive integer n,

$$n = 2^{a_1} + 2^{a_2} + \cdots + 2^{a_k}$$

for some nonnegative integers $a_1 > a_2 > \cdots > a_k$.

Proof (using the Well-Ordering Principle). Suppose, for contradiction, that the statement is **false**. Then the set X of positive integers that **cannot** be written as a sum of distinct powers of 2 is nonempty. By the Well-Ordering Principle, X has a **smallest element**; call it s.

So every positive integer less than s **can** be written this way, but s itself **cannot**.

Note that s is not itself a power of 2 (a single power of 2 is such a sum), so $2^k < s < 2^{k+1}$ for some k. Now define $r = s - 2^k$. Then $1 \leq r < 2^k$, since $s < 2^{k+1} = 2^k + 2^k$.

Since $1 \leq r < s$, by minimality of s, r **can** be written as a sum of distinct powers of 2, each less than 2^k (since $r < 2^k$). So $s = 2^k + (\text{distinct powers of 2, each less than } 2^k)$ is a sum of distinct powers of 2, which is a contradiction!

Proof by smallest counterexample: Binary representation

Statement. *Every positive integer can be written as a sum of* **distinct** *powers of 2.* That is, for any positive integer n,

$$n = 2^{a_1} + 2^{a_2} + \cdots + 2^{a_k}$$

for some nonnegative integers $a_1 > a_2 > \cdots > a_k$.

Proof (using the Well-Ordering Principle). Suppose, for contradiction, that the statement is **false**. Then the set X of positive integers that **cannot** be written as a sum of distinct powers of 2 is nonempty. By the Well-Ordering Principle, X has a **smallest element**; call it s.

So every positive integer less than s **can** be written this way, but s itself **cannot**.

Note that s is not itself a power of 2 (a single power of 2 is such a sum), so $2^k < s < 2^{k+1}$ for some k. Now define $r = s - 2^k$. Then $1 \leq r < 2^k$, since $s < 2^{k+1} = 2^k + 2^k$.

Since $1 \leq r < s$, by minimality of s, r **can** be written as a sum of distinct powers of 2, each less than 2^k (since $r < 2^k$). So $s = 2^k + (\text{distinct powers of 2, each less than } 2^k)$ is a sum of distinct powers of 2, which is a contradiction!

Therefore X is empty, and the statement holds. ✓

Proof by the Well-Ordering Principle

Scheinerman's Proof Template 16.

Proof by the Well-Ordering Principle.

To prove a statement about natural numbers:

Proof. Suppose, for the sake of contradiction, that the statement is false. Let $X \subseteq \mathbb{N}$ be the set of counterexamples to the statement. (I like the letter X for eXceptions.) Since we have supposed the statement is false, $X \neq \emptyset$. By the Well-Ordering Principle, X contains a least element, x.

(Basis step.) We know that $x \neq 0$ because *show that the result holds for* 0*; this is usually easy.*

Consider $x - 1$. Since $x > 0$, we know that $x - 1 \in \mathbb{N}$ and the statement is true for $x - 1$ (because $x - 1 < x$). *From here we argue to a contradiction, often that x both is and is not a counterexample to the statement.*$\Rightarrow\Leftarrow$ ■

Proof by the Well-Ordering Principle

Scheinerman's Proof Template 16.

Proof by the Well-Ordering Principle.

To prove a statement about natural numbers:

Proof. Suppose, for the sake of contradiction, that the statement is false. Let $X \subseteq \mathbb{N}$ be the set of counterexamples to the statement. (I like the letter X for eXceptions.) Since we have supposed the statement is false, $X \neq \emptyset$. By the Well-Ordering Principle, X contains a least element, x.

(Basis step.) We know that $x \neq 0$ because *show that the result holds for* 0*; this is usually easy.*

Consider $x - 1$. Since $x > 0$, we know that $x - 1 \in \mathbb{N}$ and the statement is true for $x - 1$ (because $x - 1 < x$). *From here we argue to a contradiction, often that x both is and is not a counterexample to the statement.*$\Rightarrow\Leftarrow$ ■

Remarks.

- The basis step could be a different number than 0 (e.g. 1) if the statement is about a subset of the natural numbers (e.g. the positive integers).
- The contradiction needn't come from considering $x - 1$.

Group exercises

Ashley, Isabelle: 1
Bailey, Tanna: 5
Blaisdell, Olyver: 11
Brown, Ashton: 11
Brown, Jonathan: 6
Cardenas, Dominic: 12
Cutler, George: 2
Fabik, Roland: 9
Fellows, Audrey: 2
Fitzgerald, Ryan: 12
Fuller, Ryan: 9
Griffen, Clara: 4
Grutsch, Tanner: 11
Hager, Nathan: 6
Horner, Jayden: 3
Johnson, Jaret: 3
Kintzel, Samantha: 2
Lameres, Alexis: 13
Lee, Holden: 5
Mayala, Reece: 10
McLean, Connor: 7
Mcbride, Daniel: 10
Miller, Maverick: 8
Moss, Emily: 3
Mousseau, Matthew: 8
Neuman, Evan: 13
Putnam, John: 8
Ramos, Vydel: 7
Rathbun, Jedidiah: 1
Roberts, Matthew: 9
Roller, Nathan: 7
Ryan, Otis: 4
Shevtsov, Timothy: 6
Sinclair, Aidan: 5
Stensland, Emma: 12
Thiessen, Logan: 4
Thompson, Jaiden: 1
Vath, Ethan: 10

Group exercises

1. Use proof by smallest counterexample to show that $n < 2^n$ for all $n \in \mathbb{N}$.
2. Use proof by smallest counterexample to show that

$$1 + 2 + 3 + \cdots + n = \frac{1}{2}n^2 + \frac{1}{2}n$$

for all positive integers n.

Solution to group exercise #1

Problem. Use proof by smallest counterexample to show that $n < 2^n$ for all $n \in \mathbb{N}$.

Solution. Suppose, for the sake of contradiction, that the statement is false. Let $X \subseteq \mathbb{N}$ be the set of counterexamples to the statement. Since we have supposed the statement is false, $X \neq \emptyset$. So by the Well Ordering Principle, X contains a smallest element, s.

(Basis step). We know that $s > 0$ since $0 < 2^0 = 1$.

Since $s > 0$, the statement must be true for $s - 1$; that is:

$$s - 1 < 2^{s-1}$$

But

$$\begin{aligned} & s - 1 < 2^{s-1} \\ \implies & s < 2^{s-1} + 1 \\ \implies & s < (2^{s-1})2 \\ \implies & s < 2^s. \end{aligned}$$

So the statement is true for s. $\Rightarrow\Leftarrow$.

Solution to group exercise #2

Problem. Use proof by smallest counterexample to show that $1 + 2 + 3 + \cdots + n = \frac{1}{2}n^2 + \frac{1}{2}n$ for all positive integers n.

Solution. Suppose, for the sake of contradiction, that the statement is false. Let $X \subseteq \mathbb{N}$ be the set of counterexamples to the statement. Since we have supposed the statement is false, $X \neq \emptyset$. So by the Well Ordering Principle, X contains a smallest element, s.

(Basis step). We know that $s > 1$ since $1 = \frac{1}{2} \cdot 1^2 + \frac{1}{2} \cdot 1$.

Since $s > 1$, the statement must be true for $s - 1$; that is:

$$1 + 2 + 3 + \cdots + s - 1 = \frac{1}{2}(s-1)^2 + \frac{1}{2}(s-1)$$

So, adding s to the LHS and RHS, we have

$$\begin{aligned} 1 + 2 + 3 + \cdots + s - 1 + s &= \frac{1}{2}(s-1)^2 + \frac{1}{2}(s-1) + s \\ &= \frac{1}{2}(s^2 - 2s + 1) + \frac{1}{2}s - \frac{1}{2} + s \\ &= \frac{1}{2}s^2 + \cancel{-s} + \cancel{\frac{1}{2}} + \frac{1}{2}s + \cancel{-\frac{1}{2}} + \cancel{s} \\ &= \frac{1}{2}s^2 + \frac{1}{2}s \end{aligned}$$

So the statement is true for s. $\Rightarrow\Leftarrow$.

Appendix

Optional Remark About Group Exercise #2

In class, group exercise # 2 was presented slightly differently as:

Use proof by smallest counterexample to show that $1 + 2 + 3 + \cdots + n = \binom{n+1}{2}$ *for all positive integers* n.

However, note that

$$\binom{n+1}{2} = \frac{(n+1)!}{2!(n-1)!} = \frac{(n+1)n}{2} = \frac{1}{2}n^2 + \frac{1}{2}n$$

So the original version simplifies to the current version.