

01/29/2026: Discrete Random Variables (Part 2)

CSCI 546: Diffusion Models

Textbook reference: Sec 3.6-3.10

Announcement (Sign-in Sheet)


Please sign the sign-in sheet.

Review Problem Set #4

Concepts for Problem Set #5


Correlation Coefficient

The **correlation coefficient** ρ shows the strength and direction of the **linear** relationship between two random variables X and Y .



Correlation Coefficient

The **correlation coefficient** ρ shows the strength and direction of the **linear** relationship between two random variables X and Y .



Remark. We have

$$|\rho(X, Y)| \leq 1,$$

with equality if and only if X and Y are linearly related (with probability 1).


Correlation Coefficient

Poll. Can uncorrelated random variables be dependent?

Correlation Coefficient

Poll. Can uncorrelated random variables be dependent?

Solution. Yes.



The correlation reflects the noisiness and direction of a linear relationship (top row), but not the slope of that relationship (middle), nor many aspects of nonlinear relationships (bottom).

Random Groups

Aubrey Williams: 1

Austin Barton : 4

Blake Sigmundstad: 7

Diego Moylan: 2

Dillon Shaffer: 1

Ismoiljon Muzaffarov: 3

Jacob Tanner: 6

Josh Stoneback: 1

Joshua Bowen: 5

Joshua Calwell: 8

Laura Banaszewski: 3

Lina Hammel: 8

Logan Racz: 4

Matt Hall: 6

Micah Miller: 7

Mike Kadoshnikov: 5

Owen Cool: 2

Racquel Bowen: 2

Samuel Mocabee: 4

Tatiana Kirillova: 3

Group exercises - Problem Set 5

1. (3.11.16) Let X and Y be independent Bernoulli random variables with parameter $\frac{1}{2}$. Show that $X + Y$ and $|X - Y|$ are dependent though uncorrelated.
2. (Sec 3.6) Use the Cauchy-Schwarz Inequality to show that the correlation coefficient ρ satisfies $|\rho(X, Y)| \leq 1$ with equality if and only if $Y = mX + b$ with probability 1 for some $m, b \in \mathbb{R}$. (Hint: Apply Cauchy-Schwarz to the variables $X - E[X]$ and $Y - E[Y]$.)
3. (3.7.6) Let X_1, X_2, \dots be identically distributed random variables with mean μ and variance σ^2 , and let N be a random variable taking values in the non-negative integers and independent of the X_i . Let $S = X_1 + X_2 + \dots + X_N$.
 - a. Show that $\mathbb{E}[S \mid N] = \mu N$, and deduce that $\mathbb{E}[S] = \mu \mathbb{E}[N]$.
 - b. Find $\text{Var}[S]$ using the conditional variance formula $\text{Var}[S] = \mathbb{E}[\text{Var}[S \mid N]] + \text{Var}[\mathbb{E}[S \mid N]]$.