

{ Surreal Numbers }

2

{ L | R }

Every surreal number consists of a left set and a right set (**L** and **R**), which contain other surreal numbers. Each half can be empty, finite or infinite, but must always satisfy the following condition: ***every number in L must be less than every number in R.***

$$\forall x \in L, \forall y \in R : x < y$$

Such a form { **L** | **R** } represents the simplest surreal number that lies between the sets **L** and **R**.

It is possible for the same number to have different, but equivalent representations.

4

Comparison and order

Suppose we have two surreal numbers, **X** = { **Lx** | **Rx** } and **Y** = { **Ly** | **Ry** }.

We say that **X** ≤ **Y** if and only if everything in **Lx** is less than **Y**, and everything in **Ry** is greater than **X**.

If **X** ≤ **Y**, but **Y** ≠ **X**, then we say that **X** < **Y**. For example, consider **2** = { **1** | } and **ω** = { **1, 2, 3, 4, ...** | }. **1** is less than **ω**, so everything in the left set of **2** is less than **ω**. The right side of **ω** is empty, therefore both conditions are met and **2** ≤ **ω**. However, the reverse statement **ω** ≤ **2** is false, since the left set of **ω** contains **3**, which is greater than **2**.

If **X** ≤ **Y** and **Y** ≤ **X**, then we say that **X** = **Y**. For example, { | } and { -1 | 1 } are equivalent ways to write **0**, since { | } ≤ { -1 | 1 } and { -1 | 1 } ≤ { | }.

6

Multiplication

$$\begin{aligned} \mathbf{X \cdot Y} &= \{ \mathbf{Lx} \mid \mathbf{Rx} \} \cdot \{ \mathbf{Ly} \mid \mathbf{Ry} \} = \\ &= \{ \mathbf{Lx \cdot Y + X \cdot Ly - Lx \cdot Ry, Rx \cdot Y + X \cdot Ry - Rx \cdot Ly} \mid \\ &\quad \mathbf{Lx \cdot Y + X \cdot Ry - Lx \cdot Ry, X \cdot Ly + Rx \cdot Y - Rx \cdot Ly} \} \end{aligned}$$

Multiplication, just like addition, involves multiplying pairs of surreal numbers drawn from both sides of **X** and **Y**.

For example, to compute the term (**Lx**·**Y** + **X**·**Ly** - **Lx**·**Ry**), pick an element **lx** ∈ **Lx**, an element **ly** ∈ **Ly**, and compute (**lx**·**Y** + **X**·**ly** - **lx**·**ly**). Then repeat for all such pairs of elements, and aggregate the results.

This multiplication formula is consistent with the real numbers:

$$\frac{1}{2} \cdot 4 = 2 \quad \{ 0 \mid 1 \} \{ 3 \mid \} = \{ 1 \mid \}$$

1

What’s that?

The field of surreal numbers, usually denoted as **No**, was introduced in 1969 by John H. Conway. It is an extention of the field of Real numbers **R**.

The field of Surreals houses the Reals, the Hyperreals, all of Cantor’s Ordinal numbers and their reciprocals. In a sense, it is the largest possible extention of the Real numbers.

3

Constructing the numbers

$$\{ \mid \} = 0$$

To get **0**, keep both sides empty.

$$\{ \mathbf{n} \mid \} = \mathbf{n + 1} \quad \{ \mid \mathbf{n} \} = \mathbf{n - 1}$$

To get **n + 1**, put **n** in the left set and keep the right set empty. To get **n - 1**, put **n** in the right set instead. This way, we can obtain every integer.

$$\{ \mathbf{a} \mid \mathbf{b} \} = \frac{\mathbf{a + b}}{2}$$

A number exactly between **a** and **b** is expressed as { **a** | **b** }, for example:

$$\{ 0 \mid 1 \} = \frac{1}{2} \quad \{ 0 \mid \frac{1}{2} \} = \frac{1}{4}$$

This gives us all the dyadic rational numbers, which have the form $\frac{a}{2^b}$.

What about $\frac{1}{3}$ or $\sqrt{2}$? We can get any real number **x** by using an increasing and a decreasing sequence of binary fractions, both of which converge to **x**.

$$\sqrt{2} = \{ 1, \frac{5}{4}, \frac{11}{8}, \frac{45}{32} \dots \mid \dots \frac{91}{64}, \frac{23}{16}, \frac{3}{2} \}$$

increasing → ← *decreasing*

We can obtain a number greater than any integer by placing all of them in **L**:

$$\omega = \{ 1, 2, 3, 4, 5 \dots \mid \}$$

This is Cantor’s first ordinal - **ω**. We can also get the first infinitesimal **ε**:

$$\epsilon = \{ 0 \mid \dots \frac{1}{32}, \frac{1}{16}, \frac{1}{8}, \frac{1}{4}, \frac{1}{2} \}$$

5

Addition and subtraction

Let’s add two surreal numbers together: **X** = { **Lx** | **Rx** } and **Y** = { **Ly** | **Ry** }.

$$\mathbf{X + Y} = \{ \mathbf{Lx + Y, X + Ly} \mid \mathbf{Rx + Y, X + Ry} \}$$

For each element in **Lx**, add with **Y**, and put the result in the *left* set of the sum. Also take every element in **Ly**, add it with **X**, and put the result in the *left* set of the sum. For the *right* set, we take every element in **Rx** plus **Y**, and every element in **Ry** plus **X**.

For example, let **X** = (**ω** + **1**) and **Y** = **1**. We should expect to get (**ω** + **2**):

$$\begin{aligned} (\omega + 1) + 1 &= \\ &= \{ \omega \mid \} + \{ 0 \mid \} \\ &= \{ \omega + 1, \omega + 1 \mid \} \\ &= \{ \omega + 1 \mid \} = \omega + 2 \end{aligned}$$

Write **X** as { **ω** | }, and **Y** as { **0** | }.

For the left part, we take **X** and add it to every element in **Ly**. Since **Ly** only contains **0**, this gets us **X** + **0**, which is (**ω** + **1**). We also add **Y** to the elements of **Lx** (which only contains **ω**), and get (**ω** + **1**) again.

Since **Rx** and **Ry** are empty, there is nothing to add for the right part of the sum - it’s empty.

What about subtraction? Let’s see how to negate a number **X** = { **Lx** | **Rx** }:

$$-\mathbf{X} = \{ -\mathbf{Rx} \mid -\mathbf{Lx} \}$$

To get **-X**, swap the right and left sets, then negate every element in both sets. For example, **-2** = - { **1** | } = { | -1 }. To subtract **Y** from **X**, just take the negative of **Y** and add it to **X**:

$$\mathbf{X - Y} = \mathbf{X + (-Y)}$$