


Protein: YNL245C


IUPred2A:



Function:

Involved in pre-mRNA splicing.

Structure:




Subcellular Location: Nucleus

Naša procjena: IDP

Protein: YDL201W


IUPred2A:



Function:

Methyltransferase that catalyzes the formation of N7-methylguanine at position 46 (m7G46) in tRNA, a modification required to maintain stability of tRNAs; its absence resulting in tRNA decay. Both the D-stem and T-stem structures of tRNAs are required for efficient methyltransferase activity.

Structure:




Subcellular Location: Nucleus

Naša procjena: IDP

Protein: YDR424C

IUPred2A:



Function: Acts as one of several non-catalytic accessory components of the cytoplasmic dynein complex that are thought to be involved in linking dynein to cargos and to adapter proteins that regulate dynein function. Cytoplasmic dynein 1 acts as a motor for the intracellular retrograde motility of vesicles and organelles along microtubules. May play a role in changing or maintaining the spatial distribution of cytoskeletal structures (By similarity).

Also a component of the nuclear pore complex where it may contribute to the stable association of the Nup82 subcomplex with the NP

Structure:




Subcellular Location: Cytoplasm, cytoskeleton Nucleus, nuclear pore complex

Naša procjena: IDP

Protein: YKL103C


IUPred2A:



Function:

Resident vacuolar enzyme that catalyzes the removal of amino acids from the N-terminus of peptides and proteins. Also acts as the major cargo protein of the cytoplasm-to-vacuole targeting (Cvt) pathway. The precursor form of aminopeptidase 1 (prApe1) assembles into dodecamers and the propeptide mediates the aggregation of dodecamers into higher multimers. The multimers are then recognized via the propeptide by their receptor ATG19, and ATG19 further interacts with ATG11, which tethers the APE1-ATG19 complex to the pre-autophagosomal structure (PAS). The cargo-receptor complex (also Cvt complex) is selectively enwrapped by a double-membrane structure termed the Cvt vesicle under vegetative growth conditions and by a similar but larger double-membrane structure termed the autophagosome under nitrogen starvation conditions. The Cvt vesicle or the autophagosome fuses with the vacuolar membrane and release its content in the vacuolar lumen. In the vacuole, prApe1 is processed into mature aminopeptidase 1 (mApe1)

Structure:




Subcellular Location: Vacuole

Naša procjena: non-IDP

Protein: YFR011C


IUPred2A:



Function:

Component of the MICOS complex, a large protein complex of the mitochondrial inner membrane that plays crucial roles in the maintenance of crista junctions, inner membrane architecture, and formation of contact sites to the outer membrane

Structure:




Subcellular Location: Mitochondrion inner membrane

Naša procjena: non-IDP???

Protein: YKL172W

IUPred2A:



Function:

Required for the processing of the 27S pre-rRNA. Probably involved in the step of the processing of the 27 SA precursor into the 27 SB intermediate

Structure:



Subcellular Location: Nucleus, nucleolus

Naša procjena: IDP