

# Introductory Functional Analysis with Applications Solutions

Michael You



# Contents

<b>I</b>	<b>Metric Spaces</b>	<b>1</b>
<b>1</b>	<b>Metric Spaces</b>	<b>3</b>
1.1	Metric Space . . . . .	3
1.2	. . . . .	8
1.3	. . . . .	9
1.4	. . . . .	10
1.5	. . . . .	11
1.6	. . . . .	12
<b>2</b>	<b>Normed Spaces. Banach Spaces</b>	<b>13</b>
2.1	. . . . .	13
2.2	. . . . .	14
2.3	. . . . .	15
2.4	. . . . .	16
2.5	. . . . .	17
2.6	. . . . .	18
2.7	. . . . .	19
2.8	. . . . .	20
2.9	. . . . .	21
2.10	. . . . .	22
<b>II</b>	<b>Banach Spaces</b>	<b>23</b>
<b>3</b>	<b>Inner Product Spaces. Hilbert Spaces</b>	<b>25</b>
3.1	. . . . .	25
3.2	. . . . .	26
3.3	. . . . .	27
3.4	. . . . .	28
3.5	. . . . .	29
3.6	. . . . .	30
3.7	. . . . .	31
3.8	. . . . .	32
3.9	. . . . .	33
3.10	. . . . .	34
<b>4</b>	<b>Fundamental Theorems for Normed and Banach Spaces</b>	<b>35</b>
4.1	. . . . .	35
4.2	. . . . .	36
4.3	. . . . .	37
4.4	. . . . .	38
4.5	. . . . .	39
4.6	. . . . .	40

4.7	41
4.8	42
4.9	43
4.10	44
4.11	45
4.12	46
4.13	47
<b>5 Further Applications: Banach Fixed Point Theorem</b>	<b>49</b>
5.1	49
5.2	50
5.3	51
5.4	52
 <b>III Hilbert Spaces</b>	 <b>53</b>
<b>6 Further Applications: Approximation Theory</b>	<b>55</b>
6.1	55
6.2	56
6.3	57
6.4	58
<b>7 Spectral Theory of Linear Operators in Normed Spaces</b>	<b>59</b>
7.1	59
7.2	60
7.3	61
7.4	62
7.5	63
7.6	64
<b>8 Compact Linear Operators on Normed Spaces and Their Spectrum</b>	<b>65</b>
8.1	65
8.2	66
8.3	67
8.4	68
8.5	69
8.6	70
 <b>IV Operators on Hilbert Spaces</b>	 <b>71</b>
<b>9 Spectral Theory of Bounded Self-Adjoint Linear Operators</b>	<b>73</b>
9.1	73
9.2	74
9.3	75
9.4	76
9.5	77
9.6	78
9.7	79
9.8	80
9.9	81
9.10	82
9.11	83

<b>10 Unbounded Linear Operators in Hilbert Space</b>	<b>85</b>
10.1 . . . . .	85
10.2 . . . . .	86
10.3 . . . . .	87
10.4 . . . . .	88
10.5 . . . . .	89
10.6 . . . . .	90
<b>11 Unbounded Linear Operators in Quantum Mechanics</b>	<b>91</b>
11.1 . . . . .	91
11.2 . . . . .	92
11.3 . . . . .	93
11.4 . . . . .	94
11.5 . . . . .	95
11.6 . . . . .	96



Part I

Metric Spaces



# Chapter 1

## Metric Spaces

### 1.1 Metric Space

We spend this chapter introducing a metric space and building some intuition around it. From the definition below, if we can show they are true, then we have a metric space.

**Definition 1** A **metric space** is a pair  $(X, d)$  where  $X$  is a set, and  $d$  is a *metric on  $X$*  (or *distance function on  $X$* ), that is, a function defined on  $X \times X$  such that for all  $x, y, z \in X$  we have:

1.  $d$  is real-valued, finite and nonnegative
2.  $d(x, y) = 0$  if and only if  $x = y$
3.  $d(x, y) = d(y, x)$  (Symmetry)
4.  $d(x, y) \leq d(x, z) + d(z, y)$  (**Triangle inequality**)

#### Exercise 1.1.1

Let's show that  $\mathbb{R}$  is a metric space. Since the author didn't specify, I assume they mean the standard metric  $d(x, y) = |x - y|$ .

In this case, we can show

1.  $d(x, y) = |x - y| \geq 0$ , by definition of absolute value. It is finite, since  $x, y$  are finite, and you can bound  $|x - y|$  by  $|x| + |y|$  which is also finite. And it is  $\in \mathbb{R}$  by definition of absolute value.
2. If  $x = y$  then  $d(x, y) = |x - y| = 0$ . If  $d(x, y) = 0$ , then  $|x - y| = 0$ , which means  $x - y = 0$ , or  $x = y$ .
3.  $d(x, y) = |x - y| = |y - x| = d(y, x)$ .
4.  $d(x, y) = |x - y| = |x - z + z - y| \leq |x - z| + |z - y| = d(x, z) + d(z, y)$

#### Exercise 1.1.2

Same as 1.1.1, but for  $d(x, y) = (x - y)^2$ .

Counterexample for Triangle Inequality: Let  $x = 0, y = 2, z = 1$ . Then, we have

$$\begin{aligned}d(x, y) &= (2 - 0)^2 = 4 \\d(x, z) + d(z, y) &= (0 - 1)^2 + (1 - 2)^2 = 1 + 1 = 2 \\4 &\not\leq 2\end{aligned}$$

So it is **not a metric**.

#### Exercise 1.1.3

Same as 1.1.1, but for  $d(x, y) = \sqrt{|x - y|}$ .

We can show

1.  $d(x, y) = \sqrt{|x - y|} \geq 0$ , by definition of absolute value and square root. It is finite, since  $x, y$  are finite, and you can bound  $\sqrt{|x - y|}$  by  $\sqrt{|x| + |y|}$  which is also finite. And it is  $\in \mathbb{R}$  by definition of absolute value and square root.
2. If  $x = y$  then  $d(x, y) = \sqrt{|x - y|} = 0$ . If  $d(x, y) = 0$ , then  $\sqrt{|x - y|} = 0$ , which means  $|x - y| = 0$ , or  $x - y = 0$ , or  $x = y$ .
3.  $d(x, y) = \sqrt{|x - y|} = \sqrt{|y - x|} = d(y, x)$ .
4.  $d(x, y) = \sqrt{|x - y|} = \sqrt{|x - z + z - y|} \leq \sqrt{|x - z|} + \sqrt{|z - y|} = d(x, z) + d(z, y)$   
The inequality step comes from the fact that  $\sqrt{a + b} \leq \sqrt{a} + \sqrt{b}$  for nonnegative  $a, b$ , because

$$\begin{aligned} & (\sqrt{a} + \sqrt{b})^2 \leq (\sqrt{a + b})^2 \\ & a + b + 2\sqrt{ab} \geq a + b \end{aligned}$$

So it is a metric.

#### Exercise 1.1.4

We need to find all metrics on a set  $X$  consisting of two points.

Let  $X = \{a, b\}$  with  $a \neq b$ . Suppose we have the following metric:

$$d(x, y) = \begin{cases} 0 & x = y \\ c & x \neq y, c > 0 \end{cases}$$

We can verify 1, 2, 3, from 1 very easily. The main caveat, we needed  $c > 0$  for 2, for  $d(x, y) = 0$  iff  $x = y$ . Then, for 4 (Triangle Inequality), we can check all possible cases:

1. If  $x = y$ , then  $d(x, y) = 0 \leq d(x, z) + d(z, y)$
2. if  $x \neq y$ , then we have two more cases
  - (a) if  $x = z$ , then  $z \neq y$  and we have  $d(x, z) + d(z, y) = c \geq c = d(x, y)$
  - (b) if  $x \neq z$ , then we have  $z = y$  and similarly  $d(x, z) + d(z, y) = c \geq c = d(x, y)$

If  $X$  consists of one point, then the only metric is the trivial one where  $d(x, x) = 0$ . This is because, if we have any other metric, then  $d(x, x) > 0$  which violates property 2.

#### Exercise 1.1.5

Suppose  $d$  is a metric on  $X$ .

1. Suppose we have  $kd$ . Then this is still a metric if  $k > 0$ . We can verify the four properties:
  - (a)  $kd(x, y) \geq 0$  since  $d(x, y) \geq 0$  and  $k > 0$ . It is finite since  $d$  is finite and  $k$  is a finite scalar.
  - (b)  $kd(x, y) = 0$  iff  $d(x, y) = 0$  iff  $x = y$ .
  - (c)  $kd(x, y) = kd(y, x)$ , we can just scale property 3
  - (d)  $kd(x, y) \leq kd(x, z) + kd(z, y)$  we can just scale property 4 the triangle inequality.
2. Suppose we have  $d + k$ . Then this is only a metric for  $k = 0$ . If  $k \neq 0$  then property 2 is immediately dissatisfied, since  $d(x, x) + k = k \neq 0$ .

#### Exercise 1.1.6

We want to show the sequence space  $l^\infty$  is a metric space.

1.  $d(x, y) = \sup_{n \in \mathbb{N}} |x_n - y_n| \geq 0$  since absolute value is nonnegative. It is finite since  $x, y \in l^\infty$  means they are bounded sequences by  $c_x \in \mathbb{R}$ , so their difference is also bounded. And it is  $\in \mathbb{R}$  since absolute value produces real numbers.

2. If  $x = y$ , then  $d(x, y) = \sup |x_n - y_n| = 0$ . If  $d(x, y) = 0$ , then  $\sup |x_n - y_n| = 0$ , which means for all  $n$ ,  $|x_n - y_n| = 0$ , or  $x_n = y_n$  for all  $n$ , so  $x = y$ .
3.  $d(x, y) = \sup |x_n - y_n| = \sup |y_n - x_n| = d(y, x)$ .
4.  $d(x, y) = \sup |x_n - y_n| = \sup |x_n - z_n + z_n - y_n| \leq \sup |x_n - z_n| + \sup |z_n - y_n| = d(x, z) + d(z, y)$ .

**Exercise 1.1.7**

Supposed  $A$  is the subset of  $l^\infty$  where each sequence is composed of 0 or 1.

The induced metric then behaves like this:

$$d(x, y) = \begin{cases} 0 & x = y \\ 1 & x \neq y \end{cases}$$

**Exercise 1.1.8**

We have  $X$  as the set of all real-valued functions, and are defined and continuous on a closed interval  $J = [a, b]$ . We now choose the metric

$$d(x, y) = \int_a^b |x(t) - y(t)| dt \quad (1.1)$$

Let's check this is a metric

1.  $d(x, y) = \int_a^b |x(t) - y(t)| dt \geq 0$  since absolute value is nonnegative. It is finite since  $x, y$  are continuous on a closed interval, so they are bounded, and the integral of a bounded function over a finite interval is finite. And it is  $\in \mathbb{R}$  since the integral of a real-valued function is real.
2. If  $x = y$ , then  $d(x, y) = \int_a^b |x(t) - y(t)| dt = \int_a^b |0| dt = 0$ . If  $d(x, y) = 0$ , then  $\int_a^b |x(t) - y(t)| dt = 0$ , which implies  $|x(t) - y(t)| = 0$  for all  $t \in [a, b]$ , so  $x(t) = y(t)$  for all  $t \in [a, b]$ .
3.  $d(x, y) = \int_a^b |x(t) - y(t)| dt = \int_a^b |y(t) - x(t)| dt = d(y, x)$ .
4.  $d(x, y) = \int_a^b |x(t) - y(t)| dt = \int_a^b |x(t) - z(t) + z(t) - y(t)| dt \leq \int_a^b |x(t) - z(t)| dt + \int_a^b |z(t) - y(t)| dt = d(x, z) + d(z, y)$ .

**Exercise 1.1.9**

We want to show that the discrete metric for  $X$  is a metric, this is

$$d(x, x) = 0, d(x, y) = 1 \text{ for } x \neq y \quad (1.2)$$

1.  $d(x, y)$  is either 0 or 1, so it is nonnegative, finite, and real-valued.
2. By definition,  $d(x, y) = 0$  iff  $x = y$ .
3. By definition,  $d(x, y) = d(y, x)$ .
4. For the triangle inequality, we have two cases:
  - (a) If  $x = y$ , then  $d(x, y) = 0 \leq d(x, z) + d(z, y)$
  - (b) If  $x \neq y$ , then we have two more cases
    - i. if  $x = z$ , then  $z \neq y$  and we have  $d(x, z) + d(z, y) = 1 \geq 1 = d(x, y)$
    - ii. if  $x \neq z$ , then we have  $z = y$  and similarly  $d(x, z) + d(z, y) = 1 \geq 1 = d(x, y)$

**Exercise 1.1.10**

**(Hamming distance)** Let  $X$  be all set of ordered triples of 0s and 1s.

We immediately see that  $X$  has  $2^3 = 8$  elements.

Now,  $d(x, y)$  = number of positions where  $x$  and  $y$  differ.

1.  $d(x, y) = 0, 1, 2, 3$  so it is nonnegative, finite, and real-valued.

2. If  $x = y$ , then they differ in 0 positions, so  $d(x, y) = 0$ . If  $d(x, y) = 0$ , then they differ in 0 positions, so  $x$  and  $y$  must be the same in all positions, so  $x = y$ .
3. The number of positions where  $x$  and  $y$  differ is the same as the number of positions where  $y$  and  $x$  differ, so  $d(x, y) = d(y, x)$ .
4. For the triangle inequality, we can prove this by induction using the discrete metric space, as the base case. Then, suppose we have  $n$ -tuples, and we add one more position to each tuple to make it an  $(n + 1)$ -tuple. There are two cases:
  - (a) If the new position is the same for all three tuples, then the number of differing positions is unchanged, so the triangle inequality holds by the inductive hypothesis.
  - (b) If the new position is different for at least one of the tuples, then the left-hand side of the triangle inequality increases by at most 1, and the right-hand side also increases by at least 1, because in  $z$ , its new position must be different from at least  $x$  or  $y$ .

**Exercise 1.1.11**

We want to prove the generalized triangle inequality:

$$d(x_1, x_n) \leq d(x_1, x_2) + d(x_2, x_3) + \cdots + d(x_{n-1}, x_n) \quad (1.3)$$

We can do this by induction.

Base case:  $n = 2$  is just the regular triangle inequality. Inductive step: Suppose it is true for  $n = k$ . Then, for  $n = k + 1$ , we have

$$\begin{aligned} d(x_1, x_{k+1}) &\leq d(x_1, x_k) + d(x_k, x_{k+1}) && \text{(Triangle Inequality)} \\ &\leq d(x_1, x_2) + d(x_2, x_3) + \cdots + d(x_{k-1}, x_k) + d(x_k, x_{k+1}) && \text{(Inductive Hypothesis)} \end{aligned}$$

Thus, by induction, the generalized triangle inequality holds for all  $n \geq 2$ .

**Exercise 1.1.12**

We can see that

$$\begin{aligned} |d(x, y) - d(z, w)| &\leq d(x, z) + d(y, w) \\ d(x, y) - d(z, w) &\leq d(x, z) + d(y, w) && \text{(Case 1, we'll do the other case next)} \\ d(x, y) &\leq d(x, z) + d(y, w) + d(z, w) \\ d(x, y) &\leq d(x, z) + d(w, y) + d(z, w) && \text{(Symmetry)} \\ d(x, y) &\leq d(x, z) + d(z, w) + d(w, y) && \text{(Triangle Inequality)} \end{aligned}$$

We need to do the other case as well:

$$\begin{aligned} d(z, w) - d(x, y) &\leq d(x, z) + d(y, w) \\ d(z, w) &\leq d(x, z) + d(y, w) + d(x, y) \\ d(z, w) &\leq d(z, x) + d(x, y) + d(y, w) && \text{(Symmetry)} \\ d(z, w) &\leq d(z, x) + d(x, y) + d(y, w) && \text{(Triangle Inequality)} \end{aligned}$$

**Exercise 1.1.13**

$$\begin{aligned} |d(x, z) - d(y, z)| &\leq d(x, y) \\ d(x, z) - d(y, z) &\leq d(x, y) && \text{(Case 1, we'll do the other case next)} \\ d(x, z) &\leq d(x, y) + d(y, z) && \text{(Triangle Inequality)} \end{aligned}$$

Let's do the other case:

$$\begin{aligned}
 d(y, z) - d(x, z) &\leq d(x, y) \\
 d(y, z) &\leq d(x, y) + d(x, z) \\
 d(y, z) &\leq d(y, x) + d(x, z) && \text{(Symmetry)} \\
 d(y, z) &\leq d(y, x) + d(x, z) && \text{(Triangle Inequality)}
 \end{aligned}$$

**Exercise 1.1.14**

We want to show that properties 2-4 in the 1 definition imply that the metric must be nonnegative.

AFSOC that there exists some  $x, y \in X$  such that  $d(x, y) < 0$ . Then, by property 4, we have

$$d(x, x) \leq d(x, y) + d(y, x)$$

By property 3, we have  $d(y, x) = d(x, y)$ , so

$$d(x, x) \leq 2d(x, y)$$

But by property 2, we have  $d(x, x) = 0$ , so

$$0 \leq 2d(x, y)$$

However, we assumed  $d(x, y) < 0$ , so  $2d(x, y) < 0$ , which is a contradiction. Therefore, our assumption is false, and we conclude that for all  $x, y \in X$ ,  $d(x, y) \geq 0$ .

**1.2**

## 1.3

**1.4**

## 1.5

**1.6**

## Chapter 2

# Normed Spaces. Banach Spaces

### 2.1

**2.2**

**2.3**

**2.4**

**2.5**

**2.6**

**2.7**

**2.8**

**2.9**

**2.10**

**Part II**

**Banach Spaces**



## Chapter 3

# Inner Product Spaces. Hilbert Spaces

### 3.1

**3.2**

**3.3**

**3.4**

**3.5**

**3.6**

**3.7**

**3.8**

**3.9**

**3.10**

## Chapter 4

# Fundamental Theorems for Normed and Banach Spaces

### 4.1

**4.2**

**4.3**

**4.4**

**4.5**

**4.6**

**4.7**

**4.8**

**4.9**

**4.10**

## 4.11

**4.12**

## 4.13



## Chapter 5

# Further Applications: Banach Fixed Point Theorem

### 5.1

**5.2**

**5.3**

**5.4**

Part III

Hilbert Spaces



## Chapter 6

# Further Applications: Approximation Theory

### 6.1

**6.2**

**6.3**

**6.4**

## Chapter 7

# Spectral Theory of Linear Operators in Normed Spaces

### 7.1

**7.2**

**7.3**

**7.4**

**7.5**

**7.6**

## Chapter 8

# Compact Linear Operators on Normed Spaces and Their Spectrum

### 8.1

## 8.2

**8.3**

## 8.4

**8.5**

## 8.6

## Part IV

# Operators on Hilbert Spaces



## Chapter 9

# Spectral Theory of Bounded Self-Adjoint Linear Operators

### 9.1

**9.2**

## 9.3

**9.4**

## 9.5

**9.6**

## 9.7

**9.8**

## 9.9

**9.10**

## 9.11



## Chapter 10

# Unbounded Linear Operators in Hilbert Space

### 10.1

**10.2**

## 10.3

**10.4**

## 10.5

**10.6**

## Chapter 11

# Unbounded Linear Operators in Quantum Mechanics

### 11.1

**11.2**

## 11.3

**11.4**

## 11.5

**11.6**