

Notes on Noncommutative Geometry and Particle Physics

Popovic Milutin

Week 5: 12.03 - 19.03

Contents

1 Noncommutative Geometric Spaces	1
1.1 Exercises	1
1.2 Properties of Matrix Algebras	3
1.3 Morphisms Between Finite Spectral Triples	4
1.4 Graphing Finite Spectral Triples	5

1 Noncommutative Geometric Spaces

1.1 Exercises

Exercise 1. Make the proof of the last theorem (see week4.pdf) explicit for $N = 3$

Solution 1. For the C^* algebra we have $A = \mathbb{C}^3$ For H we have $H = (\mathbb{C}^2)^{\oplus 3} = H_2 \oplus H_2^1 \oplus H_2^2$. The symmetric operator D acting on H and the representation $\pi(a)$:

$$\begin{aligned} \pi((a(1), a(2), a(3))) &= \begin{pmatrix} a(1) & 0 \\ 0 & a(2) \end{pmatrix} \oplus \begin{pmatrix} a(1) & 0 \\ 0 & a(3) \end{pmatrix} \oplus \begin{pmatrix} a(2) & 0 \\ 0 & a(2) \end{pmatrix} \\ &= \begin{pmatrix} a(1) & 0 & 0 & 0 & 0 & 0 \\ 0 & a(2) & 0 & 0 & 0 & 0 \\ 0 & 0 & a(1) & 0 & 0 & 0 \\ 0 & 0 & 0 & a(3) & 0 & 0 \\ 0 & 0 & 0 & 0 & a(2) & 0 \\ 0 & 0 & 0 & 0 & 0 & a(3) \end{pmatrix} \end{aligned} \quad (1)$$

$$\begin{aligned} D &= \begin{pmatrix} 0 & x_1 \\ x_1 & 0 \end{pmatrix} \oplus \begin{pmatrix} 0 & x_1 \\ x_1 & 0 \end{pmatrix} \oplus \begin{pmatrix} 0 & x_1 \\ x_1 & 0 \end{pmatrix} \\ &= \begin{pmatrix} 0 & x_1 & 0 & 0 & 0 & 0 \\ x_1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & x_2 & 0 & 0 \\ 0 & 0 & x_2 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & x_3 \\ 0 & 0 & 0 & 0 & x_3 & 0 \end{pmatrix} \end{aligned} \quad (2)$$

(3)

Then the norm of the commutator would be the largest eigenvalue

$$\begin{aligned}
||[D, \pi(a)]|| &= ||D\pi(a) - \pi(a)D|| \\
&= \left\| \begin{pmatrix} 0 & x_1(a(2) - a(1)) & 0 & 0 & 0 & 0 \\ -x_1(a(2) - a(1)) & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & -x_2(a(3) - a(1)) & x_2(a(3) - a(1)) & 0 & 0 \\ 0 & 0 & 0 & 0 & -x_3(a(2) - a(3)) & x_3(a(3) - a(2)) \\ 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix} \right\| \\
&\quad (4)
\end{aligned}$$

The matrix in Equation ?? is a skew symmetric matrix its eigenvalues are $i\lambda_1, i\lambda_2, i\lambda_3, i\lambda_4$, where the λ 's are on the upper and lower diagonal check https://en.wikipedia.org/wiki/Skew-symmetric_matrix#Skew-symmetrizable_matrix. The matrix norm of would be the maximum of the norm of the larges eigenvalues:

$$||[D, \pi(a)]|| = \max_{a \in A} \{x_i | a(j) - a(k)\} \quad (5)$$

Exercise 2. Compute the metric on the space of three points given by $d_{ij} = \sup_{a \in A} \{|a(i) - a(j)| : ||[D, \pi(a)]|| \leq 1\}$ for the set of data $A = \mathbb{C}^3$ acting in the defining representation $H = \mathbb{C}^3$, and

$$D = \begin{pmatrix} 0 & d^{-1} & 0 \\ d^{-1} & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$$

for some $d \in \mathbb{R}$

Solution 2. We have $A = \mathbb{C}^3$, $H = \mathbb{C}^3$ and D from above, then

$$||[D, \pi(a)]|| = d^{-1} \left\| \begin{pmatrix} 0 & a(2) - a(1) & 0 \\ -(a(2) - a(1)) & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \right\| \quad (6)$$

$$= d^{-1} |a(2) - a(1)| \quad (7)$$

Exercise 3. Show that d_{ij} from Equation 11 is a metric on \hat{A} by establishing that:

$$d_{ij} = 0 \Leftrightarrow i = j \quad (8)$$

$$d_{ij} = d_{ji} \quad (9)$$

$$d_{ij} \leq d_{ik} + d_{kj} \quad (10)$$

$$d_{ij} = \sup_{a \in A} \{|Tr(a(i)) - Tr(a(j))| : ||[D, a]|| \leq 1\} \quad (11)$$

Solution 3. For Equation 8 set $i = j$ in 11.

$$\begin{aligned}
d_{ii} &= \sup_{a \in A} \{|Tr(a(i)) - Tr(a(i))| : ||[D, a]|| \leq 1\} \\
&= \sup_{a \in A} \{0 : ||[D, a]|| \leq 1\} = 0
\end{aligned}$$

For Equation 9 obviously we have the commuting property of addition.

For Equation 10, for $k = j$ then $d_{kj} = 0$ and the equality holds. For $i = k$ then $d_{ik} = 0$ and equality holds. Else set $d_{ik} = 1$ and $d_{kj} = 1$ then $d_{ij} = 1 \leq d_{ik} + d_{kj} = 2$

1.2 Properties of Matrix Algebras

Lemma 1. *If A is a unital C^* algebra that acts faithfully on a finite dimensional Hilbert space, then A is a matrix algebra of the Form:*

$$A \simeq \bigoplus_{i=1}^N M_{n_i}(\mathbb{C}) \quad (12)$$

Proof. Since A acts faithfully on a Hilbert space, then A is a C^* subalgebra of a matrix algebra $L(H) = M_{\dim(H)}(\mathbb{C}) \Rightarrow A \simeq \text{Matrix algebra}$. \square

Question 1. What does the author mean when he says 'acts faithfully on a Hilbertspace'? Then the representation is fully reducible, or that the presentation is irreducible?

Example 1. $A = M_n(\mathbb{C})$ and $H = \mathbb{C}^n$, A acts on H with matrix multiplication and standard inner product. D on H is a hermitian matrix $n \times n$ matrix.

D is referred to as a finite Dirac operator as in as its ∞ dimensional on Riemannian Spin manifolds coming in Chapter 4. Now we introduce it as

$$\frac{a(i) - a(j)}{d_{ij}} \quad (13)$$

for each pair $i, j \in X$ the finite dimensional discrete space. This appears in the entries in the commutator $[D, a]$ in the above exercises.

Definition 1. Given an finite spectral triple (A, H, D) , the A -bimodule of Connes' differential one form is:

$$\Omega_D^1(A) := \left\{ \sum_k a_k [D, b_k] : a_k, b_k \in A \right\} \quad (14)$$

Question 2. Is the Conne's differential one form the set of all '1st order differential operators' given A , that act on H ?

Then there is a map $d : A \rightarrow \Omega_D^1(A)$, $d = [D, \cdot]$.

Lemma 2. *Let $(A, H, D) = (M_n(\mathbb{C}, \mathbb{C}^n, D)$, with D a hermitian $n \times n$ matrix. If D is not a multiple of the identity then:*

$$\Omega_D^1(A) \simeq M_n(\mathbb{C}) = A \quad (15)$$

Proof. Assume $D = \sum_i \lambda_i e_{ii}$ (diagonal), $\lambda_i \in \mathbb{R}$ and $\{e_{ij}\}$ the basis of $M_n(\mathbb{C})$. For fixed i, j choose k such that $\lambda_k \neq \lambda_j$ then

$$\left(\frac{1}{\lambda_k - \lambda_j} e_{ik} \right) [D, e_{kj}] = e_{ij}$$

$e_{ij} \in \Omega_D^1(A)$ by the above definition. And $\Omega_D^1(A) \subset L(\mathbb{C}^n) = H \simeq M_n(\mathbb{C}) = A$ \square

1.3 Morphisms Between Finite Spectral Triples

Definition 2. two finite spectral triples (A_1, H_1, D_1) and (A_2, H_2, D_2) are called unitarily equivalent if

- $A_1 = A_2$
- $\exists U : H_1 \rightarrow H_2$, unitary with
 1. $U\pi_1(a)U^* = \pi_2(a)$ with $a \in A_1$
 2. $UD_1U^* = D_2$

Some remarks

- the above is an equivalence relation
- spectral unitary equivalence is given by the unitaries of the matrix algebra itself
- for any such U then $(A, H, D) \sim (A, H, UDU^*)$
- $UDU^* = D + U[D, U^*]$ of the form of elements in $\Omega_D^1(A)$.

Extending the this relation we look again at the notion of equivalence from Morita equivalence of Matrix Algebras.

Given a Hilbert bimodule $E \in KK_f(B, A)$ and (A, H, D) we construct a finite spectral triple on B , (B, H', D')

$$H' = E \otimes_A H \quad (16)$$

This extends the left action on B with the right action and inherits the \mathbb{C} valued inner product space.

$$D'(e \otimes \xi) = e \otimes D\xi + \nabla(e)\xi \quad e \in E, a \in A \quad (17)$$

Where ∇ is called the *connection on the right A -module E* associated with the derivation $d = [D, \cdot]$ and satisfying the *Leibnitz Rule* which is

$$\nabla(ae) = \nabla(e)a + e \otimes [D, a] \quad e \in E, a \in A \quad (18)$$

Then the linearity of the balanced tensor product $E \otimes_A H$ is satisfied

$$\begin{aligned} D'(ea \otimes \xi - e \otimes a\xi) &= D'(ea \otimes \xi) - D'(e \otimes a\xi) \\ &= ea \otimes D\xi + \nabla(ea)\xi - e \otimes D(a\xi) - \nabla(e)a\xi \\ &= 0 \end{aligned}$$

With the information thus far we can prove the following theorem

Theorem 1. *If (A, H, D) a finite spectral triple, $E \in KK_f(B, A)$. Then $(B, E \otimes_A H, D')$ is a finite spectral triple, provided that ∇ satisfies the compatibility condition*

$$\langle e_1, \nabla e_2 \rangle_E - \langle \nabla e_1, e_2 \rangle_E = d\langle e_1, e_2 \rangle_E \quad e_1, e_2 \in E \quad (19)$$

Proof. $E \otimes_A H$ was shown in the previous section (text before the theorem). The only thing left is to show that D' is a symmetric operator, this we can just compute. Let

$e_1, e_2 \in E$ and $\xi_1, \xi_2 \in H$ then

$$\begin{aligned}
\langle e_1 \otimes \xi_1, D'(e_2 \otimes \xi_2) \rangle_{E \otimes_A H} &= \langle \xi_1, \langle e_1, \nabla e_2 \rangle_E \xi_2 \rangle + \langle \xi_1, \langle e_1, e_2 \rangle_E D \xi_2 \rangle_H \\
&= \langle \xi_1, \langle \nabla e_1, e_2 \rangle_E \xi_2 \rangle_H + \langle \xi_1, d \langle e_1, e_2 \rangle_E \xi_2 \rangle_H \\
&+ \langle D \xi_1, \langle e_1, e_2 \rangle_E \xi_2 \rangle_H - \langle \xi_1, [D, \langle e_1, e_2 \rangle_E] \xi_2 \rangle_H \\
&= \langle D'(e_1 \otimes \xi_1), e_2 \otimes \xi_2 \rangle_{E \otimes_A H}
\end{aligned}$$

□

1.4 Graphing Finite Spectral Triples

Definition 3. A *graph* is a ordered pair $(\Gamma^{(0)}, \Gamma^{(1)})$. Where $\Gamma^{(0)}$ is the set of vertices (nodes) and $\Gamma^{(1)}$ a set of pairs of vertices (edges)




Figure 1: A simple graph with three vertices and three edges

Definition 4. A Λ -decorated graph is given by an ordered pair (Γ, Λ) of a finite graph Γ and a set of positive integers Λ with the labeling

- of the vetices $v \in \Gamma^{(0)}$ given by $n(v) \in \Lambda$
- of the edges $e = (v_1, v_2) \in \Gamma^{(1)}$ by operators
 - $D_e : \mathbb{C}^{n(v_1)} \rightarrow \mathbb{C}^{n(v_2)}$
 - and $D_e^* : \mathbb{C}^{n(v_2)} \rightarrow \mathbb{C}^{n(v_1)}$ its conjugate traspose (pullback?)

such that

$$n(\Gamma^{(0)}) = \Lambda \quad (20)$$

Question 3. Would then D_e be the pullback?

Question 4. These graphs are important in the next chapter I should look into it more, I don't understand much here, specific how to construct them with the abstraction of a spectral triple...

The operator D_e between \mathbf{n}_i and \mathbf{n}_j add up to D_{ij}

$$D_{ij} = \sum_{\substack{e=(v_1, v_2) \\ n(v_1)=\mathbf{n}_i \\ n(v_2)=\mathbf{n}_j}} D_e$$

Theorem 2. *There is a on to one correspondence between finite spectral triples modulo unitary equivalence and Λ -decorated graphs, given by associating a finite spectral triples (A, H, D) to a Λ decorated graph (Γ, Λ) in the following way:*

$$A = \bigoplus_{n \in \Lambda} M_n(\mathbb{C}); \quad H = \bigoplus_{v \in \Gamma^{(0)}} \mathbb{C}^{n(v)}; \quad D = \sum_{e \in \Gamma^{(1)}} D_e + D_e^* \quad (21)$$

Example 2.




Figure 2: A Λ -decorated Graph of $(M_n(\mathbb{C}), \mathbb{C}^n, D = D_e + D_e^*)$