

1 Univerzita Karlova v Praze, Filozofická fakulta
2 Katedra logiky

3 MIKULÁŠ MRVA

4 REFLECTION PRINCIPLES AND LARGE
5 CARDINALS
6 Bakalářská práce

7 Vedoucí práce: Mgr. Radek Honzík, Ph.D.

8 2016

¹⁰ Prohlašuji, že jsem bakalářskou práci vypracoval samostatně a že jsem uvedl
¹¹ všechny použité prameny a literaturu.

¹² V Praze 22. května 2016

Abstract

Práce zkoumá vztah tzv. principů reflexe a velkých kardinálů. Lévy ukázal, že v ZFC platí tzv. věta o reflexi a dokonce, že věta o reflexi je ekvivalentní schématu nahrazení a axiomu nekonečna nad teorií ZFC bez axiomu nekonečna a schématu nahrazení. Tedy lze na větu o reflexi pohlížet jako na svého druhu axiom nekonečna. Práce zkoumá do jaké míry a jakým způsobem lze větu o reflexi zobecnit a jaký to má vliv na existenci tzv. velkých kardinálů. Práce definuje nedosažitelné, Mahlovy a nepopsatelné kardinály a ukáže, jak je lze zavést pomocí reflexe. Přirozenou limitou kardinálů získaných reflexí jsou kardinály nekonzistentní s L. Práce nabídne intuitivní zdůvodnění, proč tomu tak je.

Abstract

This thesis aims to examine relations between the so called Reflection Principles and Large cardinals. Lévy has shown that the Reflection Theorem is a sound theorem of ZF and it is equivalent to the Replacement Scheme and the Axiom of Infinity. From this point of view, Reflection theorem can be seen a specific version of an Axiom of Infinity. This paper aims to examine the Reflection Principle and its generalisations with respect to the existence of Large Cardinals. This thesis will establish the Inaccessible, Mahlo and Indescribable cardinals and show how can those be defined via reflection. A natural limit of Large Cardinals obtained via reflection are cardinals inconsistent with L. This thesis will offer an intuitive explanation of why this holds.

Contents

39	Contents	
40	1 Introduction	4
41	1.1 Motivation and Origin	4
42	1.2 Notation and Terminology	4
43	1.2.1 The Language of Set Theory	4
44	1.2.2 The Axioms	5
45	1.2.3 The Transitive Universe	8
46	1.2.4 Cardinal Numbers	10
47	1.2.5 Relativisation and Absoluteness	12
48	1.2.6 More Functions	14
49	1.2.7 Structure, Substructure and Embedding	15
50	2 Levy's First-Order Reflection	17
51	2.1 Lévy's Original Paper	17
52	2.2 Contemporary Restatement	21
53	3 Reflection And Large Cardinals	29
54	3.1 Regular Fixed-Point Axioms	29
55	3.2 Inaccessible Cardinal	32
56	3.3 Mahlo Cardinals	34
57	3.4 Indescribable Cardinals	37
58	4 Conclusion	41

1 Introduction

1.1 Motivation and Origin

“The Universe of sets cannot be uniquely characterised (i. e. distinguished from all its initial elements) by any internal structural property of the membership relation in it, which is expressible in any logic of finite or transfinite type, including infinitary logics of any cardinal order.”

— Kurt Gödel [Wang, 1997]

1.2 Notation and Terminology

1.2.1 The Language of Set Theory

This text assumes the knowledge of basic terminology and some results from first-order predicate logic, for example [Hamilton, 1988]. We won’t introduce the notions of *language*, *function symbol*, *predicate*, *term*, *model* and *interpretation* that are used in (1.42).

All proofs are based on [Jech, 2006] unless explicitly stated otherwise. Notable amount of inspiration is also drawn from [Kanamori, 2003] and [Drake, 1974].

We will now shortly review the basic notions that allow us to define the *Zermelo-Fraenkel* set theory.

When we talk about a *class*, we have the notion of a definable class in mind. If $\varphi(x, p_1, \dots, p_n)$ is a formula in the language of set theory, we call

$$A = \{x : \varphi(x, p_1, \dots, p_n)\} \quad (1.1)$$

a class of all sets satisfying $\varphi(x, p_1, \dots, p_n)$ in a sense that

$$x \in A \leftrightarrow \varphi(x, p_1, \dots, p_n) \quad (1.2)$$

Given classes A, B , one can easily define the elementary set operations such as $A \cap B$, $A \cup B$, $A \setminus B$, $\bigcup A$, see the first part of [Jech, 2006] for details. Axioms are the tools by which we can decide whether a particular class is “small enough” to be considered a set¹. A class that fails to be considered a set is called a *proper class*.

We will often write something like “ M is a limit ordinal”, it should always be clear that this can be rewritten as a formula that was introduced earlier.

¹“Small enough” means that it doesn’t introduce a paradox similar to Russell’s.

88 **1.2.2 The Axioms**89 **Definition 1.1** (*The Existence of a Set*)

$$\exists x(x = x) \quad (1.3)$$

90 **Definition 1.2** (*Axiom of Extensionality*)

$$\forall x, y(x = y \leftrightarrow \forall z(z \in x \leftrightarrow z \in y)) \quad (1.4)$$

91 **Definition 1.3** (*Axiom Schema of Specification*)

92 *The following yields an axiom for every first-order formula $\varphi(x, p_1, \dots, p_n)$*
 93 *with no free variables other than x, p_1, \dots, p_n .*

$$\forall x, p_1, \dots, p_n \exists y \forall z(z \in y \leftrightarrow z \in x \ \& \ \varphi(z, p_1, \dots, p_n)) \quad (1.5)$$

94 We will now provide two definitions that are not axioms, but will be
 95 helpful in establishing some axioms in a more comprehensible way.

96 **Definition 1.4** ($x \subseteq y, x \subset y$)

$$x \subseteq y \leftrightarrow (\forall z \in x)z \in y \quad (1.6)$$

97

$$x \subset y \leftrightarrow x \subseteq y \ \& \ x \neq y \quad (1.7)$$

98 *We read $x \subseteq y$ as x is a subset of y and $x \subset y$ as x is a proper subset of y .*

99 **Definition 1.5** (*Empty Set*) *For an arbitrary set x , the empty set, repre-*
 100 *sented by the symbol " \emptyset ", is the set defined by the following formula:*

$$(\forall y \in x)(y \in \emptyset \leftrightarrow \neg(y = y)) \quad (1.8)$$

101 \emptyset is a set due to Specification, there is only one such set due to Extension-
 102 ality.

103 **Definition 1.6** (*Axiom of Pairing*)

$$\forall x, y \exists z \forall q(q \in z \leftrightarrow q = x \vee q = y) \quad (1.9)$$

104 **Definition 1.7** (*Axiom of Union*)

$$\forall x \exists y \forall z(z \in y \leftrightarrow \exists q(z \in q \ \& \ q \in x)) \quad (1.10)$$

105 Now we can introduce more axioms.

106 **Definition 1.8** (*Axiom of Foundation*)

$$\forall x(x \neq \emptyset \rightarrow (\exists y \in x)(x \cap y = \emptyset)) \quad (1.11)$$

107 **Definition 1.9** (*Axiom of Powerset*)

$$\forall x \exists y \forall z (z \in y \leftrightarrow z \subseteq x) \quad (1.12)$$

108 **Definition 1.10** (*Axiom of Infinity*)

$$\exists x(\emptyset \in x \ \& \ (\forall y \in x)(y \cup \{y\} \in x)) \quad (1.13)$$

109 The least set satisfying this is denoted “ ω ”.

110 **Definition 1.11** (*Function*)

111 Given arbitrary first-order formula $\varphi(x, y, p_1, \dots, p_n)$, we say that φ is a func-
112 tion iff

$$\forall x, y, z, p_1, \dots, p_n (\varphi(x, y, p_1, \dots, p_n) \ \& \ \varphi(x, z, p_1, \dots, p_n) \rightarrow y = z) \quad (1.14)$$

113 When a $\varphi(x, y)$ is a function, we also write the following:

$$f(x) = y \leftrightarrow \varphi(x, y) \quad (1.15)$$

114 Alternatively, $f = \{\langle x, y \rangle : \varphi(x, y)\}$ is a class.

115 Let us introduce a few more definitions that will make the two remaining
116 axioms more comprehensible.

117 **Definition 1.12** (*Powerset Function*)

118 Given a set x , the powerset of x , denoted $\mathcal{P}(x)$ and satisfying (1.9), is
119 defined as follows:

$$\mathcal{P}(x) \stackrel{\text{def}}{=} \{y : y \subseteq x\} \quad (1.16)$$

120 **Definition 1.13** (*Domain of a Function*)

121 Let f be a function. We call the domain of f the set of all sets for which f
122 is defined. We use “ $\text{Dom}(f)$ ” to refer to this set.

$$x \in \text{Dom}(f) \leftrightarrow \exists y(f(x) = y) \quad (1.17)$$

123 We say “ f is a function on A ”, A being a class, if $A = \text{dom}(f)$.

124 **Definition 1.14** (*Range of a Function*)

125 Let f be a function. We call the range of f the set of all sets that are images
126 of other sets via f . We use “ $\text{Rng}(f)$ ” to refer to this set.

$$x \in \text{Rng}(f) \leftrightarrow \exists y(f(y) = x) \quad (1.18)$$

127 We say that f is a *function into* A , A being a class, iff $\text{rng}(f) \subseteq A$. We say
 128 that f is a *function onto* A iff $\text{rng}(f) = A$. We say a function f is a *one to*
 129 *one function*, iff

$$(\forall x_1, x_2 \in \text{dom}(f))(f(x_1) = f(x_2) \rightarrow x_1 = x_2) \quad (1.19)$$

130 We say that f is a *bijection* iff it is a one to one function that is onto.

131 Note that $\text{Dom}(f)$ and $\text{Rng}(f)$ are not definitions in a strict sense, they
 132 are in fact definition schemas that yield definitions for every function f given.
 133 Also note that they can be easily modified for φ instead of f , with the only
 134 difference being the fact that it is then defined only for those φ s that are
 135 functions, which must be taken into account. This is worth noting as we will
 136 use the notions of *function* and *formula* interchangeably.

137 **Definition 1.15** (*Function Defined For All Ordinals*)

138 We say a function f is defined for all ordinals, this is sometimes written
 139 $f : \text{Ord} \rightarrow A$ for any class A , if $\text{Dom}(f) = \text{Ord}$. Alternatively,

$$(\forall \alpha \in \text{Ord})(\exists y \in A)(f(\alpha) = y) \quad (1.20)$$

140 And now for the axioms.

141 **Definition 1.16** (*Axiom Schema of Replacement*)

142 The following is an axiom for every first-order formula $\varphi(x, p_1, \dots, p_n)$ with
 143 no free variables other than x, p_1, \dots, p_n .

$$“\varphi \text{ is a function}” \rightarrow \forall x \exists y \forall z (z \in y \leftrightarrow (\exists q \in x)(\varphi(x, y, p_1, \dots, p_n))) \quad (1.21)$$

144 **Definition 1.17** (*Choice function*)

145 We say that a function f is a *choice function* on x iff

$$\text{dom}(f) = x \setminus \{\emptyset\} \ \& \ (\forall y \in \text{dom}(f))(f(y) \in y) \quad (1.22)$$

146 **Definition 1.18** (*Axiom of Choice*)

147 For every set x there is a function f that is a *choice function* on x .

148 One might be unsettled by the fact that this definition quantifies over func-
 149 tions, which are generally classes, but in this particular case, since $\text{dom}(f) =$
 150 x and x is a set, f is also a set due to *Replacement*².

151 We will refer to the axioms by their name, written in italic type, e.g.
 152 *Foundation* refers to the Axiom of Foundation. Now we need to define the
 153 set theories to be used in the article.

²If the underlying theory includes of implies *Replacement*.

154 **Definition 1.19** (S)

155 We call **S** an axiomatic theory in the language $\mathcal{L} = \{=, \in\}$ with exactly the
 156 following axioms:

- 157 (i) Existence of a set (see (1.1))
- 158 (ii) Extensionality (see (1.2))
- 159 (iii) Specification (see (1.3))
- 160 (iv) Foundation (see (1.8))
- 161 (v) Pairing (see (1.6))
- 162 (vi) Union (see (1.7))
- 163 (vii) Powerset (see (1.9))

164 **Definition 1.20** (ZF)

165 We call **ZF** an axiomatic theory in the language $\mathcal{L} = \{=, \in\}$ that contains
 166 all the axioms of **S** in addition to the following:

- 167 (i) Replacement schema (see (1.16))
- 168 (ii) Infinity (see (1.10))

169 Existence of a set is usually left out because it is a consequence of infinity.

170 **Definition 1.21** (ZFC)

171 **ZFC** is an axiomatic theory in the language $\mathcal{L} = \{=, \in\}$ that contains all the
 172 axioms of **ZF** plus Choice (1.18).

173

174 **1.2.3 The Transitive Universe**175 **Definition 1.22** (Transitive Class)

176 We say a class A is transitive iff

$$(\forall x \in A)(x \subseteq A) \quad (1.23)$$

177 **Definition 1.23** (Well Ordered Class) A class A is said to be well ordered
 178 by \in iff the following hold:

- 179 (i) $(\forall x \in A)(x \not\subseteq x)$ (Antireflexivity)
- 180 (ii) $(\forall x, y, z \in A)(x \in y \ \& \ y \in z \rightarrow x \in z)$ (Transitivity)
- 181 (iii) $(\forall x, y \in A)(x = y \vee x \in y \vee y \in x)$ (Linearity)
- 182 (iv) $(\forall x \subseteq A)(x \neq \emptyset \rightarrow (\exists y \in x)(\forall z \in x)(z = y \vee z \in y))$ (Existence of the
 183 least element)

184 **Definition 1.24** (Ordinal Number)

185 A set x is said to be an ordinal number if it is transitive and well-ordered
 186 by \in .

For the sake of brevity, we usually just say “ x is an *ordinal*”. Note that “ x is an ordinal” is a well-defined formula in the language of set theory, since 1.22 is a first-order formula and 1.23 is in fact a conjunction of four first-order formulas. Ordinals will be usually denoted by lower case greek letters, starting from the beginning of the alphabet: $\alpha, \beta, \gamma, \dots$. Given two different ordinals α, β , we will write $\alpha < \beta$ for $\alpha \in \beta$, see Lemma 2.11 in [Jech, 2006] for technical details.

Definition 1.25 (*Non-Zero Ordinal*) We say an ordinal α is non-zero iff $\alpha \neq \emptyset$.

Definition 1.26 (*Successor Ordinal*) Consider the following function defined for all ordinals. Let β be an arbitrary ordinal. We call S the successor function.

$$S(\beta) = \beta \cup \{\beta\} \quad (1.24)$$

An ordinal α is called a successor ordinal iff there is an ordinal β , such that $\alpha = S(\beta)$. We also write $\alpha = \beta + 1$.

Definition 1.27 (*Limit Ordinal*) A non-zero ordinal α is called a limit ordinal iff it is not a successor ordinal.

Definition 1.28 (*Ord*) The class of all ordinal numbers, which we will denote “ Ord ”³ is the proper class defined as follows.

$$x \in \text{Ord} \leftrightarrow x \text{ is an ordinal} \quad (1.25)$$

Definition 1.29 (*Von Neumann’s Hierarchy*) The Von Neumann’s Hierarchy is a collection of sets indexed by elements of Ord , defined recursively in the following way:

$$(i) \quad V_0 = \emptyset \quad (1.26)$$

$$(ii) \quad V_{\alpha+1} = \mathcal{P}(V_\alpha) \text{ for any ordinal } \alpha \quad (1.27)$$

$$(iii) \quad V_\lambda = \bigcup_{\beta < \lambda} V_\beta \text{ for a limit ordinal } \lambda \quad (1.28)$$

³Other authors use “ On ”, we will stick to the notation used in [Jech, 2006].

209 We will also refer to the Von Neumann's Hierarchy as Von Neumann's Uni-
 210 verse or the Cumulative Hierarchy. This definition is only correct in a theory
 211 that contains or implies Replacement because otherwise it's not clear that the
 212 successor step is a set.

213 **Definition 1.30** (*Rank*)

214 Given a set x , we say that the rank of x (written as $\text{rank}(x)$) is the least
 215 ordinal α such that $x \in V_{\alpha+1}$

216 Due to *Regularity*, every set has a rank.⁴ The Von Neumann's hierarchy
 217 defined above can also be defined by the fact that every V_α is a set of all set
 218 with rank less than α .

219 **Definition 1.31** (*Order-type*)

220 Given an arbitrary well-ordered set x , we say that an ordinal α is the order-
 221 type of x iff x and α are isomorphic.

222

223 1.2.4 Cardinal Numbers

224 **Definition 1.32** (*Cardinality*)

225 Given a set x , let the cardinality of x , written $|x|$, be defined as the smallest
 226 ordinal number such that there is a one to one mapping from x onto α .

227 **Definition 1.33** (*Aleph function*)

228 Let ω be the set defined by ???. We will recursively define the function \aleph for
 229 all ordinals.

230 (i) $\aleph_0 = \omega$

231 (ii) $\aleph_{\alpha+1}$ is the least cardinal larger than \aleph_α ⁵

232 (iii) $\aleph_\lambda = \bigcup_{\beta < \lambda} \aleph_\beta$ for a limit ordinal λ

233 If $\kappa = \aleph_\alpha$ and α is a successor ordinal, we call κ a successor cardinal. If α
 234 is a limit ordinal, we call κ a limit cardinal.

235 **Definition 1.34** (*Cardinal number*)

236

237 (i) A set x is called a finite cardinal iff $x \in \omega$.

238 (ii) A set is called an infinite cardinal iff there is an ordinal α such that

239 $\aleph_\alpha = x$

⁴See chapter 6 of [Jech, 2006] for details.

⁵"The least cardinal larger than \aleph_α " is sometimes notated as \aleph_α^+ .

(iii) A set is called a cardinal iff it is either a finite cardinal or an infinite cardinal.

We say κ is an uncountable cardinal iff it is an infinite ordinal and $\aleph_0 < \kappa$. Infinite cardinals will be notated by lowercase greek letters from the middle of the alphabet, e.g. κ, μ, ν, \dots with the exception of λ , which is next to κ in the greek alphabet, but is also sometimes used for limit ordinal.

For formal details as well as why every set can be well-ordered assuming *Choice*, and therefore has a cardinality, see [Jech, 2006].

Definition 1.35 (*Sequence*)

We say that a function $\varphi(x, y)$ is a sequence iff there is an ordinal α such that $\text{dom}(\varphi) = \alpha$. In other words, a function is called a sequence if it is defined exactly for every ordinal from 0 to some α . We then say it is an α -sequence. We usually write $\langle \beta_i : i \in \alpha \rangle$ or $\langle \beta_0, \beta_1, \dots \rangle$ when referring to a sequence, β_i then denotes the elements of $\text{rng}(\varphi)$ for every $i \in \text{dom}(\varphi)$.

Definition 1.36 (*Cofinal Subset*)

Given a class A of ordinals, we say that $B \subseteq A$ is cofinal in A iff

$$(\forall x \in A)(\exists y \in B)(x \in y) \quad (1.29)$$

In other words, B is cofinal in A iff it is unbounded in A .

Definition 1.37 (*Cofinality of a Limit Ordinal*)

Let λ be a limit ordinal. We say that the cofinality of λ is κ iff κ is the least ordinal, such that there is a cofinal κ -sequence $\langle \beta_\xi : \xi < \kappa \rangle$, such that

$$\sup(\{\beta_\xi : \xi < \kappa\}) = \lambda \quad (1.30)$$

We write $cf(\lambda) = \kappa$.

Note that $cf(\alpha)$ is always a cardinal⁶.

Definition 1.38 (*Regular Cardinal*)

We say an infinite cardinal κ is regular iff $cf(\kappa) = \kappa$.

Definition 1.39 (*Strong Limit Cardinal*)

We say that an ordinal κ is a strong limit cardinal if it is a limit cardinal and

$$(\forall \alpha \in \kappa)(|\mathcal{P}(\alpha)| \in \kappa). \quad (1.31)$$

⁶If $cf(\alpha)$ is not a cardinal, so $|cf(\alpha)| < cf(\alpha)$, then there is a mapping from $|cf(\alpha)|$ onto $cf(\alpha)$. But then the range of this mapping is a cofinal subset of $cf(\alpha)$ that is strictly smaller than $cf(\alpha)$.

267 **Definition 1.40** (*Generalised Continuum Hypothesis*)

268

$$(\forall \alpha \in \text{Ord}) \aleph_{\alpha+1} = |\mathcal{P}(\aleph_\alpha)| \quad (1.32)$$

269 If *GCH* holds (for example in Gödel's L , see chapter 3), the notions of limit
270 cardinal and strong limit cardinal are equivalent.

271

272 1.2.5 Relativisation and Absoluteness

273 **Definition 1.41** (*Relativization*)

274 Let M be a class, $R \subseteq M \times M$ and let $\varphi(p_1, \dots, p_n)$ be a first-order formula
275 with no free variables besides p_1, \dots, p_n . The relativization of φ to M and R
276 is the formula, written as $\varphi^{M,R}$, defined in the following inductive manner:

- 277 (i) $(x \in y)^{M,R} \leftrightarrow R(x, y)$
- 278 (ii) $(x = y)^{M,R} \leftrightarrow x = y$
- 279 (iii) $(\neg \varphi)^{M,R} \leftrightarrow \neg \varphi^{M,R}$
- 280 (iv) $(\varphi \ \& \ \psi)^{M,R} \leftrightarrow \varphi^{M,R} \ \& \ \psi^{M,R}$
- 281 (v) $(\varphi \vee \psi)^{M,R} \leftrightarrow \varphi^{M,R} \vee \psi^{M,R}$
- 282 (vi) $(\varphi \rightarrow \psi)^{M,R} \leftrightarrow \varphi^{M,R} \rightarrow \psi^{M,R}$
- 283 (vii) $(\exists x \varphi(x))^{M,R} \leftrightarrow (\exists x \in M) \varphi^{M,R}(x)$
- 284 (viii) $(\forall x \varphi(x))^{M,R} \leftrightarrow (\forall x \in M) \varphi^{M,R}(x)$

285 When $R = \in \cap (M \times M)$, we usually write φ^M instead of $\varphi^{M,R}$. When we
286 talk about $\varphi^M(p_1, \dots, p_n)$, it is understood that $p_1, \dots, p_n \in M$.

287 **Definition 1.42** (*Satisfaction in a Structure*)

288 Let M be a set and R a binary relation on M . Let *Terms* be the set of all
289 terms, let $e : \text{Terms} \rightarrow M$ any evaluation function. Let φ be a first-order
290 formula in the language of set theory.

291 iff any of the following hold

292 We say that φ holds in $\langle M, R \rangle$ under the evaluation e , we write $\langle M, R \rangle \models \varphi[e]$,

293 iff any of the following hold

- 294 (i) φ is the formula " $s = t$ ", s, t are terms and $e(s) = e(t)$.
- 295 (ii) φ is the formula " $s \in t$ ", s, t are terms and the pair $\langle e(s), e(t) \rangle$ is in R .
- 296 (iii) φ is the formula " $\neg \psi$ " and not $\langle M, R \rangle \models \psi[e]$
- 297 (iv) φ is the formula " $\psi_1 \ \& \ \psi_2$ " and both $\langle M, R \rangle \models \psi_1[e]$ and $\langle M, R \rangle \models \psi_2[e]$.
- 298 (v) φ is the formula " $\psi_1 \vee \psi_2$ " and either $\langle M, R \rangle \models \psi_1[e]$ or $\langle M, R \rangle \models \psi_2[e]$.
- 299 (vi) φ is the formula " $\psi_1 \rightarrow \psi_2$ " and either not $\langle M, R \rangle \models \psi_1[e]$ or
300 $\langle M, R \rangle \models \psi_2[e]$.
- 301 (vii) φ is the formula " $\psi_1 \rightarrow \psi_2$ " and either not $\langle M, R \rangle \models \psi_1[e]$ or
302 $\langle M, R \rangle \models \psi_2[e]$.

- 303 (viii) φ is the formula " $\forall x_1 \psi$ " and $\langle M, R \rangle \models \psi[e']$ for every e' that differs
 304 from e only in the value of x_1 .
 305 (ix) φ is the formula " $\forall x_1 \psi$ " and $\langle M, R \rangle \models \psi[e']$ for every e' that differs
 306 from e only in the value of x_1 .
 307 (x) φ is the formula " $\exists x_1 \psi$ " and $\langle M, R \rangle \models \psi[e']$ for some e' that differs
 308 from e only in the value of x_1 .
 309 We also write $\langle M, R \rangle \models \varphi$, which

310 Note that we say that M is a set.

311 We will use $\langle M, R \rangle \models \varphi(p_1, \dots, p_n)$ and $\varphi^M(p_1, \dots, p_n)$ interchangeably.

312 **Definition 1.43** (Absoluteness) Given a transitive class M , we say a for-
 313 mula φ is absolute in M if for all $p_1, \dots, p_n \in M$

$$\varphi^M(p_1, \dots, p_n) \leftrightarrow \varphi(p_1, \dots, p_n) \quad (1.33)$$

314 **Definition 1.44** (Hierarchy of First-Order Formulas)

- 315
 316 (I) A first-order formula φ is Δ_0 iff it is logically equivalent to a first-order
 317 formula φ' satisfying any of the following:
 318 (i) φ' contains no quantifiers
 319 (ii) y is a set, ψ is a Δ_0 -formula, and φ' is either $(\exists x \in y)\psi(y)$ or
 320 $(\forall x \in y)\psi(y)$.
 321 (iii) ψ_1, ψ_2 are Δ_0 -formulas and φ' is any of the following: $\psi_1 \vee \psi_2$,
 322 $\psi_1 \& \psi_2$, $\psi_1 \rightarrow \psi_2$, $\neg \psi_2$,
 323 (II) If a formula is Δ_0 it is also Σ_0 and Π_0
 324 (III) A formula φ is $\Pi_n + 1$ if it is logically equivalent to a formula φ' such
 325 that $\varphi' = \forall x \psi$ where ψ is a Σ_n -formula for any $n < \omega$.
 326 (IV) A formula φ is $\Sigma_n + 1$ if it is logically equivalent to a formula φ' such
 327 that $\varphi' = \exists x \psi$ where ψ is a Π_n -formula for any $n < \omega$.

328 Note that we can use the pairing function so that for $\forall p_1, \dots, p_n \psi(p_1, \dots, p_n)$,
 329 there is a logically equivalent formula of the form $\forall x \psi'(x)$.

330 **Lemma 1.45** (Δ_0 absoluteness) Let φ be a Δ_0 -formula, then φ is absolute
 331 in any transitive class M .

332 *Proof.* This will be proven by induction over the complexity of a given Δ_0 -
 333 formula φ . Let M be an arbitrary transitive class.

334 Atomic formulas are always absolute by the definition of relativisation,
 335 see (1.41). Suppose that Δ_0 -formulas ψ_1 and ψ_2 are absolute in M . Then

from relativization, $(\psi_1 \& \psi_2)^M \leftrightarrow \psi_1^M \& \psi_2^M$, which is, from the induction hypothesis, equivalent to $\psi_1 \& \psi_2$. The same holds for \vee, \rightarrow, \neg .

Suppose that a Δ_0 -formula ψ is absolute in M . Let y be a set and let $\varphi = (\exists x \in y)\psi(x)$. From relativization, $(\exists x\psi(x))^M \leftrightarrow (\exists x \in M)\psi^M(x)$. Since the hypotheses makes it clear that $\psi^M \leftrightarrow \psi$, we get $((\exists x \in y)\psi(x))^M \leftrightarrow (\exists x \in y \cap M)\psi(x)$, which is the equivalent of $\varphi^M \leftrightarrow \varphi$. The same applies to $\varphi = (\forall x \in y)\psi(x)$. \square

Lemma 1.46 (*Downward Absoluteness*)

Let φ be a Π_1 -formula and M a transitive class. Then the following holds:

$$(\forall p_1, \dots, p_n \in M)(\varphi(p_1, \dots, p_n) \rightarrow \varphi(p_1, \dots, p_n)^M) \quad (1.34)$$

Proof. Since $\varphi(p_1, \dots, p_n)$ is Π_1 , there is a Δ_0 -formula $\psi(p_1, \dots, p_n, x)$ such that $\varphi = \forall x\psi(p_1, \dots, p_n, x)$. From relativization and lemma (1.45), $\varphi^M(p_1, \dots, p_n) \leftrightarrow (\forall x \in M)\psi(p_1, \dots, p_n, x)$.

Assume that for $p_1, \dots, p_n \in M$ fixed, that $\forall x\psi(p_1, \dots, p_n, x)$ holds, but $(\forall x \in M)\psi(p_1, \dots, p_n, x)$ does not. Therefore $\exists x\neg\psi(p_1, \dots, p_n, x)$, which contradicts $\forall x\psi(p_1, \dots, p_n, x)$. \square

Lemma 1.47 (*Upward Absoluteness*)

Let φ be a Σ_1 -formula and M a transitive class. Then the following holds:

$$(\forall p_1, \dots, p_n \in M)(\varphi^M(p_1, \dots, p_n) \rightarrow \varphi(p_1, \dots, p_n)) \quad (1.35)$$

Proof. Since $\varphi(p_1, \dots, p_n)$ is Σ_1 , there is a Δ_0 -formula $\psi(p_1, \dots, p_n, x)$ such that $\varphi = \exists x\psi(p_1, \dots, p_n, x)$. From relativization and lemma (1.45), $\varphi^M(p_1, \dots, p_n) \leftrightarrow (\exists x \in M)\psi(p_1, \dots, p_n, x)$.

Assume that for $p_1, \dots, p_n \in M$ fixed, that $(\exists x \in M)\psi(p_1, \dots, p_n, x)$ holds, but $\exists x\psi(p_1, \dots, p_n, x)$ does not. This is an obvious contradiction. \square

1.2.6 More Functions

Definition 1.48 (*Strictly Increasing Function*)

A function $f : Ord \rightarrow Ord$ is said to be strictly increasing iff

$$\forall \alpha, \beta \in Ord(\alpha < \beta \rightarrow f(\alpha) < f(\beta)). \quad (1.36)$$

Definition 1.49 (*Continuous Function*)

A function $f : Ord \rightarrow Ord$ is said to be continuous iff

$$\lambda \text{ is limit} \rightarrow f(\lambda) = \bigcup_{\alpha < \lambda} f(\alpha). \quad (1.37)$$

363 **Definition 1.50** (*Normal Function*)

364 A function $f : \text{Ord} \rightarrow \text{Ord}$ is said to be normal iff it is strictly increasing
365 and continuous.

366 **Definition 1.51** (*Fixed Point*)

367 We say x is a fixed point of a function f iff $x = f(x)$.

368 **Definition 1.52** (*Unbounded Class*)

369 We say a class A of ordinals is unbounded iff

$$\forall x(\exists y \in A)(x < y) \quad (1.38)$$

370 **Definition 1.53** (*Limit Point*)

371 Given a class $x \subseteq \text{Ord}$, we say that $\alpha \neq \emptyset$ is a limit point of x iff

$$\alpha = \bigcup (x \cap \alpha) \quad (1.39)$$

372 **Definition 1.54** (*Closed Class*)

373 We say a class $A \subseteq \text{Ord}$ is closed iff it contains all its limit points.

374 **Definition 1.55** (*Club set*)

375 For a regular uncountable cardinal κ , a set $x \subset \kappa$ is a closed unbounded
376 subset, abbreviated as a club set, iff x is both closed and unbounded in κ .

377 **Definition 1.56** (*Stationary set*)

378 For a regular uncountable cardinal κ , we say a set $A \subset \kappa$ is stationary in κ
379 iff it intersects every club subset of κ .

380 1.2.7 Structure, Substructure and Embedding

381 Structures will be denoted $\langle M, \in, R \rangle$ where M is a domain, \in stands for the
382 standard membership relation, it is assumed to be restricted to the domain⁷,
383 $R \subseteq M$ is a relation on the domain. When R is not needed, we can as well
384 only write M instead of $\langle M, \in \rangle$.

385 **Definition 1.57** (*Elementary Embedding*)

386 Given the structures $\langle M_0, \in, R \rangle$, $\langle M_1, \in, R \rangle$ and a one-to-one function $j :$
387 $M_0 \rightarrow M_1$, we say j is an elementary embedding of M_0 into M_1 , we write
388 $j : M_0 \prec M_1$, when the following holds for every formula $\varphi(p_1, \dots, p_n)$ and
389 every $p_1, \dots, p_n \in M_0$:

$$\langle M_0, \in, R \rangle \models \varphi(p_1, \dots, p_n) \leftrightarrow \langle M_1, \in, R \rangle \models \varphi(j(p_1), \dots, j(p_n)) \quad (1.40)$$

⁷To be totally explicit, we should write $\langle M, \in \cap M \times M, R \rangle$.

390 **Definition 1.58** (*Elementary Substructure*)

391 *Given the structures $\langle M_0, \in, R \rangle$, $\langle M_1, \in, R \rangle$ and a one-to-one function $j :$
 392 $M_0 \rightarrow M_1$ such that $j : M_0 \prec M_1$, we say that M_0 is an elementary sub-
 393 structure of M_1 , denoted as $M_0 \prec M_1$, iff j is an identity on M_0 . In other
 394 words*

$$\langle M_0, \in, R \rangle \models \varphi(p_1, \dots, p_n) \leftrightarrow \langle M_1, \in, R \rangle \models \varphi(p_1, \dots, p_n) \quad (1.41)$$

395 *for $p_1, \dots, p_n \in M_0$*

2 Levy's First-Order Reflection

2.1 Lévy's Original Paper

This section is based on Lévy's paper *Axiom Schemata of Strong Infinity in Axiomatic Set Theory*, [Lévy, 1960]. It presents Lévy's general reflection principle and its equivalence to *Replacement* and *Infinity* under S^8 .

First, we should point out that set theory has changed over the last 66 years and show a few notable, albeit only formal, differences. One might be confused by the fact that Lévy treats the *Subsets* axiom, which we call *Specification*, as a single axiom rather than a schema. He even takes the conjunction of all axioms of ZF and treats it like a formula. This is possible because the underlying logic calculus is different. Lévy works with set theories formulated in the *non-simple applied first order functional calculus*, see beginning of *Chapter IV* in [Church, 1996] for details. For now, we only need to know that the calculus contains a substitution rule for functional variables. This way, *Subsets* is de facto a schema even though it sometimes treated as a single formula⁹ but the logic is still first-order since one can't quantify over functional variables. We will use the usual first-order axiomatization of ZFC as seen on [Jech, 2006]. It should also be noted that the logical connectives look different. The now usual symbol for an universal quantifier does not appear, $\forall x\varphi(x)$ would be written as $(x)\varphi(x)$. The symbol for negation is " \sim ", implication is written as " \supset " and equivalence is " \equiv ". We will use standard notation with " \neg ", " \rightarrow " and " \leftrightarrow " respectively when presenting Lévy's results.

This subsection uses ZF instead of the usual ZFC as the underlying theory.

Definition 2.1 (Standard Complete Model of a Set Theory)

Let Q be an arbitrary axiomatic set theory. We say that u is a standard complete model of Q iff

- (i) $(\forall \sigma \in Q)(\langle u, \in \rangle \models \sigma)$
- (ii) $\forall y(y \in u \rightarrow y \subset u)$ (u is transitive)

We write $Scm^Q(u)$.

Definition 2.2 (Cardinals Inaccessible With Respect to Q)

Let Q be an arbitrary axiomatic set theory. We say that a cardinal κ is inaccessible with respect to theory Q iff

$$Scm^Q(V_\kappa) \quad (2.42)$$

⁸See definition (1.19).

⁹This way, the conjunction of all axioms is then in fact an axiom schema.

429 We write $In^Q(\kappa)$.¹⁰

430 **Definition 2.3** (*Inaccessible Cardinal With Respect to ZF*)

431 When a cardinal κ is inaccessible with respect to ZF, we only say that it is
432 inaccessible. We write $In(\kappa)$.

$$In(\kappa) \leftrightarrow In^{ZF}(\kappa) \quad (2.43)$$

433 The above definition of inaccessibles is used because it doesn't require *Choice*.

434 For the definition of relativization, see (1.41). The notation used by Lévy is
435 " $Rel(u, \varphi)$ ", we will stick to " φ^u ".

436 **Definition 2.4** (N)

437 The following is an axiom schema of complete reflection over ZF, denoted N .
438 For every first-order formula φ in the language of set theory with no free variables
439 except for p_1, \dots, p_n , the following is an instance of schema N .

$$\exists u(Scm^{ZF}(u) \ \& \ (\forall p_1, \dots, p_n \in u)(\varphi \leftrightarrow \varphi^u)) \quad (2.44)$$

440 **Definition 2.5** (N')

441 For any first-order formulas $\varphi_1, \dots, \varphi_m$ in the language of set theory with no
442 free variables except for p_1, \dots, p_n , the following is an instance of schema N' .

$$\exists u(z \in u \ \& \ Scm^{ZF}(u) \ \& \ (\forall p_1, \dots, p_n \in u)(\varphi_1 \leftrightarrow \varphi_1^u) \ \& \ \dots \ \& \ \varphi_m \leftrightarrow \varphi_m^u)) \quad (2.45)$$

443 **Definition 2.6** (N')

444 For any first-order formulas $\varphi_1, \dots, \varphi_m$ in the language of set theory with no
445 free variables except for p_1, \dots, p_n , the following is an instance of schema N' .

$$\exists u(Scm^{ZF}(u) \ \& \ (\forall p_1, \dots, p_n \in u)(\varphi_1 \leftrightarrow \varphi_1^u) \ \& \ \dots \ \& \ \varphi_m \leftrightarrow \varphi_m^u)) \quad (2.46)$$

446 Let S be an axiomatic set theory defined in (1.19).

447 This is *Theorem 2* in [Lévy, 1960]

448 **Lemma 2.7** ($N \leftrightarrow N'' \leftrightarrow N'$)

449 The schemas N , N' and N'' are equivalent under S .

¹⁰To be able to define V_κ , we need to work in a logic that contains the *Replacement Schema* or any of it's equivalents. It should be noted that we don't work in an arbitrary theory Q , but in ZF, which contains the *Replacement Schema*. $Scm^Q(V_\kappa)$ in fact says "ZF thinks that V_κ is a transitive model of Q ".

450 *Proof.* We will execute this proof in the theory ZF, but the reader should note
 451 that we are neither using *Replacement* nor *Infinity*, so for schemas similar to N ,
 452 N' , N'' but with " $Scm^S(u)$ " instead of " $Scm^{ZF}(u)$ ", the proof works equally
 453 well.

454 Clearly, $N' \rightarrow N'' \rightarrow N$.

455 Now, assuming N and given the formulas $\varphi_1, \dots, \varphi_n$, we will prove N'' .
 456 Consider the following formula:

$$\psi = \bigvee_{i=1}^t t = i \ \& \ \varphi_i \quad (2.47)$$

457 We will take advantage of the fact that natural numbers are defined by atomic
 458 formulas and therefore absolute in transitive structures. From N , we get such
 459 u that $Scm^{ZF}(u) \ \& \ (\forall p_1, \dots, p_n \in u)(\bigvee_{i=1}^t t = i \ \& \ \varphi_i \leftrightarrow \bigvee_{i=1}^t t = i \ \& \ \varphi_i^u)$. This
 460 already satisfies N'' .

461 In order to prove N' from N'' , let's add two more formulas. Given p_1, \dots, p_n ,
 462 we denote

$$\varphi_{m+1} = \exists u(z \in u \ \& \ Scm^{ZF}(u) \ \& \ (\forall p_1, \dots, p_n \in u)(\bigvee_{i=1}^m \varphi_i = \varphi_i^u)) \quad (2.48)$$

$$\varphi_{m+2} = \forall z \varphi_{m+1} \quad (2.49)$$

464 So, by N'' , we have a set u that satisfies $Scm^{ZF}(u)$ as well as the following:

$$(\forall p_1, \dots, p_n \in u)(\varphi_i \leftrightarrow \varphi_i^u) \text{ for } 1 \leq i \leq m \quad (2.50)$$

$$z \in u \rightarrow \varphi_{m+1} \leftrightarrow \varphi_{m+1}^u \quad (2.51)$$

$$\varphi_{m+2} \leftrightarrow \varphi_{m+2}^u \quad (2.52)$$

467 By $Scm^{ZF}(u)$ and (2.50), we get $(\forall z \in u)\varphi_{m+1}$, so together with (2.51), we get
 468 $(\forall z \in u)\varphi_{m+1}^u$, exactly φ_{m+2}^u , so by (2.52) we get φ_{m+2} . But φ_{m+2} is exactly
 469 the instance of N' we were looking for. \square

470 Definition 2.8 (N_0)

471 Axiom schema N_0 is similar to N defined above, but with S instead of ZF. For
 472 every φ , a first-order formula in the language of set theory with no free variables
 473 except p_1, \dots, p_n , the following is an instance of N_0 .

$$\exists u(Scm^S(u) \ \& \ (\forall p_1, \dots, p_n \in u)(\varphi \leftrightarrow \varphi^u)) \quad (2.53)$$

474 We will now show that in S , N_0 implies both *Replacement* and *Infinity*.

475
476 Let N_0 be defined as in (2.8), for *Infinity* see (1.10).

477 **Theorem 2.9** *In S , the axiom schema N_0 implies Infinity.*

478 *Proof.* Let $\varphi = \forall x \exists y (y = x \cup \{x\})$. This clearly holds in S because given a
479 set x , there is a set $y = x \cup \{x\}$ obtained via *Pairing* and *Union*. From N_0 ,
480 there is a set u such that φ^u holds. This u satisfies the conditions required by
481 *Infinity*. \square

482 Lévy proves this theorem in a different way. He argues that for an arbitrary
483 formula φ , N_0 gives us $\exists u Scm^S(u)$ and this u already satisfies *Infinity*. To do
484 this, we would need to prove lemma (2.15) earlier on, we will do that later in
485 this chapter.

486
487 Let S be a set theory defined in (1.19), N_0 a schema defined in (2.8) and
488 *Replacement* a schema defined in (1.16).

489 **Theorem 2.10** *In S , the axiom schema N_0 implies Replacement.*

490 *Proof.* Let $\varphi(x, y, p_1, \dots, p_n)$ be a formula with no free variables except x, y, p_1, \dots, p_n .
491 Let a set x be given and let χ be an instance of the *Replacement* schema for
492 the φ given. We want to verify in S that given a formula φ , the instance of N_0
493 for φ implies χ .

$$\begin{aligned} \chi = \forall x', y', z (\varphi(x', y', p_1, \dots, p_n) \ \& \ \varphi(x', z, p_1, \dots, p_n) \rightarrow y' = z') \\ \rightarrow \exists y \forall z (z \in y \leftrightarrow (\exists q \in x) (\varphi(x, y, p_1, \dots, p_n))) \end{aligned} \quad (2.54)$$

494 Since it can be shown that N_0 is equivalent to N'_0 similar to N' in lemma
495 (2.7), there is a set u such that $Scm^S(u)$, $x \in u$ and all of the following hold:

- 496 (i) $\varphi \leftrightarrow \varphi^u$
497 (ii) $\exists y \varphi \leftrightarrow (\exists y \varphi)^u$

498 From relativization, $(\exists y \varphi)^u$ is equivalent to $(\exists y \in u) \varphi^u$, together with (i) and
499 (ii), we get

$$(\exists y \in u) \varphi \leftrightarrow \exists y \varphi \quad (2.55)$$

500 If φ is a function, it maps the elements of x , which are also elements of u
501 due to transitivity of u , to elements of u . From *Specification*, $y = \{z \in u\}$ we
502 can find y , a set of all images of the elements of x via φ . So we have satisfied
503 *Replacement* – given a function and a set, we have proven that image of the set
504 via given function is also a set. \square

505 What we have just proven is only a single theorem from Lévy's aforementioned
506 article, we will introduce other interesting results, dealing with inaccessible and
507 Mahlo cardinals, later in their appropriate context in chapter 3.

2.2 Contemporary Restatement

We will now introduce and prove a theorem that is called Lévy's Reflection in contemporary set theory. The only difference is that while Lévy originally reflects a formula φ from V to a set u which is a *standard complete model* of S , we say that there is a V_λ for a limit λ that reflects φ . Those two conditions are equivalent due to lemma (2.15).

Lemma 2.11 *Let $\varphi_1, \dots, \varphi_n$ be first-order formulas in the language of set theory, all with m free variables¹¹.*

- (i) *For each set M_0 there is such set M that $M_0 \subset M$ and the following holds for every i , $1 \leq i \leq n$:*

$$\exists x \varphi_i(p_1, \dots, p_{m-1}, x) \rightarrow (\exists x \in M) \varphi_i(p_1, \dots, p_{m-1}, x) \quad (2.56)$$

for every $p_1, \dots, p_{m-1} \in M$.

- (ii) *Furthermore, there is a limit ordinal λ such that $M_0 \subset V_\lambda$ and the following holds for each i , $1 \leq i \leq n$:*

$$\exists x \varphi_i(p_1, \dots, p_{m-1}, x) \rightarrow (\exists x \in V_\lambda) \varphi_i(p_1, \dots, p_{m-1}, x) \quad (2.57)$$

for every $p_1, \dots, p_{m-1} \in M$.

- (iii) *Assuming Choice, there is M , $M_0 \subset M$ such that (2.56) holds for every M , $i \leq n$ and $|M| \leq |M_0| \cdot \aleph_0$.*

Proof. We will simultaneously prove statements (i) and (ii), denoting M^T the transitive set required by part (ii). Steps in the construction of M^T that are not explicitly included are equivalent to steps for M .

Let us first define an operation $H_i(p_1, \dots, p_{m-1})$ that yields the set of x 's with minimal rank¹² satisfying $\varphi_i(p_1, \dots, p_{m-1}, x)$ for p_1, \dots, p_{m-1} and for every i , $1 \leq i \leq n$.

$$H_i(p_1, \dots, p_n) = \{x \in C_i : (\forall z \in C)(\text{rank}(x) \leq \text{rank}(z))\} \quad (2.58)$$

for each $1 \leq i \leq n$, where

$$C_i = \{x : \varphi_i(p_1, \dots, p_{m-1}, x)\} \text{ for } 1 \leq i \leq n \quad (2.59)$$

¹¹For formulas with a different number of free variables, take for m the highest number of parameters among those formulas. Add spare parameters to every formula that has less than m parameters in a way that preserves the last parameter, which we will denote x . E.g. let φ'_i be the a formula with k parameters, $k < m$. Let us set $\varphi_i(p_1, \dots, p_{m-1}, x) = \varphi'_i(p_1, \dots, p_{k-1}, x)$, notice that the parameters p_k, \dots, p_{m-1} are not used.

¹²Rank is defined in (1.30).

531

532

Next, let's construct M from given M_0 by induction.

$$M_{i+1} = M_i \cup \bigcup_{j=0}^n \{H_j(p_1, \dots, p_{m-1}) : p_1, \dots, p_{m-1} \in M_i\} \quad (2.60)$$

533

534

535

536

537

In other words, in each step we include into the construction the elements satisfying $\varphi(p_1, \dots, p_{m-1}, x)$ for p_1, \dots, p_{m-1} from the previous step. For statement (ii), this is the only part that differs from (i). To end up with a transitive M , we need to extend every step to it's transitive closure transitive closure of M_{i+1} from (i). In other words, let γ be the smallest ordinal such that

$$(M_i^T \cup \bigcup_{j=0}^n \{\bigcup \{H_j(p_1, \dots, p_{m-1}) : p_1, \dots, p_{m-1} \in M_i\}\}) \subset V_\gamma \quad (2.61)$$

538

Then the incremental step is

$$M_{i+1}^T = V_\gamma \quad (2.62)$$

539

and the final M is obtained by joining the previous steps.

$$M = \bigcup_{i=0}^{\infty} M_i, \quad M^T = \bigcup_{i=0}^{\infty} M_i^T = V_\lambda \text{ for some limit } \lambda. \quad (2.63)$$

540

541

542

543

544

545

546

547

548

549

550

551

552

553

We have yet to finish part (iii). Let's try to construct a set M' that satisfies the same conditions like M but is kept as small as possible. Assuming the Axiom of Choice, we can modify the construction so that the cardinality of M' is at most $|M_0| \cdot \aleph_0$. Note that the size of M in the previous construction is determined by the size of M_0 and, most importantly, by the size of $H_i(p_1, \dots, p_{m-1})$ for every i , $1 \leq i \leq n$ in individual iterations of the construction. Since (i) only ensures the existence of an x that satisfies $\varphi_i(p_1, \dots, p_{m-1}, x)$ for any i , $1 \leq i \leq n$, we only need to add one x for every set of parameters but $H_i(u_1, \dots, u_{m-1})$ can be arbitrarily large. Let F be a choice function on $\mathcal{P}(M')$. Also let $h_i(p_1, \dots, p_{m-1}) = F(H_i(p_1, \dots, p_{m-1}))$ for i , where $1 \leq i \leq n$, which means that h is a function that outputs an x that satisfies $\varphi_i(p_1, \dots, p_{m-1}, x)$ for i such that $1 \leq i \leq n$ and has minimal rank among all such sets. The induction step needs to be redefined to

$$M'_{i+1} = M'_i \cup \bigcup_{j=0}^n \{h_j(p_1, \dots, p_{m-1}) : p_1, \dots, p_{m-1} \in M'_i\} \quad (2.64)$$

554

555

This way, the amount of elements added to M'_{i+1} in each step of the construction is the same as the amount of m -tuples of parameters that yielded elements not

556 included in M'_i . It is easy to see that if M_0 is finite, M' is countable because
 557 it was constructed as a countable union of sets that are themselves at most
 558 countable. If M_0 is countable or larger, the cardinality of M' is equal to the
 559 cardinality of M_0 .¹³ Therefore $|M'| \leq |M_0| \cdot \aleph_0$ \square

560 **Theorem 2.12** (*Lévy's first-order reflection theorem*)

561 Let $\varphi(p_1, \dots, p_n)$ be a first-order formula.

562 (i) For every set M_0 there exists a set M such that $M_0 \subset M$ and the following
 563 holds:

$$\varphi^M(p_1, \dots, p_n) \leftrightarrow \varphi(p_1, \dots, p_n) \quad (2.65)$$

564 for every $p_1, \dots, p_n \in M$.

565 (ii) For every set M_0 there is a transitive set M , $M_0 \subset M$ such that the
 566 following holds:

$$\varphi^M(p_1, \dots, p_n) \leftrightarrow \varphi(p_1, \dots, p_n) \quad (2.66)$$

567 for every $p_1, \dots, p_n \in M$.

568 (iii) For every set M_0 there is a limit ordinal λ such that $M_0 \subset V_\lambda$ and the
 569 following holds:

$$\varphi^{V_\lambda}(p_1, \dots, p_n) \leftrightarrow \varphi(p_1, \dots, p_n) \quad (2.67)$$

570 for every $p_1, \dots, p_n \in M$.

571 (iv) Assuming Choice, for every set M_0 there is M such that $M_0 \subset M$ and
 572 $|M| \leq |M_0| \cdot \aleph_0$ and the following holds:

$$\varphi^M(p_1, \dots, p_n) \leftrightarrow \varphi(p_1, \dots, p_n) \quad (2.68)$$

573 for every $p_1, \dots, p_n \in M$.

574 *Proof.* Let's now prove (i) for given φ via induction by complexity. We can safely
 575 assume that φ contains no quantifiers besides " \exists " and no logical connectives
 576 other than " \neg " and "&". Let $\varphi_1, \dots, \varphi_n$ be all subformulas of φ . Then there
 577 is a set M , obtained by the means of lemma (2.11), for all of the formulas
 578 $\varphi_1, \dots, \varphi_n$.

579 Let's first consider atomic formulas in the form of either $x_1 = x_2$ or $x_1 \in x_2$.
 580 It is clear from relativisation¹⁴ that (2.65) holds for both cases, $(x_1 = x_2)^M \leftrightarrow$
 581 $(x_1 = x_2)$ and $(x_1 \in x_2)^M \leftrightarrow (x_1 \in x_2)$.
 582

¹³It can not be smaller because $|M'_{i+1}| \geq |M'_i|$ for every i . It may not be significantly larger because the maximum of elements added is the number of n -tuples in M'_i , which is of the same cardinality as M'_i .

¹⁴See (1.41.). This only holds for relativization to $M, \in \cap M \times M$, not M, R for an arbitrary R .

583 We now want to verify the inductive step. First, take $\varphi = \neg\varphi'$. From
 584 relativization, we get $(\neg\varphi')^M \leftrightarrow \neg(\varphi'^M)$. Because the induction hypothesis tells
 585 us that $\varphi'^M \leftrightarrow \varphi'$, the following holds:

$$(\neg\varphi')^M \leftrightarrow \neg(\varphi'^M) \leftrightarrow \neg\varphi' \quad (2.69)$$

586 The same holds for $\varphi = \varphi_1 \& \varphi_2$. From the induction hypothesis, we know
 587 that $\varphi_1^M \leftrightarrow \varphi_1$ and $\varphi_2^M \leftrightarrow \varphi_2$, which together with relativization for formulas in
 588 the form of $\varphi_1 \& \varphi_2$ gives us

$$(\varphi_1 \& \varphi_2)^M \leftrightarrow \varphi_1^M \& \varphi_2^M \leftrightarrow \varphi_1 \& \varphi_2 \quad (2.70)$$

589 Let's now examine the case when $\varphi = \exists x\varphi'(p_1, \dots, p_n, x)$. The induction
 590 hypothesis tells us that $\varphi'^M(p_1, \dots, p_n, x) \leftrightarrow \varphi'(p_1, \dots, p_n, x)$, so, together with
 591 above lemma (2.11), the following holds:

$$\begin{aligned} & \varphi(p_1, \dots, p_n, x) \\ & \leftrightarrow \exists x\varphi'(p_1, \dots, p_n, x) \\ & \leftrightarrow (\exists x \in M)\varphi'(p_1, \dots, p_n, x) \\ & \leftrightarrow (\exists x \in M)\varphi'^M(p_1, \dots, p_n, x) \\ & \leftrightarrow (\exists x\varphi'(p_1, \dots, p_n, x))^M \\ & \leftrightarrow \varphi^M(p_1, \dots, p_n, x) \end{aligned} \quad (2.71)$$

592 Which is what we wanted to prove for part (i).

593

594 We now need to verify that the same holds for any finite number of formulas
 595 $\varphi_1, \dots, \varphi_n$. This has in fact been already done since lemma (2.11) gives us a set
 596 M for any finite amount of formulas and given M_0 . We can therefore find a set
 597 M for the union of all of their subformulas. When we obtain such M , it should
 598 be clear that it also reflects every formula in $\varphi_1, \dots, \varphi_n$.

599

600 Since V_λ is a transitive set, by proving (iii) we also satisfy (ii). To do so, we
 601 only need to look at part (ii) of lemma (2.11). All of the above proof also holds
 602 for $M = V_{\lambda \text{ambda}}$.

603 To finish part (iv), we take M of size $\leq |M_0| \cdot \aleph_0$, which exists due to part
 604 (iii) of lemma (2.11), the rest being identical. \square

605

606 Let S be a set theory defined in (1.19), for ZFC see definition (1.21).

607 The two following lemmas are based on [Drake, 1974][Chapter 3, Theorem
 608 1.2].

609 **Lemma 2.13** *If M is a transitive set, then $\langle M, \in \rangle \models$ Extensionality.*

610 *Proof.* Given a transitive set M , we want to show that the following holds.

$$\langle M, \in \rangle \models \forall x, y (x = y \leftrightarrow \forall z (z \in x \leftrightarrow z \in y)) \quad (2.72)$$

611 Given arbitrary $x, y \in M$, we want to prove that $\langle M, \in \rangle \models (x = y \leftrightarrow \forall z (z \in$
 612 $x \leftrightarrow z \in y))$. This is equivalent to $\langle M, \in \rangle \models x = y$ iff $\langle M, \in \rangle \models \forall z (z \in$
 613 $x \leftrightarrow z \in y)$, which is the same as $x = y$ iff $\langle M, \in \rangle \models \forall z (z \in x \leftrightarrow z \in y)$.

614 So all elements of x are also elements of y in M , and vice versa. Because M is
 615 transitive, all elements of x and y are in M , so $\langle M, \in \rangle \models \forall z (z \in x \leftrightarrow z \in y)$
 616 holds iff x and y contain the same elements and are therefore equal. \square

617 **Lemma 2.14** *If M is a transitive set, then $\langle M, \in \rangle \models$ Foundation.*

618 *Proof.* We want to prove the following:

$$\langle M, \in \rangle \models \forall x (x \neq \emptyset \rightarrow (\exists y \in x)(x \cap y = \emptyset)) \quad (2.73)$$

619 Given an arbitrary non-empty $x \in M$ let's show that $\langle M, \in \rangle \models (\exists y \in$
 620 $x)(x \cap y = \emptyset)$.

621 Because M is transitive, every element of x is an element of M . Take for
 622 y the element of x with the lowest rank¹⁵. It should be clear that there is no
 623 $z \in y$ such that $z \in x$, because then $\text{rank}(z) < \text{rank}(y)$, which would be a
 624 contradiction. \square

625 Let S be a set theory as defined in (1.19).

626 **Lemma 2.15** *The following holds for every λ .*

$$"\lambda \text{ is a limit ordinal}" \rightarrow \langle V_\lambda, \in \rangle \models S \quad (2.74)$$

627 *Proof.* Given an arbitrary limit ordinal λ , we will verify the axioms of S one by
 628 one.

629 (i) *The existence of a set* comes from the fact that V_λ is a non-empty set
 630 because limit ordinal is non-zero by definition.

631 (ii) *Extensionality* holds from (2.13).

632 (iii) *Foundation* holds from (2.14).

633 (iv) *Union*:

634 Given any $x \in V_\lambda$, we want verify that $y = \bigcup x$ is also in V_λ . Note that
 635 $y = \bigcup x$ is a Δ_0 -formula.

$$y = \bigcup x \leftrightarrow (\forall z \in y)(\exists q \in x)z \in q \ \& \ (\forall z \in x)(\forall q \in z)q \in y \quad (2.75)$$

636 So by lemma (1.45)

$$y = \bigcup x \leftrightarrow \langle V_\lambda, \in \rangle \models y = \bigcup x \quad (2.76)$$

¹⁵Rank is defined in (1.30).

637 (v) *Pairing*:

638 Given two sets $x, y \in V_\lambda$, we want to show that $z = \{x, y\}$ is also an
639 element of V_λ .

$$z = \{x, y\} \leftrightarrow x \in z \ \& \ y \in z \ \& \ (\forall q \in z)(q = x \vee q = y) \quad (2.77)$$

640 So $z = \{x, y\}$ is a Δ_0 -formula, and thus by lemma (1.45) it holds that

$$z = \{x, y\} \leftrightarrow \langle V_\lambda, \in \rangle \models z = \{x, y\} \quad (2.78)$$

641 (vi) *Powerset*:

642 Given any $x \in V_\lambda$, we want to make sure that $\mathcal{P}(x) \in V_\lambda$. Let $\varphi(y)$ denote
643 the formula $y \in \mathcal{P}(x) \leftrightarrow y \subset x$. according to definition of subset (1.4),
644 $y \subset x$ is Δ_0 , so for any given $x, y \in V_\lambda$, $y = \mathcal{P}(x) \leftrightarrow \langle V_\lambda, \in \rangle \models y =$
645 $\mathcal{P}(x)$. Because λ is limit and $\text{rank}(\mathcal{P}(x)) = \text{rank}(x) + 1$, if $\mathcal{P}(x) \in V_\lambda$
646 for every $x \in V_\lambda$.

647 (vii) *Specification*:

648 Given a first-order formula φ , we want to show the following:

$$\langle V_\lambda, \in \rangle \models \forall x, p_1, \dots, p_n, \exists y \forall z (z \in y \leftrightarrow z \in x \ \& \ \varphi(z, p_1, \dots, p_n)) \quad (2.79)$$

649 Given any x along with parameters p_1, \dots, p_n in V_λ , we set

$$y = \{z \in x : \varphi^{V_\lambda}(z, p_1, \dots, p_n)\} \quad (2.80)$$

650 From transitivity of V_λ and the fact that $y \subset x$ and $x \in V_\lambda$, we know that
651 $y \in V_\lambda$, so $\langle V_\lambda, \in \rangle \models \forall z (z \in y \leftrightarrow z \in x \ \& \ \varphi(z, p_1, \dots, p_n))$.
652 □

653 **Definition 2.16** (*First-Order Reflection Schema*)

654 For every first-order formula φ , the following is an axiom:

$$\forall M_0 \exists M (M_0 \subseteq M \ \& \ (\varphi(p_1, \dots, p_n) \leftrightarrow \varphi(p_1, \dots, p_n)^M)) \quad (2.81)$$

655 We will refer to this axiom schema as First-order reflection.

656 Let *Infinity* and *Replacement* be as defined in (1.10) and (1.16) respectively.

657 **Theorem 2.17** First-order reflection is equivalent to Infinity & Replacement
658 under S.

659 *Proof.* Since (2.12) already gives us one side of the implication, we are only
660 interested in showing the converse which we shall do in two parts:

661 *First-order reflection* \rightarrow *Infinity* This is done exactly like (2.9). We pick for
 662 φ the formula $(\forall y \in x)(y \cup \{y\} \in x)$, $M_0 = \{\emptyset\}$. From (2.16), there is a set M
 663 that satisfies φ , so there is an inductive set. We have picked M_0 so that $\emptyset \in M$
 664 obviously holds and M is the witness for

$$\exists x(\emptyset \in x \ \& \ (\forall y \in x)(y \cup \{y\} \in x)) \quad (2.82)$$

665 which is exactly (1.10).

666 *First-order reflection* \rightarrow *Replacement*

667 Let's first point out that while *First-order reflection* gives us a set for one
 668 formula, we can generalize it to hold for any finite number of formulas. We will
 669 show how is it done for two formulas, which is what we will use in this proof.
 670 Given two first-order formulas φ, ψ , we can suppose that there are formulas φ'
 671 and ψ' that are equivalent to φ and ψ respectively, but their free variables are
 672 different¹⁶. Let $\xi = \varphi \ \& \ \psi$, given any M_0 , we can find a M such that $\xi \leftrightarrow \xi^M$.
 673 It is easy to see that from relativisation, the following holds:
 674

$$\varphi \ \& \ \psi \leftrightarrow \varphi' \ \& \ \psi' \leftrightarrow \xi \leftrightarrow \xi^M \leftrightarrow (\varphi' \ \& \ \psi')^M \leftrightarrow \varphi'^M \ \& \ \psi'^M \leftrightarrow \varphi^M \ \& \ \psi^M \quad (2.83)$$

675 Now given a function $\varphi(x, y)$, we know from *First-order reflection* that for
 676 every M_0 , there is a set M such that $M_0 \subseteq M$ and both

$$(\forall x, y \in M)(\varphi(x, y) \leftrightarrow \varphi^M(x, y)) \quad (2.84)$$

677 and

$$(\forall x, y \in M)(\exists y \varphi(x, y) \leftrightarrow (\exists y \varphi(x, y))^M) \quad (2.85)$$

678 hold, the latter being equivalent to

$$(\forall x, y \in M)(\exists y \varphi(x, y) \leftrightarrow (\exists y \in M) \varphi^M(x, y)) \quad (2.86)$$

679 Therefore

$$(\forall x, y \in M)(\exists y \varphi(x, y) \leftrightarrow (\exists y \in M) \varphi(x, y)) \quad (2.87)$$

680 holds too. That means that we have a set M such that for every $x \in M$, if φ is
 681 defined for x , $(\exists y \in M) \varphi(x, y)$.

682 To show that *Replacement* holds for this particular φ , we need to verify that
 683 given a set M_0 , $M'_0 = \{y : (\exists x \in M_0) \varphi(x, y)\}$ is also a set. But since $M_0 \subseteq M$

¹⁶This is plausible since we can for example substitute all free variables in φ' for x_0, x_2, x_4, \dots and use x_1, x_3, x_5, \dots for free variables in ψ' , the resulting formulas will be equivalent.

684 and because given any $x \in M$, there is $y \in M$ satisfying $\varphi(x, y)$, the following
 685 is a set due to *Specification*:

$$M'_0 = \{y : (\exists x \in M_0)\varphi(x, y)\} = \{y \in M : (\exists x \in M_0)\varphi(x, y)\} \quad (2.88)$$

686

□

687

688 We have shown that *Reflection* for first-order formulas, *First-order reflection* is
 689 a theorem of ZFC. We have also shown that it can be used instead of the *Infinity*
 690 and *Replacement* scheme, but $\text{ZFC} + \text{First-order reflection}$ is a conservative
 691 extension of ZF. Besides being a starting point for more general and powerful
 692 statements, it can be used to show that ZF is not finitely axiomatizable. This
 693 follows from the fact that *Reflection* gives a model to any consistent finite set of
 694 formulas. So if $\varphi_1, \dots, \varphi_n$ would be the axioms of ZFC, *Reflection* would prove
 695 that every model of ZFC contains a smaller model of ZFC, which would in turn
 696 contradict the Second Gödel's Theorem¹⁷.

697

698 It is also worthwhile to note that, in a way, *Reflection* is dual to compactness.
 699 Compactness says that given a set of sentences, if every finite subset yields
 700 a model, so does the whole set. *Reflection*, on the other hand, says that while
 701 the whole set has no model in the underlying theory, every finite subset has a
 702 model.

702

703 Furthermore, *Reflection* can be used in ways similar to upward Löwenheim–Skolem
 704 theorem. Since *Reflection* extends any set M_0 into a model of given formulas
 705 $\varphi_1, \dots, \varphi_n$, we can choose the lower bound of the size of M by appropriately
 706 choosing M_0 .

706

707 In the next section, we will try to generalize *Reflection* in a way that tran-
 scends ZF and yields some large cardinals.

¹⁷See chapter ?? for further details.

3 Reflection And Large Cardinals

3.1 Regular Fixed-Point Axioms

Lévy's article mentions various schemata that are not instances of reflection per se, but deal with fixed points of normal ordinal functions. We will introduce them and show that they are equivalent to *First-Order Reflection*¹⁸.

Lemma 3.1 (*Fixed-point lemma for normal functions*)

Let f be a normal function defined for all ordinals¹⁹. Then all of the following hold:

- (i) $\forall \lambda (\text{"}\lambda \text{ is a limit ordinal"} \rightarrow \text{"}f(\lambda) \text{ is a limit ordinal"})$
- (ii) $\forall \alpha (\alpha \leq f(\alpha))$
- (iii) $\forall \alpha \exists \beta (\alpha < \beta \ \& \ f(\beta) = \beta)$
- (iv) *The fixed points of f form a closed unbounded class.*²⁰

Proof. Let f be a normal function defined for all ordinals.

(i) Suppose λ is a limit ordinal. For an arbitrary ordinal $\alpha < \lambda$, the fact that f is strictly increasing means that $f(\alpha) < f(\lambda)$ and for any ordinal β , satisfying $\alpha < \beta < \lambda$, $f(\alpha) < f(\beta) < f(\lambda)$. We know that there is such β from limitness of λ . Because f is continuous and λ is limit, $f(\lambda) = \bigcup_{\gamma < \lambda} f(\gamma)$. That means that if λ is limit, so is $f(\lambda)$.

(ii) This step will be proven using the transfinite induction. Since f is defined for all ordinals, there is an ordinal α such that $f(\emptyset) = \alpha$ and because \emptyset is the least ordinal, (ii) holds for \emptyset .

Suppose (ii) holds for some β from the induction hypothesis. It the holds for $\beta + 1$ because f is strictly increasing.

For a limit ordinal λ , suppose (ii) holds for every $\alpha < \lambda$. (i) implies that $f(\lambda)$ is also limit, so there is a strictly increasing κ -sequence $\langle \alpha_0, \alpha_1, \dots \rangle$ for some κ such that $\lambda = \bigcup_{i < \kappa} \alpha_i$. Because f is stricly increasing, the κ -sequence $\langle f(\alpha_0), f(\alpha_1), \dots \rangle$ is also strictly increasing, the induction hypothesis implies that $\alpha_i \leq f(\alpha_i)$ for each $i \leq \kappa$. Thus, $\lambda \leq f(\lambda)$.

(iii) For a given ordinal α , let there be an ω -sequence $\langle \alpha_0, \alpha_1, \dots \rangle$, such that $\alpha_0 = \alpha$ and $\alpha_{i+1} = f(\alpha_i)$ for each $i < \omega$. This sequence is stricly increasing because so is f . Now, there's a limit ordinal $\beta = \bigcup_{i < \omega} \alpha_i$, we want to show that this is the fixed point. So $f(\beta) = f(\bigcup_{i < \omega} \alpha_i) = \bigcup_{i < \omega} f(\alpha_i)$ because f is continuous. We have defined the above sequence so that $\beta, \bigcup_{i < \omega} f(\alpha_i) = \bigcup_{i < \omega} \alpha_{i+1}$, which means we are done, since $\bigcup_{i < \omega} \alpha_{i+1} = \bigcup_{i < \omega} \alpha_i = \beta$.

¹⁸For definition, see (2.16).

¹⁹For the definition of normal function, see (1.50).

²⁰See (??.) for the definition of closed class, (1.52) for the definition of unboundedness.

(iv) The class of fixed points of f is obviously unbounded by (iii). It remains to show that it is closed, this is based on [Drake, 1974], chapter 4. Let Y be a non-empty set of fixed points of f such that $\bigcup Y \notin Y$. Since f is defined on ordinals, Y is a set of ordinals, so $\bigcup Y$ is an ordinal because a supremum of a set of ordinals is an ordinal. $\bigcup Y$ is a limit ordinal. If it were a successor ordinal, suppose that $\alpha + 1 = \bigcup Y$, then $\alpha \in \bigcup Y$, which means that there is some x such that $\alpha \in x \in Y$. But the least such x is $\alpha + 1$, so $\bigcup Y \in Y$.
 Note that $\alpha < \bigcup Y$ iff $\exists \xi \in Y (\alpha < \xi)$. Since f is defined for all ordinals and $\bigcup Y$ is a limit ordinal, $f(\bigcup Y) = \bigcup_{\alpha \in Y} f(\alpha)$, but because Y is a set of fixed points of f , $f(\bigcup Y) = \bigcup_{\alpha \in Y} f(\alpha) = \bigcup Y$, so $\bigcup Y$ is also a limit point of Y .

□

Lemma 3.2 *Let α be a limit ordinal. Then the following hold:*

- (i) *If C is a club set in α , then there is an ordinal β and a normal function $f : \beta \rightarrow \alpha$ such that $\text{rng}(f) = C$. We say that f enumerates C .*
- (ii) *If β is an ordinal and f is a normal function such that $f : \beta \rightarrow \alpha$ and $\text{rng}(f)$ is unbounded in α , then $\text{rng}(f)$ is a closed unbounded set in α .*

This proof comes from (<http://euclid.colorado.edu/~monkd/m6730/gradsets09.pdf> TODO cite!) *Proof.*

- (i) Let β be the order-type²¹ of C , let f be the isomorphism from β onto C . Since $C \subseteq \alpha$, f is also an increasing function from β into α . In order to be continuous, let γ be a limit ordinal under β , let $\epsilon = \bigcup_{\delta < \gamma} f(\delta)$. We want to verify that $f(\gamma) = \epsilon$. Since ϵ is a limit ordinal, we only need to show that $C \cap \epsilon$ is unbounded in ϵ .
 Take $\zeta < \epsilon$. Then there is a $\delta < \gamma$ such that $\zeta < f(\delta)$. Since γ is limit, $\delta + 1 < \gamma$ and also $f(\delta + 1) < f(\gamma)$, we know that $f(\delta) \in C \cap \epsilon$. But that means that $C \cap \epsilon$ is unbounded in ϵ , so $\epsilon \in C$. We have also shown that ϵ is closed unbounded in the image of γ over f . Therefore, $f(\gamma) = \epsilon = \bigcup_{\delta < \gamma} f(\delta)$, so f is normal.

- (ii) TODO (potrebuj to?)

□ It

should be clear that while this lemma works with club subsets of an ordinal, we can formulate analogous statement for club classes, which then yields a normal function defined for all ordinals, with the only exception that there is no such β because f is a function from Ord to Ord .

²¹See definition (1.31).

778 **Definition 3.3** (Axiom Schema M_1)

779 “Every normal function defined for all ordinals has at least one inaccessible num-
780 ber in its range.”

781 Lévy uses “ M ” to refer to this axiom but since we also use “ M ” for sets and
782 models, for example in (2.16), we will call the above axiom “Axiom Schema M_1 ”
783 to avoid confusion.

784 Let $\varphi(x, y, p_1, \dots, p_n)$ be a first-order formula with no free variables besides
785 x, y, p_1, \dots, p_n . The following is equivalent to Axiom M_1 .

$$\begin{aligned} & \text{“}\varphi \text{ is a normal function”} \ \& \ \forall x(x \in \text{Ord} \rightarrow \exists y(\varphi(x, y, p_1, \dots, p_n))) \rightarrow \\ & \rightarrow \exists y(\exists x \varphi(x, y, p_1, \dots, p_n) \ \& \ cf(y) = y \ \& \ (\forall x \in \kappa)(\exists y \in \kappa)(x > y)) \end{aligned} \quad (3.89)$$

786 **Definition 3.4** (Axiom Schema M_2)

787 “Every normal function defined for all ordinals has at least one fixed point which
788 is inaccessible.”

789 **Definition 3.5** (Axiom Schema M_3)

790 “Every normal function defined for all ordinals has arbitrarily great fixed points
791 which are inaccessible.”

792 Similar axiom is proposed in [Drake, 1974].

793 **Definition 3.6** (Axiom Schema F)

794 “Every normal function has a regular fixed point.”

795 **Lemma 3.7** Let f be a normal function defined for all ordinals.

- 796 (i) There is a normal function g_1 defined for all ordinals that enumerates the
797 class $\{\alpha : f(\alpha) = \alpha \ \& \ \alpha \in \text{Ord}\}$.
798 (ii) There is a normal function g_2 defined for all ordinals that enumerates the
799 class $\{\lambda : \text{“}f(\lambda) \text{ is a strong limit cardinal.”}\}$.

800 *Proof.* We know that (ii) holds from lemma (3.1) and lemma (3.2).

801 For (i), It should be clear that there is no largest strong limit ordinal ν ,
802 because the limit of $\nu, \mathcal{P}(\nu), \mathcal{P}(\mathcal{P}(\nu)), \dots$ is again a limit ordinal. The class of
803 limit ordinals is closed because a limit of strong limit ordinals is clearly always a
804 strong limit ordinal. Let h be a function enumerating limit ordinals which exists
805 from lemma (3.2). Then $g_1(\alpha) = f(h(\alpha))$ for every ordinal α is normal and
806 defined for all ordinals. \square

807 The following is *Theorem 1* in [Lévy, 1960], the parts dealing with Axiom
808 Schema F come from [Drake, 1974].

809 **Theorem 3.8** *The following are all equivalent:*

- 810 (i) Axiom Schema M_1
- 811 (ii) Axiom Schema M_2
- 812 (iii) Axiom Schema M_3
- 813 (iv) Axiom Schema F

814 *Proof.* It is clear that *Axiom Schema M_3* is a stronger version of *Axiom Schema*
 815 *M_2* , which is in turn a stronger version of both *Axiom Schema M_1* and *Axiom*
 816 *Schema F_1* .

817 We will now prove that *Axiom Schema $F \rightarrow$ Axiom Schema M_2* . Lemma
 818 (3.7) tells us that given a normal function f defined for all ordinals, there is a
 819 normal function g_1 defined for all ordinals that enumerates the fixed-points of f .
 820 There is also a function g_2 that enumerates the strong limit ordinals in $rng(f)$.
 821 By *Axiom Schema F* , g_2 has a regular fixed-point κ , which is also a strong limit
 822 ordinal, so

$$f(\kappa) = g_2(\kappa) = \kappa \text{ and } \kappa \text{ is inaccessible.} \quad (3.90)$$

823 So every normal function d.f.a.o. has a regular fixed-point.

824 We have yet to show *Axiom Schema $M_1 \rightarrow$ Axiom Schema M_3* . Again by
 825 lemma (3.7), there is a normal function g defined for all ordinals that enumerates
 826 the fixed points of f . Let $h_\alpha(\beta) = g(\alpha + \beta)$ for any given ordinal α , then h_α
 827 is a normal function defined for all ordinals. Then, given an arbitrary α , from
 828 *Axiom Schema M_1* , there is a β such that $\gamma = h_\alpha(\beta)$ is inaccessible. Because
 829 $\gamma = g(\alpha + \beta)$, $f(\gamma) = \gamma$. Since $\alpha \leq f'(\alpha)$ for any ordinal α and any normal
 830 function f' , we know that $\alpha \leq \alpha + \gamma \leq \gamma$, so γ is inaccessible and arbitrarily
 831 large, depending on the choice of α . \square

832 But how do those schemata relate to reflection? Let's introduce a stronger
 833 version of *First-order reflection schema* from the previous chapter to see it more
 834 clearly. But in order to do this, we must establish the inaccessible cardinal first.

835 3.2 Inaccessible Cardinal

836 **Definition 3.9** *An uncountable cardinal κ is inaccessible iff it is regular and*
 837 *strongly limit. We write $In(\kappa)$ to say that κ is an inaccessible cardinal.*

838 An uncountable cardinal that is regular and limit is called a *weakly inaccessible*
 839 *cardinal*, we will only use the (strongly) inaccessible cardinal, but most of the
 840 results are similar for weakly inaccessible, including higher types of ordinals that
 841 will be presented later in this chapter.

842 **Theorem 3.10** *Let κ be an inaccessible cardinal.*

$$\langle V_\kappa, \in \rangle \models \text{ZFC} \quad (3.91)$$

843 We will prove this theorem in a way similar to [Kanamori, 2003]. *Proof.* Most
 844 of this is already done in lemma (2.15), we only need to verify that *Replacement*
 845 and *Infinity* axioms hold in V_κ .

846 *Infinity* holds because κ is uncountable, so $\omega \in V_\kappa$.

847 To verify *Replacement*, let x be an element of V_κ and f a function from x to
 848 V_κ . Let $y = \{z \in V_\kappa : (\exists q \in x)f(q) = z\}$, so $y \subset V_\kappa$, it remains to show that
 849 $y \in V_\kappa$. Because f is a function, we know that $|y| \leq |x| \leq \kappa$. But since κ
 850 is regular, $\{rank(z) : z \in y\} \subseteq \alpha$ for some $\alpha < \kappa$, and so $x \in V_{\alpha+1} \subseteq V_\kappa$.
 851 Therefore $y \in V_\kappa$. \square

852 **Definition 3.11** (*Inaccessible Reflection Schema*)

853 For every first-order formula φ , the following is an axiom:

$$\forall M_0 \exists \kappa (M_0 \subseteq V_\kappa \ \& \ In(\kappa) \ \& \ (\varphi(p_1, \dots, p_n) \leftrightarrow \varphi(p_1, \dots, p_n)^{V_\kappa})) \quad (3.92)$$

854 We will refer to this axiom schema as Inaccessible reflection schema.

855 We have added the requirement that α is inaccessible, which trivially means
 856 that there is an inaccessible cardinal. By taking appropriate M_0 , it can be shown
 857 that in a theory that includes the *Inaccessible reflection schema*, there is a closed
 858 unbounded class of inaccessible cardinals. Since we know that for an inaccessible
 859 κ , V_κ is a model of ZFC, *Inaccessible reflection schema* is equivalent to

$$\forall M_0 \exists \kappa (M_0 \subseteq V_\kappa \ \& \ \langle V_\kappa, \in \rangle \models \text{ZFC} \ \& \ (\varphi(p_1, \dots, p_n) \leftrightarrow \varphi(p_1, \dots, p_n)^{V_\kappa})) \quad (3.93)$$

860 because we have proven in the last section that for an inaccessible κ , $\langle V_\kappa, \in$
 861 $\rangle \models \text{ZFC}$.

862 **Theorem 3.12** Inaccessible reflection schema is equivalent to Axiom schema
 863 F .

864 This is *Theorem 4.1* in chapter four of [Drake, 1974], also equivalent to
 865 *Theorem 3* in [Lévy, 1960]. *Proof.* Let's start by showing that *Inaccessible*
 866 *reflection schema* implies *Axiom schema F*. It should be clear that we can
 867 reflect two formulas to a single set, just form a new formula as a conjunction of
 868 universal closures of the two.

869 Given a normal function f defined for all ordinals, we want to show that it
 870 has a regular fixed point. For any ordinal α , there is an ordinal κ such that

$$\alpha < \kappa \ \& \ In(\kappa) \ \& \ (\forall \gamma, \delta \in V_\kappa) (f(\gamma) = \delta \leftrightarrow (f(\gamma) = \delta)^{V_\kappa}) \quad (3.94)$$

871 and

$$\alpha < \kappa \ \& \ In(\kappa) \ \& \ \forall \gamma \exists \delta (f(\gamma) = \delta) \leftrightarrow (\forall \gamma \exists \delta f(\gamma) = \delta)^{V_\kappa} \quad (3.95)$$

872 Since V_κ is the set of all sets of rank less than κ and since every ordinal is the
873 rank of itself, there is an inaccessible ordinal κ such that

$$\forall \gamma < \kappa \exists \delta < \kappa (f^{V_\kappa}(\gamma) = \delta) \quad (3.96)$$

874 We also know that $f(\gamma) = \delta \leftrightarrow (f(\gamma) = \delta)^{V_\kappa}$. Now since κ is a limit ordinal
875 and f is continuous we get

$$f(\kappa) = \bigcup_{\gamma < \kappa} f^{V_\kappa}(\gamma) = \bigcup_{\gamma < \kappa} f(\gamma). \quad (3.97)$$

876 From (3.96) and the fact that f is increasing, we know that $\kappa \leq \bigcup_{\gamma < \kappa} f(\gamma) \leq \kappa$.
877 Therefore κ is an inaccessible fixed point of f .

878 For the opposite direction, it suffices to show that since there is an inacces-
879 sible cardinal from *Axiom schema F*, given a first-order formula φ , there is an
880 arbitrarily large inaccessible cardinal κ for which

$$\varphi \leftrightarrow \langle V_\kappa, \in \rangle \models \varphi. \quad (3.98)$$

881 Note that the arbitrary size of κ means given an arbitrary ordinal α , there is a
882 κ satisfying (3.98). In the previous chapter, in theorem (2.12), we have shown
883 that we can easily obtain a limit ordinal satisfying (3.98). Note that since for
884 any set M_0 , there is such α that $M_0 \subseteq V_\alpha$, there is a closed unbounded class of
885 sets satisfying (3.98), which are levels in the cumulative hierarchy, so there is a
886 club sets of κ s satisfying (3.98).

887 Let f be a normal function defined for all ordinals that enumerates this club
888 class, there is such by lemma (3.2). Let g be the function that enumerates
889 strong limit ordinals in $\text{rng}(f)$. Then g has a regular fixed point κ , which is also
890 a regular fixed point of f , so (3.98) holds for κ .

891

□

892 Definition 3.13 (ZMC)

893 We will call ZMC an axiomatic set theory that contains all axioms and schemas
894 of ZFC together with Axiom Schema M_1 .

895 We have decided to call it ZMC, because Lévy uses ZM, derived from ZF, which
896 is more intuitive, but we also need the axiom of choice, thus, ZMC.

897 3.3 Mahlo Cardinals

898 We have shown that ZMC contains arbitrarily large inaccessible cardinals. To
899 return to reflection-style argument, is there a set that satisfies this property? To
900 be able to properly answer this question, we have to formulate the notion of

"containing arbitrarily large cardinals" more carefully. While we have previously used club sets, this is not an option because inaccessible don't form a club class in ZMC^{22} , we could try to formulate stronger versions of *Axiom Schema M_1* .

Let's shortly review what *Axiom Schema M_1* says. We have shown earlier in this chapter that there is a simple relation between normal function defined for all ordinals and closed unbounded classes. So by saying that for a class of ordinals C , a normal function f has at least one element of C in its range, we say that C is stationary. Or, as Drake puts it for C , the class of inaccessible cardinals, and a κ , in which C is stationary:

"The class of inaccessible cardinals is so rich that there are members κ of the class such that no normal function on κ can avoid this class; however we climb through κ , provided we are continuous at limits (so that we are enumerating a closed subset of κ), we shall eventually have to hit an inaccessible."

Definition 3.14 (*Mahlo Cardinal*)

We say that κ is a Mahlo Cardinal iff it is an inaccessible cardinal and the set $\{\lambda < \kappa : \lambda \text{ is inaccessible}\}$ is stationary in κ .

Alternatively, κ is Mahlo iff $\langle V_\kappa, \in \rangle \models ZMC$ as shown above, this is also sometimes written as *Ord is Mahlo*. There are also *weakly Mahlo cardinals*, that are defined via weakly inaccessible cardinal below them, Mahlo cardinals are then also called *strongly Mahlo* to highlight the difference, but we will only use the term *Mahlo cardinal*.

Mahlo cardinals are related to reflection principles in an interesting way. Note that given a formula φ , reflection gave us a club set of ordinals α such that V_α reflects φ , all below the first inaccessible cardinal. We have then used a different reflection schema to obtain arbitrarily high inaccessible cardinals. Now we have a cardinal in which this reflection schema holds, so we're in fact reflecting reflection. Beware that this is done rather informally, because *Axiom Schema M_1* is a countable set of axioms, which is too large to be reflected via the schemas introduced so far. One way to deal with this would be to extend reflection for second- and possibly higher-order formulas, but we would have to be very careful with the notion of satisfaction. For now, explore where can stationary sets take us because as we have shown, their connection to reflection is quite clear.

What would happen if we strengthened *Axiom Schema M_1* to say that every normal function has a Mahlo cardinal in its range?

Definition 3.15 (*hyper-Mahlo cardinal*)

We say that κ is a hyper-Mahlo cardinal iff it is inaccessible and the set $\{\lambda < \kappa : \lambda \text{ is Mahlo}\}$ is stationary in κ .

²²Note that cofinality of the limit of the first ω inaccessibles is ω , which makes it singular.

939 **Definition 3.16** (*hyper-hyper-Mahlo cardinal*)

940 We say that κ is a hyper-hyper-Mahlo cardinal iff it is inaccessible and the set
 941 $\{\lambda < \kappa : \lambda \text{ is hyper-Mahlo}\}$ is stationary in κ .

942 It is clear that one can continue in this direction, but the nomenclature gets
 943 increasingly overwhelming even if we introduce *hyper $^\alpha$ -Mahlo cardinals*. To see
 944 this, let's try to establish an operation that would exhaust all such cardinals in a
 945 more unified manner.

946 **Definition 3.17** (*Mahlo Operation*)

947 Let A be a class of ordinals. Let

$$H(A) = \{\alpha \in A : A \cap \alpha \text{ is stationary in } \alpha\}. \quad (3.99)$$

948 We call H the Mahlo's operation.

949 If we pick for A the class of all inaccessible cardinals, $H(A)$ is the class of
 950 Mahlo cardinals. It is easy to see that if A is the class of all α -Mahlo cardinals,
 951 $H(A)$ is the class of $\alpha + 1$ -Mahlo cardinals, $H(H(A))$ is the class of $\alpha + 2$ -
 952 Mahlo cardinals and so on.

953 **Definition 3.18** (*Iterated Mahlo Operation*)

954 Let A be a class of ordinals. We shall extend the Mahlo operation in the following
 955 way:

- 956 (i) $H^0(A) = A$,
- 957 (ii) $H^{\alpha+1}(A) = H(H^\alpha(A))$,
- 958 (iii) $H^\lambda(A) = \bigcap_{\alpha < \lambda} H^\alpha(A)$ for limit λ .

959 Clearly if A is the class of inaccessibles, $H^\alpha(A)$ is the class of α -Mahlo
 960 cardinals. To get to hyper-Mahlo cardinals, we can diagonalize the operation.

961 **Definition 3.19** (*Diagonal Mahlo Operation*)

962 Let A be again a class of ordinals. Then the diagonal Mahlo operation is defined
 963 as follows:

$$H^\Delta(A) = \{\alpha : \forall \beta < \alpha (\alpha \in H^\beta(A))\}. \quad (3.100)$$

964 We can further diagonalize the diagonal version and continue this process ad
 965 libitum. All of the above steps can be turned into axioms similar to ... TODO ?

3.4 Indescribable Cardinals

Definition 3.20 (Describability)

We say an ordinal α is described by a sentence φ in the language \mathcal{L} with relation symbols P_1, \dots, P_n given iff

$$\langle V_\alpha, \in, P_1, \dots, P_n \rangle \models \varphi \quad (3.101)$$

but for every $\beta < \alpha$

$$\langle V_\beta, \in, P_1 \cap V_\beta, \dots, P_n \cap V_\beta \rangle \not\models \varphi \quad (3.102)$$

For the definition of a Π_n^m -formula and a Σ_n^m -formula, see TODO and TODO respectively.

TODO nezavest spis pro provoradove fle?

TODO Kanamori to dela pro vyssi rady, coz nas nesere?

Definition 3.21 (Π_n^m -Indescribable Cardinal)

We say that κ is Π_n^m -indescribable iff it is not described by any Π_n^m -formula.

Definition 3.22 (Σ_n^m -Indescribable Cardinal)

We say that κ is Σ_n^m -indescribable iff it is not described by any Σ_n^m -formula.

To see that this notion is based in reflection, note that for Π_n^m -formulas²³, a cardinal κ is Π_n^m -indescribable iff every Π_n^m -formula reflects in κ in the sense of definition (??). Informally, can also view indescribability as a property held by the universe V , in the sense that every formula aiming to describe it in fact describes an initial segment, which is similar to a reflection principle, albeit stated informally.²⁴

Since we are interested accessing cardinals from below via fixed points of normal functions, we will limit ourselves to Π_n^1 -formulas, with the exception of measurable cardinal, that is included for context.

Lemma 3.23 *Let κ be a cardinal, the following holds for any $n \in \omega$. κ is Π_n^1 -indescribable iff κ is $\Sigma_n^1 + 1$ -indescribable*

²³This holds for Σ_n^m -formulas alike.

²⁴Formally, we have to be once again careful with “properties of V ” for the reasons mentioned in the introduction of this thesis. That’s why this chapter only reflects sentences to models with additional relations.

992 *Proof.* The forward direction is obvious, we can always add a spare quantifier
 993 over a type 2 variable to turn a Π_n^1 formula φ into a $\exists P\varphi$ which is obviously a
 994 $\Sigma_n^1 + 1$ formula.²⁵

995 To prove the opposite direction, suppose that $\langle V_\kappa, \in \rangle \models \exists X\varphi(X)$ where
 996 X is a type 2 variable and φ is a Π_n^1 formula with one free variable of type
 997 2. This means that there is a set $S \subseteq V_\kappa$ that is a witness of $\exists X\varphi(X)$, in
 998 other words, $\varphi(S)$ holds. We can replace every occurrence of X in φ by a new
 999 predicate symbol S , this allows us to say that κ is Π_n^1 -indescribable (with respect
 1000 to $\langle V_\kappa, \in, R, S \rangle$).²⁶ \square

1001 The above lemma makes it clear that we can suppose that all formulas with
 1002 no higher than type 2 variables are Π_n^1 -formulas, $n \in \omega$, without the loss of
 1003 generality.

1004 **Lemma 3.24** *If κ is an inaccessible cardinal and given $R \subseteq V_\kappa$, then the*
 1005 *following is a club set in κ :*

$$\{\alpha : \alpha < \kappa \text{ \& } \langle V_\alpha, \in, R \cap V_\alpha \rangle \prec \langle V_\kappa, \in, R \rangle\} \quad (3.103)$$

1006 *Proof.* To see that (3.103) is closed, let us recall that a $A \subseteq \kappa$ is closed iff for
 1007 every ordinal $\alpha < \lambda$, $\alpha \neq \emptyset$: if $A \cap \alpha$ is unbounded in α then $\alpha \in A$. Since κ is
 1008 an inaccessible cardinal, thus strong limit, it is closed under limits of sequences
 1009 of ordinals lesser than κ .

1010 We want to verify that it is unbounded, we will use a recursively defined
 1011 sequence $\alpha_0, \alpha_1, \dots$ to build an elementary substructure of $\langle V_\kappa, \in, R \rangle$ that is
 1012 built above an arbitrary $\alpha_0 < \kappa$. Let us fix an arbitrary $\alpha_0 < \kappa$. Given α_n ,
 1013 $\alpha_n + 1$ is defined as the least β , $\alpha_n \leq \beta$ that satisfies the following for any
 1014 formula φ , $p_1, \dots, p_m \in V_{\alpha_n}$, $m \in \omega$:

$$\text{If } \langle V_\kappa, \in, R \rangle \models \exists x\varphi(p_1, \dots, p_n), \text{ then } \langle V_\kappa, \in, R \rangle \models \varphi(x, p_1, \dots, p_n) \quad (3.104)$$

1015 Let $\alpha = \bigcup_{n < \omega} \alpha_n$.

1016 Then $\langle V_\alpha, \in, R \cap V_\alpha \rangle \prec \langle V_\kappa, \in, R \rangle$, in other words, for any φ with given
 1017 arbitrary parameters $p_1, \dots, p_n \in V_\alpha$, it holds that

$$\langle V_\alpha, \in, R \cap V_\alpha \rangle \models \varphi(p_1, \dots, p_n) \leftrightarrow \langle V_\kappa, \in, R \rangle \models \varphi(p_1, \dots, p_n) \quad (3.105)$$

1018 Which should be clear from the construction of α \square

²⁵Note that unlike in previous sections, it is worth noting that φ is now a sentence so we don't have to worry whether P is free in φ .

²⁶A different yet interesting approach is taken by Tate in [Tait, 2005]. He states that for $n \geq 0$, a formula of order $\leq n$ is called a Π_0^n and a Σ_0^n formula. Then a Π_{m+1}^n is a formula of form $\forall Y\psi(Y)$ where ψ is a Σ_m^n formula and Y is a variable of type n . Finally, a Σ_{m+1}^n is the negation of a Π_m^n formula. So the above holds ad definitio.

1019 **Theorem 3.25** *Let κ be an ordinal. The following are equivalent.*

- 1020 (i) κ is inaccessible
 1021 (ii) κ is Π_0^1 -indescribable.

1022 *Proof.* Since Π_0^1 -sentences are first-order sentences, we want to prove that κ is
 1023 an inaccessible cardinal iff whenever a first-order tries to describe κ in the sense
 1024 of definition (3.20), the formula fails to do so and describes a initial segment
 1025 thereof instead. We have already shown in (??) that there is no way to reach an
 1026 inaccessible cardinal via first-order formulas in ZFC. We will now prove it again
 1027 in for formal clarity.

1028 For (i)→(ii), suppose that κ is inaccessible.

1029 Then there is, by lemma (3.24) a club set of ordinals α such that V_α is
 1030 an elementary substructures of V_κ . For κ to be Π_0^1 -indescribable, we need to
 1031 make sure that given an arbitrary first-order sentence φ satisfied in the structure
 1032 $\langle V_\kappa, \in, R \rangle$, there is an ordinal $\alpha < \kappa$, such that $\langle V_\alpha, \in, R \cap V_\alpha \rangle \models \varphi$. But this
 1033 follows from the definition of elementary substructure.

1034 For (ii)→(i), suppose κ is not inaccessible, so it is either singular, or there is
 1035 a cardinal $\nu < \kappa$ such that $\kappa \leq \mathcal{P}(\nu)$ or $\kappa = \omega$.

1036 Suppose κ is singular. Then there is a cardinal $\nu < \kappa$ and a function $f : \nu \rightarrow$
 1037 κ such that $\text{rng}(f)$ is cofinal in κ . Since $f \subseteq V_\kappa$, we can add f as a relation
 1038 to the language. We can do the same with $\{\nu\}$. That means $\langle V_\kappa, \in, P_1, P_1$
 1039 with $P_1 = f, P_2 = \{\nu\}$ is a structure, let $\varphi = P_1 \neq \emptyset \ \& \ \text{rng}(P_1) = P_2$ ²⁷.
 1040 Since for every $\alpha < \nu$, $P_1 \cap V_\alpha = \emptyset$, φ is false and therefore describes κ . That
 1041 contradicts the fact that κ was supposed to be Π_0^1 -indescribable, but φ is a
 1042 first-order formula.

1043 Suppose there a cardinal ν satisfying $\kappa \leq \mathcal{P}(\nu)$. Let there be a function
 1044 $f : \mathcal{P}(\nu) \rightarrow \kappa$ that is onto. Then, like in the previous paragraph, we can obtain
 1045 a structure $\langle V_\kappa, \in, P_1, P_2 \rangle$, where $P_1 = f$ like before, but this time $P_2 = \mathcal{P}(\nu)$.
 1046 Again, $\varphi = P_1 \neq \emptyset \ \& \ \text{rng}(P_1) = P_2$ describes κ .

1047 Finally, suppose $\kappa = \omega$, then the sentence $\varphi = \forall x \exists y (x \in y)$ describes κ ,
 1048 there is obviously no $\alpha < \omega$ such that $\langle V_\alpha, \in \rangle \models \varphi$.

1049 □

1050 Generally, it should be clear that it a cardinal κ is Π_n^m -indescribable, it is
 1051 also $\Pi_{n'}^{m'}$ -indescribable for every $m' < m, n' < n$. By the same line of thought,
 1052 if a cardinal κ satisfies property implied by Π_n^m -indescribability, it satisfies all
 1053 properties implied by $\Pi_{n'}^{m'}$ -indescribability for $m' < m, n' < n$, for example κ is
 1054 Π_n^m -indescribable for $m \geq 1, n \geq 0$, it is also an inaccessible cardinal.

1055 **Theorem 3.26** *If a cardinal κ is Π_1^1 -indescribable, then it is a Mahlo cardinal.*

²⁷ $\text{rng}(x) = y$ is a first-order formula, see (1.14).

1056 *Proof.* Assuming that κ is Π_1^1 -indescribable, we want to prove that every club
 1057 set in κ contains an inaccessible cardinal.

1058 Consider the following Π_1^1 -sentence:

$$\forall P(\text{"}P \text{ is a function"} \ \& \ \exists x(x = \text{dom}(P) \vee \mathcal{P}(x) = \text{dom}(P)) \rightarrow \rightarrow \exists y(y = \text{rng}(P))) \quad (3.106)$$

1059 where P is a type 2 variable and x, y are type 1 variables, $\text{rng}(P)$ is defined in
 1060 (1.14), $\text{dom}(P)$ in (1.13) and " P is a function" is a first-order formula defined
 1061 in (1.11). We will call this sentence *Inac*, as in "inaccessible", because, given a
 1062 cardinal μ , the following holds if and only if μ is inaccessible:

$$\langle V_\mu, \in \rangle \models \text{Inac} \quad (3.107)$$

1063 So let's fix an arbitrary $C \subset \kappa$, club set in κ . We want to show that it
 1064 contains an inaccessible cardinal. Since C is a subset of V_κ , let's add it to the
 1065 structure $\langle V_\kappa, \in \rangle$, turning it into $\langle V_\kappa, \in, C \rangle$. Then the following holds:

$$\langle V_\kappa, \in, C \rangle \models \text{Inac} \ \& \ \text{"}C \text{ in unbounded"} \quad (3.108)$$

1066 Note that this is correct, because, as we have noted just before introducing the
 1067 statement now being proven, if κ is Π_1^1 -indescribable, it is also Π_0^1 -indescribable.
 1068 So κ is itself inaccessible and therefore $\langle V_\kappa, \in, C \rangle \models \text{Inac}$. C is obviously
 1069 picked so that it is unbounded in κ ²⁸.

1070 Now because we have assumed that κ is Π_1^1 -indescribable and *Inac* is a Π_1^1 -
 1071 formula, so *Inac* & " C in unbounded" is equivalent to a Π_1^1 -formula, there must
 1072 be an ordinal α that satisfies

$$\langle V_\alpha, \in, C \cap V_\alpha \rangle \models \text{Inac} \ \& \ \text{"}C \text{ in unbounded"} \quad (3.109)$$

1073 which implies that α is inaccessible.

1074 To be finished, we need to verify that $\alpha \in C$. Since $\kappa = V_\kappa$ for inaccessible
 1075 κ , $C \cap V_\alpha = C \cap \alpha$, from unboundedness of $C \cap \alpha$ in α , $\bigcup(C \cap \alpha) = \alpha$, which,
 1076 together with the fact that C is a club set in κ and therefore closed in κ , yields
 1077 that $\alpha \in C$. \square

²⁸ " C in unbounded" is a first-order formula defined in (1.52).

¹⁰⁷⁸ **4 Conclusion**

References

- 1079
- 1080 [Church, 1996] Church, A. (1996). *Introduction to Mathematical Logic*. Annals
1081 of Mathematics Studies. Princeton University Press.
- 1082 [Drake, 1974] Drake, F. (1974). *Set theory. An introduction to large cardinals*.
1083 Studies in Logic and the Foundations of Mathematics, Volume 76. NH.
- 1084 [Hamilton, 1988] Hamilton, A. (1988). *Logic for Mathematicians*. Cambridge
1085 University Press.
- 1086 [Jech, 2006] Jech, T. (2006). *Set theory*. Springer monographs in mathematics.
1087 Springer, the 3rd millennium ed., rev. and expanded edition.
- 1088 [Kanamori, 2003] Kanamori, A. (2003). *The higher infinite: Large cardinals*
1089 *in set theory from their beginnings*. Springer Monographs in Mathematics.
1090 Springer-Verlag Berlin Heidelberg, 2 edition.
- 1091 [Lévy, 1960] Lévy, A. (1960). Axiom schemata of strong infinity in axiomatic set
1092 theory. *Pacific Journal of Mathematics*, 10.
- 1093 [Tait, 2005] Tait, W. W. (2005). Constructing cardinals from below. The Prove-
1094 nance of Pure Reason: Essays in the Philosophy of Mathematics and Its His-
1095 tory, 133-154.
- 1096 [Wang, 1997] Wang, H. (1997). *"A Logical Journey: From Gödel to Philoso-*
1097 *phy"*. A Bradford Book.