

Univerzita Karlova v Praze, Filozofická fakulta
Katedra logiky

MIKULÁŠ MRVA

REFLECTION PRINCIPLES AND LARGE
CARDINALS

Bakalářská práce

Vedoucí práce: Mgr. Radek Honzík, Ph.D.

2016

Prohlašuji, že jsem bakalářskou práci vypracoval samostatně a že jsem uvedl všechny použité prameny a literaturu.

V Praze 22. května 2016

Mikuláš Mrva

Abstract

Práce zkoumá vztah tzv. principů reflexe a velkých kardinálů. Lévy ukázal, že v ZFC platí tzv. věta o reflexi a dokonce, že věta o reflexi je ekvivalentní schématu nahrazení a axiomu nekonečna nad teorií ZFC bez axiomu nekonečna a schématu nahrazení. Tedy lze na větu o reflexi pohlížet jako na svého druhu axiom nekonečna. Práce zkoumá do jaké míry a jakým způsobem lze větu o reflexi zobecnit a jaký to má vliv na existenci tzv. velkých kardinálů. Práce definuje nedosažitelné, Mahlovy a nepopsatelné kardinály a ukáže, jak je lze zavést pomocí reflexe. Přirozenou limitou kardinálů získaných reflexí jsou kardinály nekonzistentní s L. Práce nabídne intuitivní zdůvodnění, proč tomu tak je.

Abstract

This thesis aims to examine relations between the so called Reflection Principles and Large cardinals. Lévy has shown that the Reflection Theorem is a sound theorem of ZF and it is equivalent to the Replacement Scheme and the Axiom of Infinity. From this point of view, Reflection theorem can be seen a specific version of an Axiom of Infinity. This paper aims to examine the Reflection Principle and its generalisations with respect to the existence of Large Cardinals. This thesis will establish the Inaccessible, Mahlo and Indescribable cardinals and show how can those be defined via reflection. A natural limit of Large Cardinals obtained via reflection are cardinals inconsistent with L. This thesis will offer an intuitive explanation of why this holds.

Contents

1	Introduction	4
1.1	Motivation and Origin	4
1.2	Notation and Terminology	4
1.2.1	The Language of Set Theory	4
1.2.2	The Axioms	4
1.2.3	The Transitive Universe	8
1.2.4	Cardinal Numbers	10
1.2.5	Relativisation and Absoluteness	12
1.2.6	More Functions	13
1.2.7	Structure, Substructure and Embedding	14
2	Levy's First-Order Reflection	15
2.1	Lévy's Original Paper	15
2.2	Contemporary Restatement	18
3	Conclusion	26

1 Introduction

1.1 Motivation and Origin

The Universe of sets cannot be uniquely characterised (i. e. distinguished from all its initial elements) by any internal structural property of the membership relation in it, which is expressible in any logic of finite or transfinite type, including infinitary logics of any cardinal order.

— Kurt Gödel [Wang, 1997]

1.2 Notation and Terminology

1.2.1 The Language of Set Theory

This text assumes the knowledge of basic terminology and some results from first-order predicate logic.¹

We will now shortly review the basic notions that allow us to define *Zermelo–Fraenkel* set theory.

When we talk about *class*, we have the notion of definable class in mind. If $\varphi(x, p_1, \dots, p_n)$ is a formula in the language of set theory, we call

$$A = \{x : \varphi(x)\} \tag{1.1}$$

a class of all sets satisfying $\varphi(x)$ in a sense that

$$x \in A \leftrightarrow \varphi(x) \tag{1.2}$$

Given classes A, B , one can easily define the elementary set operations such as $A \cap B, A \cup B, A \setminus C, \bigcup A$, see the first part of [Jech, 2006] for details. The following axioms are the tools by which decide whether particular classes are in fact sets. A class that fails to be considered a set is called a *proper class*.

Speaking of formulas, we will often use syntax like " M is a limit ordinal". It should be clear that this statement can be rewritten as a formula that was introduced earlier in the text.

1.2.2 The Axioms

Definition 1.1 (*The Existence of a Set*)

$$\exists x(x = x) \tag{1.3}$$

¹todo odkaz na pripadny zdroj? svejdar? neco en?

The above axiom is usually not used because it can be deduced from the axiom of *Infinity* (see below), but since we will be using set theories that omit *Infinity*, this will be useful.

Definition 1.2 (*Extensionality*)

$$\forall x, y (x = y \leftrightarrow \forall z (z \in x \leftrightarrow z \in y)) \quad (1.4)$$

Definition 1.3 (*Specification*)

The following is a schema for every first-order formula $\varphi(x, p_1, \dots, p_n)$ with no free variables other than x, p_1, \dots, p_n .

$$\forall x, p_1, \dots, p_n \exists y \forall z (z \in y \leftrightarrow z \in x \ \& \ \varphi(z, p_1, \dots, p_n)) \quad (1.5)$$

We will now provide two definitions that are not axioms, but will be helpful in establishing some of the other axioms in a more intuitive way.

Definition 1.4 ($x \subseteq y, x \subset y$)

$$x \subseteq y \leftrightarrow (\forall z \in x) z \in y \quad (1.6)$$

$$x \subset y \leftrightarrow x \subseteq y \ \& \ x \neq y \quad (1.7)$$

Definition 1.5 (*Empty Set*) For an arbitrary set x , the empty set, written as \emptyset , is defined by the following formula

$$(\forall y \in x)(y \in \emptyset \leftrightarrow \neg(y = y)) \quad (1.8)$$

We know that \emptyset is a set from specification. While it could also be defined by " $\forall y(y \in \emptyset \leftrightarrow \neg(y = y))$ ", we have decided to use the bounded version because it's Δ_0 , which we will find useful later on. The two definitions yield the same set for every x given thanks to extensionality.

Definition 1.6 (*Pairing*)

$$\forall x, y \exists z \forall q (q \in z \leftrightarrow q = x \vee q = y) \quad (1.9)$$

Definition 1.7 (*Union*)

$$\forall x \exists y \forall z (z \in y \leftrightarrow \exists q (z \in q \ \& \ q \in x)) \quad (1.10)$$

Definition 1.8 (*Set Intersection*)

$$x \cap y = \{z : z \in x \ \& \ z \in y\} \quad (1.11)$$

Definition 1.9 (*Set Union*)

$$x \cup y = \{z : z \in x \vee z \in y\} \quad (1.12)$$

Now we can introduce more axioms.

Definition 1.10 (*Foundation*)

$$\forall x(x \neq \emptyset \rightarrow (\exists y \in x)(x \cap y = \emptyset)) \quad (1.13)$$

Definition 1.11 (*Powerset*)

$$\forall x \exists y \forall z (z \subseteq x \leftrightarrow z \in y) \quad (1.14)$$

Definition 1.12 (*Infinity*)

$$\exists x(\emptyset \in x \ \& \ (\forall y \in x)(y \cup \{y\} \in x)) \quad (1.15)$$

Let us introduce a few more definitions that will make the two remaining axioms more comprehensible.

Definition 1.13 (*Powerset function*)

Given a set x , the powerset of x , denoted $\mathcal{P}(x)$ and satisfying 1.11, is defined as follows:

$$\mathcal{P}(x) \stackrel{\text{def}}{=} \{y : y \subseteq x\} \quad (1.16)$$

Definition 1.14 (*Function*)

Given arbitrary first-order formula $\varphi(x, y, p_1, \dots, p_n)$, we say that φ is a function iff

$$\forall x, y, z, p_1, \dots, p_n (\varphi(x, y, p_1, \dots, p_n) \ \& \ \varphi(x, z, p_1, \dots, p_n) \rightarrow y = z) \quad (1.17)$$

When a $\varphi(x, y)$ is a function, we also write the following:

$$f(x) = y \leftrightarrow \varphi(x, y) \quad (1.18)$$

² Alternatively, $f = \{\langle x, y \rangle : \varphi(x, y)\}$ is a class.

Definition 1.15 (*Dom(f)*)

Let f be a function. We read the following as " $\text{Dom}(f)$ is the domain of f ".

$$\text{Dom}(f) \stackrel{\text{def}}{=} \{x : \exists y (f(x) = y)\} \quad (1.19)$$

²This can also be done for φ s with more than two free variables by either setting $f(x, p_1, \dots, p_n) = y \leftrightarrow \varphi(x, y, p_1, \dots, p_n)$ or saying that φ codes more functions, determined by the various parameters, so given t_1, \dots, t_n , $f(x) = y \leftrightarrow \varphi(x, y, t_1, \dots, t_n)$.

We say " f is a function on A ", A being a class, if $A = \text{dom}(f)$.

Definition 1.16 (*Rng(f)*)

Let f be a function. We read the following as " $\text{Rng}(f)$ is the range of f ".

$$\text{Rng}(f) \stackrel{\text{def}}{=} \{x : \exists y(f(x) = y)\} \quad (1.20)$$

We say that f is a *function into* A , A being a class, if $\text{rng}(f) \subseteq A$. We say that f is a *function onto* A if $\text{rng}(f) = A$, in other words,

$$(\forall y \in A)(\exists x \in \text{dom}(f))(f(x) = y) \quad (1.21)$$

We say a function f is a *one to one function*, iff

$$(\forall x_1, x_2 \in \text{dom}(f))(f(x_1) = f(x_2) \rightarrow x_1 = x_2) \quad (1.22)$$

f is a bijection iff it is a one to one function that is onto.

Note that $\text{Dom}(f)$ and $\text{Rng}(f)$ are not definitions in a strict sense, they are in fact definition schemas that yield definitions for every function f given. Also note that they can be easily modified for φ instead of f , with the only difference being the fact that it is then defined only for those φ s that are functions, which must be taken into account. This is worth noting as we will sometimes interchange the notions of *function* and *formula*.

Definition 1.17 (*Function Defined For All Ordinals*)

We say a function f is defined for all ordinals, this is sometimes written $f : \text{Ord} \rightarrow A$ for any class A , if $\text{Dom}(f) = \text{Ord}$. Alternatively,

$$(\forall \alpha \in \text{Ord})(\exists y \in A)(f(\alpha) = y) \quad (1.23)$$

And now for the axioms.

Definition 1.18 (*Replacement*)

The following is a schema for every first-order formula $\varphi(x, p_1, \dots, p_n)$ with no free variables other than x, p_1, \dots, p_n .

$$"\varphi \text{ is a function}" \rightarrow \forall x \exists y \forall z (z \in y \leftrightarrow (\exists q \in x)(\varphi(x, y, p_1, \dots, p_n))) \quad (1.24)$$

Definition 1.19 (*Choice*)

$$\begin{aligned} &\forall x \exists f ((f \text{ is a choice function with } \text{dom}(f) = x \setminus \{\emptyset\}) \\ &\quad \& \forall y ((y \in x \& y \neq \emptyset) \rightarrow f(y) \in y)) \end{aligned} \quad (1.25)$$

We will refer the axioms by their name, written in italic type, e.g. *Foundation* refers to the Axiom of Foundation. Now we need to define some basic set theories to be used in the article.

Definition 1.20 (S)

We call **S** an axiomatic theory in the language $\mathcal{L} = \{=, \in\}$ with exactly the following axioms:

- (i) Existence of a set (see 1.1)
- (ii) Extensionality (see 1.2)
- (iii) Specification (see 1.3)
- (iv) Foundation (see 1.10)
- (v) Pairing (see 1.6)
- (vi) Union (see 1.7)
- (vii) Powerset (see 1.11)

Definition 1.21 (ZF)

We call **ZF** an axiomatic theory in the language $\mathcal{L} = \{=, \in\}$ that contains all the axioms of **S** in addition to the following

- (i) Replacement schema (see 1.18)
- (ii) Infinity (see 1.12)

Existence of a set is usually left out because it is a consequence of infinity.

Definition 1.22 (ZFC)

ZFC is an axiomatic theory in the language $\mathcal{L} = \{=, \in\}$ that contains all the axioms of **ZF** plus Choice (1.19).

1.2.3 The Transitive Universe

Definition 1.23 (Transitive Class)

We say a class A is transitive iff

$$(\forall x \in A)(x \subseteq A) \tag{1.26}$$

Definition 1.24 (Well Ordered Class) A class A is said to be well ordered by \in iff the following hold:

- (i) $(\forall x \in A)(x \not\in x)$ (*Antireflexivity*)
- (ii) $(\forall x, y, z \in A)(x \in y \ \& \ y \in z \rightarrow x \in z)$ (*Transitivity*)
- (iii) $(\forall x, y \in A)(x = y \vee x \in y \vee y \in x)$ (*Linearity*)
- (iv) $(\forall x \subseteq A)(x \neq \emptyset \rightarrow (\exists y \in x)(\forall z \in x)(z = y \vee z \in y))$ (*Existence of the least element*)

Definition 1.25 (*Ordinal Number*)

A set x is said to be an ordinal number if it is transitive and well-ordered by \in .

For the sake of brevity, we usually just say " x is an ordinal". Note that " x is an ordinal" is a well-defined formula in the language of set theory, since 1.23 is a first-order formula and 1.24 is in fact a conjunction of four first-order formulas. Ordinals will be usually denoted by lower case greek letters, starting from the beginning: $\alpha, \beta, \gamma, \dots$. Given two different ordinals α, β , we will write $\alpha < \beta$ for $\alpha \in \beta$, see [Jech, 2006] Lemma 2.11 for technical details.

Definition 1.26 (*Non-Zero Ordinal*) We say an ordinal α is non-zero iff $\alpha \neq \emptyset$.

Definition 1.27 (*Successor Ordinal*)

Consider the following operation, let β be an ordinal.

$$S(\beta) = \beta \cup \{\beta\} \quad (1.27)$$

An ordinal α is called a successor ordinal iff there is an ordinal β , such that $\alpha = S(\beta)$. We will sometimes also write $\alpha = \beta + 1$.

Definition 1.28 (*Limit Ordinal*)

A non-zero ordinal α is called a limit ordinal iff it is not a successor ordinal.

Definition 1.29 (*Ord*)

The class of all ordinal numbers, which we will denote Ord^3 is the proper class defined as follows.

$$\text{Ord} = \{x : x \text{ is an ordinal}\} \quad (1.28)$$

Definition 1.30 (*Von Neumann's Hierarchy*)

The Von Neumann's Hierarchy is a collection of sets indexed by elements of Ord , defined recursively in the following way:

(i)

$$V_0 = \emptyset \quad (1.29)$$

(ii)

$$V_{\alpha+1} = \mathcal{P}(V_\alpha) \text{ for any ordinal } \alpha \quad (1.30)$$

³It is sometimes denoted On , but we will stick to the notation used in [Jech, 2006]

(iii)

$$V_\lambda = \bigcup_{\beta < \lambda} V_\beta \text{ for a limit ordinal } \lambda \quad (1.31)$$

We will also refer to the Von Neumann's Hierarchy as Von Neumann's Universe or the Cumulative Hierarchy.

Definition 1.31 (*Rank*)

Given a set x , we say that the rank of x (written as $\text{rank}(x)$) is the least ordinal α such that

$$x \in V_{\alpha+1} \quad (1.32)$$

Due to *Regularity*, every set has a rank.⁴

Definition 1.32 (ω)

$$\omega \stackrel{\text{def}}{=} \bigcap \{x : "x \text{ is a limit ordinal}"\} \quad (1.33)$$

ω is non-empty if *Infinity* or any equivalent holds.

Definition 1.33 (*Lévy's Hierarchy*)

!!! pozor na konflikt s analytickou (vyres podle kanamoriho) TODO

1.2.4 Cardinal Numbers**Definition 1.34** (*Cardinality*)

Given a set x , let the cardinality of x , written $|x|$, be defined as the smallest ordinal number such that there is a one to one mapping from x to α .

Definition 1.35 (*Aleph function*)

Let ω be the set defined by 1.32. We will recursively define the function \aleph for all ordinals.

(i) $\aleph_0 = \omega$

(ii) $\aleph_{\alpha+1}$ is the least cardinal larger than \aleph_α ⁵

(iii) $\aleph_\lambda = \bigcup_{\beta < \lambda} \aleph_\beta$ for a limit ordinal λ

If $\kappa = \aleph_\alpha$ and α is a successor ordinal, we call κ a *successor cardinal*. If α is a limit ordinal, we call κ a *limit cardinal*.

⁴See chapter 6 of [Jech, 2006] for details.

⁵"The least cardinal larger than \aleph_α " is sometimes notated as \aleph_α^+

Definition 1.36 (*Cardinal number*)

- (i) A set x is called a finite cardinal iff $x \in \omega$.
- (ii) A set is called an infinite cardinal iff there is an ordinal α such that $\aleph_\alpha = x$
- (iii) A set is called a cardinal iff it is either a finite cardinal or an infinite cardinal.

We say κ is an uncountable cardinal if it is an infinite ordinal and $\aleph_0 > \kappa$. Infinite cardinals will be notated by lowercase greek letters from the middle of the alphabet, e.g. κ, μ, ν, \dots ⁶

For formal details as well as why every set can be well-ordered assuming *Choice*, and therefore has a cardinality, see [Jech, 2006].

Definition 1.37 (*Cofinality of a Limit Ordinal*)

Let λ be a limit ordinal. We say that the cofinality of λ is α iff α is the smallest limit ordinal, such that there is an α -sequence $\langle \beta_\xi : \xi < \alpha \rangle$, such that

$$\sup(\beta_\xi : \xi < \alpha) = \lambda \quad (1.34)$$

We write $cf(\lambda) = \alpha$.

Definition 1.38 (*Regular Cardinal*)

We say a cardinal κ is regular iff $cf(\kappa) = \kappa$

Definition 1.39 (*Limit Cardinal*)

We say that a cardinal κ is a limit cardinal if

$$(\exists \alpha \in Ord)(\kappa = \aleph_\alpha) \quad (1.35)$$

Definition 1.40 (*Strong Limit Cardinal*)

We say that an ordinal κ is a strong limit cardinal if it is a limit cardinal and

$$(\forall \alpha \in \kappa)(\mathcal{P}(\alpha) \in \kappa) \quad (1.36)$$

Definition 1.41 (*Generalised Continuum Hypothesis*)

$$\aleph_{\alpha+1} = \mathcal{P}(\aleph_\alpha) \quad (1.37)$$

If *GCH* holds (for example in Gödel's *L*, see chapter 3), the notions of a limit cardinal and a strong limit cardinal are equivalent.

⁶Except λ which is preferably used for limit ordinals.

1.2.5 Relativisation and Absoluteness

Definition 1.42 (*Relativization*)

Let M be a class, $R \subseteq M \times M$ and let $\varphi(p_1, \dots, p_n)$ be a first-order formula with no free variables besides p_1, \dots, p_n . The relativization of φ to M and R is the formula, written as $\varphi^{M,R}(p_1, \dots, p_n)$, defined in the following inductive manner:

- (i) $(x \in y)^{M,R} \leftrightarrow R(x, y)$
- (ii) $(x = y)^{M,R} \leftrightarrow x = y$
- (iii) $(\neg \varphi)^{M,R} \leftrightarrow \neg \varphi^{M,R}$
- (iv) $(\varphi \& \psi)^{M,R} \leftrightarrow \varphi^{M,R} \& \psi^{M,R}$
- (v) $(\varphi \vee \psi)^{M,R} \leftrightarrow \varphi^{M,R} \vee \psi^{M,R}$
- (vi) $(\varphi \rightarrow \psi)^{M,R} \leftrightarrow \varphi^{M,R} \rightarrow \psi^{M,R}$
- (vii) $(\exists x \varphi(x))^{M,R} \leftrightarrow (\exists x \in M) \varphi^{M,R}(x)$
- (viii) $(\forall x \varphi(x))^{M,R} \leftrightarrow (\forall x \in M) \varphi^{M,R}(x)$

When $R = \in \cap (M \times M)$, we usually write φ^M instead of $\varphi^{M,R}$. When we talk about $\varphi^M(p_1, \dots, p_n)$, it is understood that $p_1, \dots, p_n \in M$. We will also use $M \models \varphi(p_1, \dots, p_n)$ and $\varphi^M(p_1, \dots, p_n)$ interchangeably.

Definition 1.43 (*Absoluteness*) Given a transitive class M , we say a formula φ is absolute in M if for all $p_1, \dots, p_n \in M$

$$\varphi^M(p_1, \dots, p_n) \leftrightarrow \varphi(p_1, \dots, p_n) \quad (1.38)$$

Definition 1.44 (*Hierarchy of First-Order Formulas*)

A first-order formula φ is Δ_0 iff it is logically equivalent to a first-order formula φ' satisfying any of the following:

- (i) φ' contains no quantifiers
- (ii) y is a set, ψ is a Δ_0 formula, and φ' is either $(\exists x \in y)\psi(y)$ or $(\forall x \in y)\psi(y)$.
- (iii) ψ_1, ψ_2 are Δ_0 formulas and φ' is any of the following: $\psi_1 \vee \psi_2$, $\psi_1 \& \psi_2$, $\psi_1 \rightarrow \psi_2$, $\neg \psi_2$,
- (I) If a formula is Δ_0 it is also Σ_0 and Π_0
- (II) A formula φ is $\Pi_n + 1$ if it is logically equivalent to a formula φ' such that $\varphi' = \forall x \psi$ where ψ is a Σ_n -formula for any $n < \omega$.
- (III) A formula φ is $\Sigma_n + 1$ if it is logically equivalent to a formula φ' such that $\varphi' = \exists x \psi$ where ψ is a Π_n -formula for any $n < \omega$.

Note that we can use the pairing function so that for $\forall p_1, \dots, p_n \psi(p_1, \dots, p_n)$, there a logically equivalent formula of the form $\forall x \psi'(x)$.

Lemma 1.45 (Δ_0 absoluteness) *Let φ be a Δ_0 formula, then φ is absolute in any transitive class M .*

Proof. This will be proven by induction over the complexity of a given Δ_0 formula φ . Let M be an arbitrary transitive class. Suppose, that

Atomic formulas are always absolute by the definition of relativisation, see 1.42. Suppose that Δ_0 formulas ψ_1 and ψ_2 are absolute in M . Then from relativization, $(\psi_1 \& \psi_2)^M \leftrightarrow \psi_1^M \& \psi_2^M$, which is, from the induction hypothesis, equivalent to $\psi_1 \& \psi_2$. The same holds for \vee, \rightarrow, \neg .

Suppose that a Δ_0 formula ψ is absolute in M . Let y be a set and let $\varphi = (\exists x \in y)\psi(x)$. From relativization, $(\exists x\psi(x))^M \leftrightarrow (\exists x \in M)\psi^M(x)$. Since the hypotheses makes it clear that $\psi^M \leftrightarrow \psi$, we get $((\exists x \in y)\psi(x))^M \leftrightarrow (\exists x \in y \cap M)\psi(x)$, which is the equivalent of $\varphi^M \leftrightarrow \varphi$. The same applies to $\varphi = (\forall x \in y)\psi(x)$. \square

Lemma 1.46 (*Downward Absoluteness*)

Let φ be a Π_1 formula and M a transitive class. Then the following holds:

$$(\forall p_1, \dots, p_n \in M)(\varphi(p_1, \dots, p_n) \rightarrow \varphi(p_1, \dots, p_n)^M) \quad (1.39)$$

Proof. Since $\varphi(p_1, \dots, p_n)$ is Π_1 , there is a Δ_0 formula $\psi(p_1, \dots, p_n, x)$ such that $\varphi = \forall x\psi(p_1, \dots, p_n, x)$. From relativization and lemma 1.45, $\varphi^M(p_1, \dots, p_n) \leftrightarrow (\forall x \in M)\psi(p_1, \dots, p_n, x)$.

Assume that for $p_1, \dots, p_n \in M$ fixed, that $\forall x\psi(p_1, \dots, p_n, x)$ holds, but $(\forall x \in M)\psi(p_1, \dots, p_n, x)$ does not. Therefore $\exists x\neg\psi(p_1, \dots, p_n, x)$, which contradicts $\forall x\psi(p_1, \dots, p_n, x)$. \square

Lemma 1.47 (*Upward Absoluteness*)

Let φ be a Σ_1 formula and M a transitive class. Then the following holds:

$$(\forall p_1, \dots, p_n \in M)(\varphi^M(p_1, \dots, p_n) \rightarrow \varphi(p_1, \dots, p_n)) \quad (1.40)$$

Proof. Since $\varphi(p_1, \dots, p_n)$ is Σ_1 , there is a Δ_0 formula $\psi(p_1, \dots, p_n, x)$ such that $\varphi = \exists x\psi(p_1, \dots, p_n, x)$. From relativization and lemma 1.45, $\varphi^M(p_1, \dots, p_n) \leftrightarrow (\exists x \in M)\psi(p_1, \dots, p_n, x)$.

Assume that for $p_1, \dots, p_n \in M$ fixed, that $(\exists x \in M)\psi(p_1, \dots, p_n, x)$ holds, but $\exists x\psi(p_1, \dots, p_n, x)$ does not. This is an obvious contradiction. \square

1.2.6 More Functions

Definition 1.48 (*Strictly Increasing Function*)

A function $f : \text{Ord} \rightarrow \text{Ord}$ is said to be strictly increasing iff

$$\forall \alpha, \beta \in \text{Ord}(\alpha < \beta \rightarrow f(\alpha) < f(\beta)). \quad (1.41)$$

Definition 1.49 (*Continuous Function*)

A function $f : \text{Ord} \rightarrow \text{Ord}$ is said to be continuous iff

$$\alpha \text{ is limit} \rightarrow f(\alpha) = \bigcup_{\alpha < \lambda} f(\alpha). \quad (1.42)$$

Definition 1.50 (*Normal Function*)

A function $f : \text{Ord} \rightarrow \text{Ord}$ is said to be normal if it is strictly increasing and continuous.

Definition 1.51 (*Fixed Point*)

We say x is a fixed point of a function f iff $x = f(x)$.

Definition 1.52 (*Unbounded Class*)

We say a class A is unbounded if

$$\forall x(\exists y \in A)(x < y) \quad (1.43)$$

Definition 1.53 (*Limit Point*)

Given a class $x \subseteq \text{On}$, we say that $\alpha \neq \emptyset$ is a limit point of x iff

$$\alpha = \bigcup (x \cap \alpha) \quad (1.44)$$

Definition 1.54 (*Closed Class*)

We say a class $A \subseteq \text{Ord}$ is closed iff it contains all of its limit points.

Definition 1.55 (*Club set*)

For a regular uncountable cardinal κ , a set $x \subset \kappa$ is a closed unbounded subset, abbreviated as a club set, iff x is both closed and unbounded in κ .

Definition 1.56 (*Stationary set*)

For a regular uncountable cardinal κ , we say a set $A \subset \kappa$ is stationary in κ iff it intersects every club subset of κ .

1.2.7 Structure, Substructure and Embedding

Structures will be denoted $\langle M, \in, R \rangle$ where M is a domain, \in stands for the standard membership relation, it is assumed to be restricted to the domain⁷, $R \subseteq M$ is a relation on the domain. When R is not needed, we may as well only write M instead of $\langle M, \in \rangle$.

⁷To be totally correct, we should write $\langle M, \in \cap M \times M, R \rangle$

Definition 1.57 (*Elementary Embedding*)

Given the structures $\langle M_0, \in, R \rangle$, $\langle M_1, \in, R \rangle$ and a one-to-one function $j : M_0 \rightarrow M_1$, we say j is an elementary embedding of M_0 into M_1 , we write $j : M_0 \prec M_1$, when the following holds for every formula $\varphi(p_1, \dots, p_n)$ and every $p_1, \dots, p_n \in M_0$:

$$\langle M_0, \in, R \rangle \models \varphi(p_1, \dots, p_n) \leftrightarrow \langle M_1, \in, R \rangle \models \varphi(j(p_1), \dots, j(p_n)) \quad (1.45)$$

Definition 1.58 (*Elementary Substructure*)

Given the structures $\langle M_0, \in, R \rangle$, $\langle M_1, \in, R \rangle$ and a one-to-one function $j : M_0 \rightarrow M_1$ such that $j : M_0 \prec M_1$, we say that M_0 is an elementary substructure of M_1 , denoted as $M_0 \prec M_1$, iff j is an identity on M_0 . In other words

$$\langle M_0, \in, R \rangle \models \varphi(p_1, \dots, p_n) \leftrightarrow \langle M_1, \in, R \rangle \models \varphi(p_1, \dots, p_n) \quad (1.46)$$

for $p_1, \dots, p_n \in M_0$

§

2 Lévy's First-Order Reflection

2.1 Lévy's Original Paper

This section is based on Lévy's paper *Axiom Schemata of Strong Infinity in Axiomatic Set Theory* [Lévy, 1960] from 1960. It presents Lévy's general reflection principle and its equivalence to *Replacement* and *Infinity* under S^8 .

First, we should point out that set theory has changed over the last 66 years, we will now point out a few notable differences. When reading Lévy's article, one should bear in mind that while the author often speaks about a model of **ZF**, usually denoted u , it doesn't necessarily mean that there is a set u that is a model of **ZF**⁹, nowadays we are used to using the notion of universal class V in similar sense, albeit independently of a particular axiomatic theory. We will review the exact meaning of the notion of a standard complete model in a moment. The theory **ZF** is almost identical to the theory we have established in (1.21). One might be confused by the fact that Lévy treats the *Subsets* axiom, which is in fact *Specification* as a single axiom rather than a schema. He even takes the conjunction of all axioms of **ZF** and treats it like a formula. This is possible because the

⁸See definition (1.20).

⁹This is indeed impossible to prove in **ZF** due to Gödel's Incompleteness.

underlying logic calculus is different. Lévy works with set theories formulated in the *non-simple applied first order functional calculus*, see Chapter IV in [Church, 1996] for details. For now, we only need to know that the calculus contains a substitution rule for functional variables. This way, *Subsets* is de facto a schema even though it sometimes treated as a single formula. It should also be noted that the logical connectives look different. The usual symbol for universal quantifier does not appear, $\forall x\varphi(x)$ would be written as $(x)\varphi(x)$. The symbol for negation is " \sim ", implication is written as " \supset " and equivalence is " \equiv ". We will use standard notation with " \neg ", " \rightarrow " and " \leftrightarrow " respectively when presenting Lévy's results.

This subsection uses ZF instead of the usual ZFC as the underlying theory.

Definition 2.1 (*Standard Complete Model of a Set Theory*)

Let \mathbf{Q} be an arbitrary set theory given. We say that u is a standard complete model of \mathbf{Q} , which is usually written as $\text{Scm}^{\mathbf{Q}}(u)$, iff

- (i) $(\forall \sigma \in \mathbf{Q})(u \models \sigma)$
- (ii) $\forall y(y \in u \rightarrow y \subset u)$

Definition 2.2 (*Inaccessible Cardinal With Respect to \mathbf{Q}*)

Let \mathbf{Q} be an arbitrary axiomatic first-order set theory. We say that a cardinal κ is inaccessible with respect to \mathbf{Q} , we write $\text{In}^{\mathbf{Q}}(\kappa)$.

$$\text{In}^{\mathbf{Q}}(\kappa) \stackrel{\text{def}}{=} \text{Scm}^{\mathbf{Q}}(V_{\kappa}). \quad (2.47)$$

Definition 2.3 (*Inaccessible Cardinal With Respect to ZF*)

When a cardinal κ is inaccessible with respect to ZF, we only say that it is inaccessible. We write $\text{In}(\kappa)$.

$$\text{In}(\kappa) \stackrel{\text{def}}{=} \text{In}^{\text{ZF}}(\kappa) \quad (2.48)$$

The above definition of inaccessibles is used because it doesn't require *Choice*.

For the definition of relativization, see (1.42). The notation used by Lévy is " $\text{Rel}(u, \varphi)$ ", we will stick to " φ^u ".

Definition 2.4 (*N*)

The following is an axiom schema of complete reflection over ZF, denoted as N :

$$\exists u(\text{Scm}^{\text{ZF}}(u) \ \& \ \forall x_1, \dots, x_n(x_1, \dots, x_n \in u \rightarrow \varphi \leftrightarrow \varphi^u)) \quad (2.49)$$

where φ is a formula which contains no free variables except for x_1, \dots, x_n .

Definition 2.5 (N_0)

The following is almost identical to axiom schema N , but with S instead of ZF . We will call it N_0 :

$$\exists u(Scm^S(u) \ \& \ \forall x_1, \dots, x_n(x_1, \dots, x_n \in u \rightarrow \varphi \leftrightarrow \varphi^u)) \quad (2.50)$$

where φ is a formula which contains no free variables except for x_1, \dots, x_n .

Let S be an axiomatic set theory defined in (1.20). We will now show that in S , N_0 implies both *Replacement* and *Infinity*.

Let N_0 be defined as in (2.5), for *Infinity* see (1.12).

Theorem 2.6 In S , the schema N_0 implies *Infinity*.

Proof. Let $\varphi = \forall x \exists y(y = x \cup \{x\})$. This clearly holds in S because given any set x , we can always obtain the set $x \cup \{x\}$ via *Powerset* and *Specification*. From N_0 , there then exists a set u such that φ^u holds. This u satisfies the conditions required by *Infinity*, so we're done. \square

Theorem 2.7 In S , the schema N_0 implies *Replacement*.
TODO jedno nebo druhe

$$S + N_0 \vdash \text{Replacement} \quad (2.51)$$

Proof. Let $\varphi(x, y, p_1, \dots, p_n)$ be a formula with no free variables except x, y, p_1, \dots, p_n for an arbitrary natural number n .

$$\begin{aligned} \chi = & \forall x, y, z(\varphi(x, y, p_1, \dots, p_n) \ \& \ \varphi(x, z, p_1, \dots, p_n) \rightarrow y = z) \\ & \rightarrow \forall x \exists y \forall z(z \in y \leftrightarrow \exists q(q \in x \ \& \ \varphi(q, z, p_1, \dots, p_n))) \end{aligned} \quad (2.52)$$

Let χ be an instance of *Replacement* schema for the above φ . Let the following formulas be instances of the N_0 schema for formulas φ , $\exists y \varphi$, χ and $\forall x, p_1, \dots, p_n \chi$ respectively:

- (i) $x, y, p_1, \dots, p_n \in u \rightarrow (\varphi \leftrightarrow \varphi^u)$
- (ii) $x, p_1, \dots, p_n \in u \rightarrow (\exists y \varphi \leftrightarrow (\exists y \varphi)^u)$
- (iii) $x, p_1, \dots, p_n \in u \rightarrow (\chi \leftrightarrow \chi^u)$
- (iv) $\forall x, p_1, \dots, p_n(\chi \leftrightarrow (\forall x, p_1, \dots, p_n \chi)^u)$

From relativization, we also know that $(\exists y \varphi)^u$ is equivalent to $(\exists y \in u) \varphi^u$. Therefore (ii) is equivalent to

$$x, p_1, \dots, p_n \in u \rightarrow (\exists y \in u) \varphi^u. \quad (2.53)$$

If φ is a function¹⁰, then for every $x \in u$, which is also $x \subset u$ by the transitivity of $Scm^S(u)$, it maps elements of x onto u . From the axiom scheme of comprehension¹¹, we can find y , a set of all images of elements of x . That gives us $x, p_1, \dots, p_n \in u \rightarrow \chi$. By (iii) we get $x, p_1, \dots, p_n \in u \rightarrow \chi^u$, the universal closure of this formula is $(\forall x, p_1, \dots, p_n \chi)^u$, which together with (iv) yields $\forall x, p_1, \dots, p_n \chi$. Via universal instantiation, we end up with χ . We have inferred replacement for a given arbitrary formula. \square

What we have just proven is just a single theorem from the above mentioned article by Lévy, we will introduce other interesting propositions, mostly related to the existence of large cardinals, later in their appropriate context in chapter 3.

2.2 Contemporary Restatement

We will now a theorem that is referred to as Lévy's Reflection in contemporary set theory. The only difference is that while Lévy reflects φ from V to a set u which is a *standard complete model of S*, we say that there is a V_α for a limit α that reflects φ . Those two conditions are equivalent due to lemma (2.12).

Lemma 2.8 *Let $\varphi_1, \dots, \varphi_n$ be first-order formulas in the language of set theory, all with m free variables¹².*

- (i) *For each set M_0 there is such set M that $M_0 \subset M$ and the following holds for every i , $1 \leq i \leq n$:*

$$\exists x \varphi_i(p_1, \dots, p_{m-1}, x) \rightarrow (\exists x \in M) \varphi_i(p_1, \dots, p_{m-1}, x) \quad (2.54)$$

for every $p_1, \dots, p_{m-1} \in M$.

- (ii) *Furthermore, there is an ordinal α such that $M_0 \subset V_\alpha$ and the following holds for each i , $1 \leq i \leq n$:*

$$\exists x \varphi_i(p_1, \dots, p_{m-1}, x) \rightarrow (\exists x \in V_\alpha) \varphi_i(p_1, \dots, p_{m-1}, x) \quad (2.55)$$

for every $p_1, \dots, p_{m-1} \in M$.

- (iii) *Assuming Choice, there is M , $M_0 \subset M$ such that (2.83) holds for every M , $i \leq n$ and $|M| \leq |M_0| \cdot \aleph_0$.*

¹⁰See definition (1.14)

¹¹Lévy uses its equivalent, axiom of subsets

¹²For formulas with a different number of free variables, take for m the highest number of parameters among those formulas. Add spare parameters to every formula that has less than m parameters in a way that preserves the last parameter, which we will denote x . E.g. let φ'_i be the a formula with k parameters, $k < m$. Let us set $\varphi_i(p_1, \dots, p_{m-1}, x) \stackrel{\text{def}}{=} \varphi'_i(p_1, \dots, p_{k-1}, x)$, notice that the parameters p_k, \dots, p_{m-1} are not used.

Proof. We will simultaneously prove statements (i) and (ii), denoting M^T the transitive set required by part (ii). Steps in the construction of M^T that are not explicitly included are equivalent to steps for M .

Let us first define an operation $H_i(p_1, \dots, p_{m-1})$ that yields the set of x 's with minimal rank¹³ satisfying $\varphi_i(p_1, \dots, p_{m-1}, x)$ for p_1, \dots, p_{m-1} and for every i , $1 \leq i \leq n$.

$$H_i(p_1, \dots, p_n) = \{x \in C_i : (\forall z \in C)(\text{rank}(x) \leq \text{rank}(z))\} \quad (2.56)$$

for each $1 \leq i \leq n$, where

$$C_i = \{x : \varphi_i(p_1, \dots, p_{m-1}, x)\} \text{ for } 1 \leq i \leq n \quad (2.57)$$

Next, let's construct M from given M_0 by induction.

$$M_{i+1} = M_i \cup \bigcup_{j=0}^n \{H_j(p_1, \dots, p_{m-1}) : p_1, \dots, p_{m-1} \in M_i\} \quad (2.58)$$

In other words, in each step we include into the construction the elements satisfying $\varphi(p_1, \dots, p_{m-1}, x)$ for p_1, \dots, p_{m-1} from the previous step. For statement (ii), this is the only part that differs from (i). To end up with a transitive M , we need to extend every step to its transitive closure transitive closure of M_{i+1} from (i). In other words, let γ be the smallest ordinal such that

$$(M_i^T \cup \bigcup_{j=0}^n \{\bigcup \{H_j(p_1, \dots, p_{m-1}) : p_1, \dots, p_{m-1} \in M_i\}\}) \subset V_\gamma \quad (2.59)$$

Then the incremental step is

$$M_{i+1}^T = V_\gamma \quad (2.60)$$

and the final M is obtained by joining the previous steps.

$$M = \bigcup_{i=0}^{\infty} M_i, \quad M^T = \bigcup_{i=0}^{\infty} M_i^T = V_\alpha \quad (2.61)$$

We have yet to finish part (iii). Let's try to construct a set M' that satisfies the same conditions like M but is kept as small as possible. Assuming the Axiom of Choice, we can modify the process so that the cardinality

¹³Rank is defined in (1.31)

of M' is at most $|M_0| \cdot \aleph_0$. Note that the size of M in the previous construction is determined by the size of M_0 and, most importantly, by the size of $H_i(p_1, \dots, p_{m-1})$ for every i , $1 \leq i \leq n$ in individual iterations of the construction. Since (i) only ensures the existence of an x that satisfies $\varphi_i(p_1, \dots, p_{m-1}, x)$ for any i , $1 \leq i \leq n$, we only need to add one x for every set of parameters but $H_i(u_1, \dots, u_{m-1})$ can be arbitrarily large. Let F be a choice function on $\mathcal{P}(M')$. Also let $h_i(p_1, \dots, p_{m-1}) = F(H_i(p_1, \dots, p_{m-1}))$ for i , where $1 \leq i \leq n$, which means that h is a function that outputs an x that satisfies $\varphi_i(p_1, \dots, p_{m-1}, x)$ for i such that $1 \leq i \leq n$ and has minimal rank among all such sets. The induction step needs to be redefined to

$$M'_{i+1} = M'_i \cup \bigcup_{j=0}^n \{H_j(p_1, \dots, p_{m-1}) : p_1, \dots, p_{m-1} \in M'_i\} \quad (2.62)$$

This way, the amount of elements added to M'_{i+1} in each step of the construction is the same as the amount of m -tuples of parameters that yielded elements not included in M'_i . It is easy to see that if M_0 is finite, M' is countable because it was constructed as a countable union of at most countable sets. If M_0 is countable or larger, the cardinality of M' is equal to the cardinality of M_0 .¹⁴ Therefore $|M'| \leq |M_0| \cdot \aleph_0$ \square

Theorem 2.9 (*Lévy's first-order reflection theorem*)

Let $\varphi(p_1, \dots, p_n)$ be a first-order formula.

- (i) For every set M_0 there exists M such that $M_0 \subset M$ and the following holds:

$$\varphi^M(p_1, \dots, p_n) \leftrightarrow \varphi(p_1, \dots, p_n) \quad (2.63)$$

for every $p_1, \dots, p_n \in M$.

- (ii) For every set M_0 there is a transitive set M , $M_0 \subset M$ such that the following holds:

$$\varphi^M(p_1, \dots, p_n) \leftrightarrow \varphi(p_1, \dots, p_n) \quad (2.64)$$

for every $p_1, \dots, p_n \in M$.

- (iii) For every set M_0 there is α such that $M_0 \subset V_\alpha$ and the following holds:

$$\varphi^{V_\alpha}(p_1, \dots, p_n) \leftrightarrow \varphi(p_1, \dots, p_n) \quad (2.65)$$

for every $p_1, \dots, p_n \in M$.

¹⁴It can not be smaller because $|M'_{i+1}| \geq |M'_i|$ for every i . It may not be significantly larger because the maximum of elements added is the number of n -tuples in M'_i , which is of the same cardinality is M'_i .

(iv) Assuming Choice, for every set M_0 there is M such that $M_0 \subset M$ and $|M| \leq |M_0| \cdot \aleph_0$ and the following holds:

$$\varphi^M(p_1, \dots, p_n) \leftrightarrow \varphi(p_1, \dots, p_n) \quad (2.66)$$

for every $p_1, \dots, p_n \in M$.

Proof.

Let's now prove (i) for given φ via induction by complexity. We can safely assume that φ contains no quantifiers besides " \exists " and no logical connectives other than " \neg " and " $\&$ ". Let $\varphi_1, \dots, \varphi_n$ be all subformulas of φ . Then there is a set M , obtained by the means of lemma 2.8, for all of the formulas $\varphi_1, \dots, \varphi_n$.

Let's first consider atomic formulas in the form of either $x_1 = x_2$ or $x_1 \in x_2$. It is clear from relativisation¹⁵ that 2.63 holds for both cases.

$$(i) \quad (x_1 = x_2)^M \leftrightarrow (x_1 = x_2)$$

$$(ii) \quad (x_1 \in x_2)^M \leftrightarrow (x_1 \in x_2)$$

We now want to verify the inductive step. First, take $\varphi = \neg\varphi'$. From the relativization, we get

$$(\neg\varphi')^M \leftrightarrow \neg(\varphi'^M) \quad (2.67)$$

Because the induction hypothesis tell us that $\varphi'^M \leftrightarrow \varphi'$, the following holds:

$$(\neg\varphi')^M \leftrightarrow \neg(\varphi'^M) \leftrightarrow \neg\varphi' \quad (2.68)$$

The same holds for $\varphi = \varphi_1 \& \varphi_2$. From the induction hypothesis, we know that $\varphi_1^M \leftrightarrow \varphi_1$ and $\varphi_2^M \leftrightarrow \varphi_2$, which together with relativization for formulas in the form of $\varphi_1 \& \varphi_2$ gives us

$$(\varphi_1 \& \varphi_2)^M \leftrightarrow \varphi_1^M \& \varphi_2^M \leftrightarrow \varphi_1 \& \varphi_2 \quad (2.69)$$

Let's now examine the case when, from the induction hypothesis, M reflects $\varphi'(p_1, \dots, p_n, x)$ and we are interested in $\varphi = \exists x \varphi'(p_1, \dots, p_n, x)$. The induction hypothesis tells us that

$$\varphi'^M(p_1, \dots, p_n, x) \leftrightarrow \varphi'(p_1, \dots, p_n, x) \quad (2.70)$$

¹⁵See 1.42. This only holds for relativization to M, \in , not M, E for an arbitrary relation E .

so, together with above lemma 2.8, the following holds:

$$\begin{aligned}
& \varphi(p_1, \dots, p_n, x) \\
& \leftrightarrow \exists x \varphi'(p_1, \dots, p_n, x) \\
& \leftrightarrow (\exists x \in M) \varphi'(p_1, \dots, p_n, x) \\
& \leftrightarrow (\exists x \in M) \varphi'^M(p_1, \dots, p_n, x) \\
& \leftrightarrow (\exists x \varphi'(p_1, \dots, p_n, x))^M \\
& \leftrightarrow \varphi^M(p_1, \dots, p_n, x)
\end{aligned} \tag{2.71}$$

Which is what we wanted to prove for part (i).

We now need to verify that the same holds for any finite number of formulas $\varphi_1, \dots, \varphi_n$. This has in fact been already done since lemma 2.8 gives us a set M for any (finite) amount of formulas and given M_0 . We can therefore find a set M for the union of all of their subformulas. When we obtain such M , it should be clear that it also reflects every formula in $\varphi_1, \dots, \varphi_n$.

Since V_α is a transitive set, by proving (iii) we also satisfy (ii). To do so, we only need to look at part (ii) of lemma 2.8. All of the above proof also holds for $M = V_\alpha$.

To finish part (iv), we take M of size $\leq |M_0| \cdot \aleph_0$, which exists due to part (iii) of lemma 2.8, the rest being identical. \square

Let S be a set theory defined in 1.20, for ZFC see 1.22.

The two following lemmas are based on [Drake, 1974][Chapter 3, Theorem 1.2].

Lemma 2.10 *Iff M is a transitive set, then $M \models$ Extensionality.*

Proof. Given a transitive set M , we want to show that the following holds.

$$M \models \forall x, y (x = y \leftrightarrow \forall z (z \in x \leftrightarrow z \in y)) \tag{2.72}$$

Given arbitrary sets $x, y \in M$, we want to prove

$$M \models (x = y \leftrightarrow \forall z (z \in x \leftrightarrow z \in y)) \tag{2.73}$$

This is equivalent to

$$M \models x = y \text{ iff } M \models \forall z (z \in x \leftrightarrow z \in y) \tag{2.74}$$

Which is the same as

$$x = y \text{ iff } M \models \forall z(z \in x \leftrightarrow z \in y) \quad (2.75)$$

So all elements of x are also elements of y in M , and vice versa. Because M is transitive, all elements of x and y are in M , so $M \models \forall z(z \in x \leftrightarrow z \in y)$ holds iff x and y contain the same elements and are therefore equal. \square

Lemma 2.11 *If M is a transitive set, then $M \models \text{Foundation}$.*

Proof. We want to prove

$$M \models \forall x(x \neq \emptyset \rightarrow (\exists y \in x)(x \cap y = \emptyset)) \quad (2.76)$$

Given an arbitrary $x \in M$ such that $x \neq \emptyset$, we want to prove

$$M \models (\exists y \in x)(x \cap y = \emptyset) \quad (2.77)$$

Because M is transitive, every element of x is an element of M . Take for y the element of x with the lowest rank. It should be clear that there is no $z \in y$ such that $z \in x$, because then $\text{rank}(z) < \text{rank}(y)$, which is a contradiction. \square

Let S be a set theory as defined in (1.20).

Lemma 2.12 *The following holds for every λ .*

$$"\lambda \text{ is a limit ordinal}" \rightarrow V_\lambda \models S \quad (2.78)$$

Proof. Given an arbitrary limit ordinal λ , we will verify the axioms of S one by one.

- (i) *The existence of a set* comes from the fact that V_λ is a non-empty set because limit ordinal is defined as non-zero.
- (ii) *Extensionality* holds from ??
- (iii) *Foundation* holds from ??
- (iv) *Union*:
(see (1.7))

$$\forall x \exists y \forall z (z \in y \leftrightarrow \exists q (z \in q \ \& \ q \in x)) \quad (2.79)$$

Given any $x \in u$, we want verify that $y = \bigcup x$ is also in u . Note that $y = \bigcup x$ is also Δ_0 .

$$y = \bigcup x \leftrightarrow (\forall z \in y)(\exists q \in x) z \in q \ \& \ (\forall z \in x)(\forall q \in z) q \in y \quad (2.80)$$

So by lemma (1.45)

$$y = \bigcup x \leftrightarrow (y = \bigcup x)^u \quad (2.81)$$

- (v) *Pairing*:
(see (1.6))

$$\forall x, y \exists z \forall q (q \in z \leftrightarrow q = x \vee q = y) \quad (2.82)$$

Given two sets $x, y \in u$, we want to show that z , defined as $z = \{x, y\}$, is also an element of u .

$$z = \{x, y\} \leftrightarrow x \in z \ \& \ y \in z \ \& \ (\forall q \in z)(q = x \vee q = y) \quad (2.83)$$

So $(z = \{x, y\})$ is Δ_0 , and thus by lemma (1.45) it holds that

$$z = \{x, y\} \leftrightarrow (z = \{x, y\})^u \quad (2.84)$$

- (vi) *Powerset*:
(see (1.11))

$$\forall x \exists y \forall z (z \subseteq x \leftrightarrow z \in y) \quad (2.85)$$

Given $x \in u$, we want to make sure that $\mathcal{P}(x) \in u$. Let φ denote the formula $y \in \mathcal{P}(x) \leftrightarrow y \subset x$. We know that $y \subset x$ is Δ_0 according to definition (1.4). We also know that given x , φ holds for every y due to the definition of $\mathcal{P}(x)$. That means that $\varphi \leftrightarrow \varphi^u$ and therefore we can conclude that $u \models \varphi$.

- (vii) *Specification*:

Given a first-order formula φ , we want to show the following

$$u \models \forall x \forall p_1, \dots, p_n, \exists y \forall z (z \in y \leftrightarrow z \in x \ \& \ \varphi(z, p_1, \dots, p_n)) \quad (2.86)$$

Given any x along with parameters p_1, \dots, p_n , we set $y = \{z \in x : \varphi^u(z, p_1, \dots, p_n)\}$. From transitivity of u and the fact that $y \subset x$ and $x \in u$, we can conclude that $y \in u$, so $u \models \forall z (z \in y \leftrightarrow z \in x \ \& \ \varphi(z, p_1, \dots, p_n))$. □

Let *Infinity* and *Replacement* be as defined in 1.12 and 1.18 respectively.

Definition 2.13 (*First-Order Reflection Schema*)

For each set M_0 there is such set M that $M_0 \subset M$ and the following holds for every i , $1 \leq i \leq n$:

$$\exists x \varphi_i(p_1, \dots, p_{m-1}, x) \rightarrow (\exists x \in M) \varphi_i(p_1, \dots, p_{m-1}, x) \quad (2.87)$$

for every $p_1, \dots, p_{m-1} \in M$.

Theorem 2.14 Reflection_1 is equivalent to *Infinity* & *Replacement* under S.

Proof. Since 2.9 already gives us one side of the implication, we are only interested in showing the converse which we shall do in two parts:

Reflection₁ → Infinity From *Reflection₁*, we know that for any first-order formula φ and a set M_0 , there is a M such that $M_0 \subseteq M$ and $\varphi^M \leftrightarrow \varphi$. Let's pick *Powerset* for φ , then by *Reflection₁* there is a set that satisfies *Powerset*, ergo there is a strong limit cardinal, which in turn satisfies *Infinity*.

Reflection → Replacement

Given a formula $\varphi(x, y, p_1, \dots, p_n)$, we can suppose that it is reflected in any M ¹⁶ What we want to obtain is the following:

$$\begin{aligned} \forall x, y, z (\varphi(x, y, p_1, \dots, p_n) \ \& \ \varphi(x, z, p_1, \dots, p_n) \rightarrow y = z) \rightarrow \\ \rightarrow \forall X \exists Y \forall y (y \in Y \leftrightarrow \exists x (\varphi(x, y, p_1, \dots, p_n) \ \& \ x \in X)) \end{aligned} \quad (2.88)$$

We do also know that $x, y \in M$, in other words for every X , $Y = \{y \mid \varphi(x, y, p_1, \dots, p_n)\}$ and we know that $X \subset M$ and $Y \subset M$, which, together with the specification schema implies that Y , the image of X over φ , is a set. \square

We have shown that *Reflection* for first-order formulas, *Reflection₁* is a theorem of **ZF**, which means that it won't yield us any large cardinals. We have also shown that it can be used instead of the *Infinity* and *Replacement* scheme, but **ZF** + *Reflection₁* is a conservative extension of **ZF**. Besides being a starting point for more general and powerful statements, it can be used to show that **ZF** is not finitely axiomatizable. That follows from the fact that *Reflection* gives a model to any finite number of (consistent) formulas. So if $\varphi_1, \dots, \varphi_n$ for any finite n would be the axioms of **ZF**, *Reflection* would always contain a model of itself, which would in turn contradict the Second Gödel's Theorem¹⁷. Notice that, in a way, reflection is complementary to compactness. Compactness argues that given a set of sentences, if every finite subset yields a model, so does the whole set. Reflection, on the other hand, says that while the whole set has no model in the underlying theory, every finite subset does have one.

Also, notice how reflection can be used in ways similar to upward Löwenheim–Skolem theorem. Since Reflection extends any set M_0 into a model of given formulas $\varphi_1, \dots, \varphi_n$, we can choose the lower bound of the size of M by appropriately choosing M_0 .

In the next section, we will try to generalize *Reflection* in a way that transcends **ZF** and finally yields some large cardinals.

¹⁶Which means that for $x, y, p_1, \dots, p_n \in M$, $\varphi^M(x, y, p_1, \dots, p_n) \leftrightarrow \varphi(x, y, p_1, \dots, p_n)$.

¹⁷See chapter ?? for further details.

3 Conclusion

References

- [Church, 1996] Church, A. (1996). *Introduction to Mathematical Logic*. Annals of Mathematics Studies. Princeton University Press.
- [Drake, 1974] Drake, F. (1974). *Set theory. An introduction to large cardinals*. Studies in Logic and the Foundations of Mathematics, Volume 76. NH.
- [Jech, 2006] Jech, T. (2006). *Set theory*. Springer monographs in mathematics. Springer, the 3rd millennium ed., rev. and expanded edition.
- [Lévy, 1960] Lévy, A. (1960). Axiom schemata of strong infinity in axiomatic set theory. *Pacific Journal of Mathematics*, 10.
- [Wang, 1997] Wang, H. (1997). *"A Logical Journey: From Gödel to Philosophy"*. A Bradford Book.