

Topological Optimization Using the SIMP Method

Mikal Nelson

University of Kansas
Department of Mathematics
mikal.nelson@ku.edu

M.A. Thesis Defense
July 26th, 2021
Zoom



Department of Mathematics



Outline

- ▶ Orthogonal Polynomials
- ▶ Chebyshev Polynomials
- ▶ Chebyshev Expansion
- ▶ Using Kernel Polynomials
- ▶ Application: Calculating the Density of States

Orthogonal Polynomials

Scalar product on $[a, b]$:

$$\langle f|g \rangle = \int_a^b w(x) f(x) g(x) \, dx$$

Given a scalar product, we get a set of polynomials, p_n , which satisfy the orthogonality relation

$$\langle p_n | p_m \rangle = \delta_{n,m} \langle p_n | p_n \rangle .$$

This allows for an expansion of any given function $f(x)$ in terms of the $p_n(x)$:

$$f(x) = \sum_{n=0}^{\infty} \alpha_n p_n(x) \quad \text{with} \quad \alpha_n = \frac{\langle p_n | f \rangle}{\langle p_n | p_n \rangle}$$

Chebyshev Polynomials

I will focus on Chebyshev polynomials of the first kind, T_n :

Defined on interval $[-1, 1]$ with weight function

$$w(x) = \left(\pi\sqrt{1-x^2}\right)^{-1}.$$

$$T_n = \cos(n \arccos(x))$$

Recursively defined:

$$T_0(x) = 1, \quad T_{-1}(x) = T_1(x) = x,$$

$$T_{m+1}(x) = 2xT_m(x) - T_{m-1}(x).$$

The polynomials also follow the relation

$$2T_m(x)T_n(x) = T_{m+n}(x) + T_{m-n}(x).$$

Chebyshev Polynomials have some particular advantages which make them ideal for use in orthogonal polynomial expansions.

Chebyshev Expansion

The expansion of $f(x)$ in terms of Chebyshev polynomials

$$f(x) = \sum_{n=0}^{\infty} \frac{\langle f|T_n \rangle_1}{\langle T_n|T_n \rangle_1} T_n(x)$$

To make this a easier to compute, we rearrange:

$$f(x) = \frac{1}{\pi\sqrt{1-x^2}} \left[\mu_0 + 2 \sum_{n=1}^{\infty} \mu_n T_n(x) \right], \quad \mu_n = \int_{-1}^1 f(x) T_n(x) dx.$$

However, calculating the moments μ_n can be quite computationally expensive.

Motivation for KPM: Gibbs Oscillations

In practice, we cannot compute an infinite series, so we need to truncate:

$$f(x) \approx \frac{1}{\pi \sqrt{1-x^2}} \left[\mu_0 + 2 \sum_{n=1}^{N-1} \mu_n T_n(x) \right]$$

Gibbs Oscillations:

Kernel Polynomials

To lessen the impact of the Gibbs Oscillations, we introduce a “damping function” in the form of a kernel polynomial, g_n :

$$f_{\text{KPM}}(x) = \frac{1}{\pi\sqrt{1-x^2}} \left[g_0\mu_0 + 2 \sum_{n=1}^{N-1} g_n\mu_n T_n(x) \right]$$

Application: Density of States Calculations

Density of States: How many energy states exist at a given energy E .

$$\rho(E) = \frac{1}{N} \sum_{k=0}^{N-1} \delta(E - E_k)$$

Scale E down to $[-1, 1]$:

$$E = aX + b$$

We can approximate $\rho(E)$ using KPM:

$$\rho(E) \approx \frac{1}{\pi\sqrt{1-x^2}} \left[g_0\mu_0 + 2 \sum_{n=1}^{N-1} g_n\mu_n T_n(X) \right]$$

$$\mu_n \approx \frac{1}{N_r} \sum_r \langle r | T_n(X) | r \rangle$$

Disilicon Si₂ Density of States

Twisted Graphene Density of States

Topics of Further Exploration

- ▶ Effects of the number of random vectors (R) used in calculating the moments.
- ▶ Optimal resolution values (N) for various Hamiltonians.
- ▶ More in-depth comparisons of various kernels.

Acknowledgments

Big thanks to Professor Cazeaux for helping me to get my program to work.

Also thanks to Aaron for talking through some aspects of the programming with me.

Bibliography



Stephen P. Boyd and Lieven Vandenberghe.
Convex Optimization.
Cambridge Univ. Pr., 2004.