

Applied Mathematics (4320001) - Summer 2022 Solution

Milav Dabgar

August 23, 2022

Question 1 [14 marks]

Fill in the blanks using appropriate choice from the given options.

Question 1.1 [1 marks]

If $A = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}$ then $A^2 = \dots\dots\dots$ Answer: (c) $\begin{bmatrix} 7 & 10 \\ 15 & 22 \end{bmatrix}$

Solution

$$A^2 = A \times A = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix} \times \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}$$
$$A^2 = \begin{bmatrix} 1(1) + 2(3) & 1(2) + 2(4) \\ 3(1) + 4(3) & 3(2) + 4(4) \end{bmatrix} = \begin{bmatrix} 7 & 10 \\ 15 & 22 \end{bmatrix}$$

Question 1.2 [1 marks]

If $A = \begin{bmatrix} 1 & 3 \\ 4 & -2 \end{bmatrix}$ then $2A - 2I = \dots\dots\dots$ Answer: (a) $\begin{bmatrix} 0 & 6 \\ 8 & -6 \end{bmatrix}$

Solution

$$2A = 2 \begin{bmatrix} 1 & 3 \\ 4 & -2 \end{bmatrix} = \begin{bmatrix} 2 & 6 \\ 8 & -4 \end{bmatrix}$$
$$2I = 2 \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 2 & 0 \\ 0 & 2 \end{bmatrix}$$
$$2A - 2I = \begin{bmatrix} 2 & 6 \\ 8 & -4 \end{bmatrix} - \begin{bmatrix} 2 & 0 \\ 0 & 2 \end{bmatrix} = \begin{bmatrix} 0 & 6 \\ 8 & -6 \end{bmatrix}$$

Question 1.3 [1 marks]

If $A = \begin{bmatrix} -8 & -6 \\ 3 & 4 \end{bmatrix}$ then $\text{Adj } A = \dots\dots\dots$ Answer: (a) $\begin{bmatrix} 4 & 6 \\ -3 & -8 \end{bmatrix}$

Solution

For a 2×2 matrix $\begin{bmatrix} a & b \\ c & d \end{bmatrix}$, $\text{Adj } A = \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}$

$$\text{Adj } A = \begin{bmatrix} 4 & 6 \\ -3 & -8 \end{bmatrix}$$

Question 1.4 [1 marks]

Order of the matrix $\begin{bmatrix} 5 & 2 & 20 & 41 & 0 \\ 15 & 4 & 30 & 40 & 1 \\ 25 & 6 & 40 & 39 & 2 \\ 35 & 8 & 50 & 38 & 3 \end{bmatrix}$ is Answer: (b) 4×5

Solution

The matrix has 4 rows and 5 columns, so the order is 4×5 .

Question 1.5 [1 marks]

$\frac{d}{dx}(\cos^2 x + \sin^2 x) = \dots\dots\dots$ Answer: (d) 0

Solution

Since $\cos^2 x + \sin^2 x = 1$ (trigonometric identity)
 $\frac{d}{dx}(1) = 0$

Question 1.6 [1 marks]

If $f(x) = \log x$ then $f'(1) = \dots\dots\dots$ Answer: (a) 1

Solution

$$f(x) = \log x \implies f'(x) = \frac{1}{x}$$

$$f'(1) = \frac{1}{1} = 1$$

Question 1.7 [1 marks]

If $x^2 + y^2 = a^2$ then $\frac{dy}{dx} = \dots\dots\dots$ Answer: (b) $-\frac{x}{y}$

Solution

Differentiating both sides with respect to x : $2x + 2y \frac{dy}{dx} = 0 \implies \frac{dy}{dx} = -\frac{x}{y}$

Question 1.8 [1 marks]

$\int x^2 dx = \dots\dots\dots$ Answer: (b) $\frac{x^3}{3}$

Solution

$$\int x^2 dx = \frac{x^{2+1}}{2+1} + c = \frac{x^3}{3} + c$$

Question 1.9 [1 marks]

$\int e^{x \log a} dx = \dots\dots\dots$ Answer: (d) $\frac{a^x}{\log a}$

Solution

$$e^{x \log a} = a^x$$

$$\int a^x dx = \frac{a^x}{\log a} + c$$

Question 1.10 [1 marks]

$\int \cot x dx = \dots\dots\dots$ Answer: (a) $\log |\sin x|$

Solution

$$\int \cot x dx = \int \frac{\cos x}{\sin x} dx$$

Let $u = \sin x$, then $du = \cos x dx$. $\int \frac{du}{u} = \log |u| + c = \log |\sin x| + c$

Question 1.11 [1 marks]

Order of differential equation $\left(\frac{d^2 y}{dx^2}\right)^4 + \left(\frac{d^2 y}{dx^2}\right)^3 = 0$ is $\dots\dots\dots$ Answer: (b) 2

Solution

The highest derivative present is $\frac{d^2 y}{dx^2}$, which is a second derivative. Therefore, the order is 2.

Question 1.12 [1 marks]

Integrating factor of differential equation $\frac{dy}{dx} + y = 3x$ is $\dots\dots\dots$ Answer: (c) e^x

Solution

For the linear differential equation $\frac{dy}{dx} + Py = Q$, where $P = 1$. Integrating factor = $e^{\int P dx} = e^{\int 1 dx} = e^x$

Question 1.13 [1 marks]

If given data is 6, 9, 7, 3, 8, 5, 4, 8, 7 and 8 then mean is $\dots\dots\dots$ Answer: (b) 6.5

Solution

$$\text{Mean} = \frac{\text{Sum of all values}}{\text{Number of values}}$$

$$\text{Sum} = 6 + 9 + 7 + 3 + 8 + 5 + 4 + 8 + 7 + 8 = 65$$

$$\text{Number of values} = 10. \text{ Mean} = \frac{65}{10} = 6.5$$

Question 1.14 [1 marks]

The mean value of first eight natural numbers is Answer: (b) 4.5

Solution

$$\text{First eight natural numbers: } 1, 2, 3, 4, 5, 6, 7, 8. \text{ Sum} = 1 + 2 + 3 + 4 + 5 + 6 + 7 + 8 = 36 \text{ Mean} = \frac{36}{8} = 4.5$$

Question 2(a) [6 marks]

Attempt any two

Question 2(a)(1) [3 marks]

If $M = \begin{bmatrix} 2 & 3 \\ 1 & 0 \end{bmatrix}$, $N = \begin{bmatrix} 4 & 1 \\ 2 & -3 \end{bmatrix}$ then prove that $(M + N)^T = M^T + N^T$

Solution

$$M + N = \begin{bmatrix} 2 & 3 \\ 1 & 0 \end{bmatrix} + \begin{bmatrix} 4 & 1 \\ 2 & -3 \end{bmatrix} = \begin{bmatrix} 6 & 4 \\ 3 & -3 \end{bmatrix}$$

$$(M + N)^T = \begin{bmatrix} 6 & 3 \\ 4 & -3 \end{bmatrix}$$

$$M^T = \begin{bmatrix} 2 & 1 \\ 3 & 0 \end{bmatrix}, N^T = \begin{bmatrix} 4 & 2 \\ 1 & -3 \end{bmatrix}$$

$$M^T + N^T = \begin{bmatrix} 2 & 1 \\ 3 & 0 \end{bmatrix} + \begin{bmatrix} 4 & 2 \\ 1 & -3 \end{bmatrix} = \begin{bmatrix} 6 & 3 \\ 4 & -3 \end{bmatrix}$$

$$\text{Therefore, } (M + N)^T = M^T + N^T. \text{ Proved.}$$

Question 2(a)(2) [3 marks]

If $A = \begin{bmatrix} 3 & 1 \\ -1 & 2 \end{bmatrix}$ then prove that $A^2 - 5A + 7I = 0$

Solution

$$A^2 = \begin{bmatrix} 3 & 1 \\ -1 & 2 \end{bmatrix} \begin{bmatrix} 3 & 1 \\ -1 & 2 \end{bmatrix} = \begin{bmatrix} 8 & 5 \\ -5 & 3 \end{bmatrix}$$

$$\begin{aligned}
 5A &= 5 \begin{bmatrix} 3 & 1 \\ -1 & 2 \end{bmatrix} = \begin{bmatrix} 15 & 5 \\ -5 & 10 \end{bmatrix} \\
 7I &= 7 \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 7 & 0 \\ 0 & 7 \end{bmatrix} \\
 A^2 - 5A + 7I &= \begin{bmatrix} 8 & 5 \\ -5 & 3 \end{bmatrix} - \begin{bmatrix} 15 & 5 \\ -5 & 10 \end{bmatrix} + \begin{bmatrix} 7 & 0 \\ 0 & 7 \end{bmatrix} \\
 &= \begin{bmatrix} 8 - 15 + 7 & 5 - 5 + 0 \\ -5 + 5 + 0 & 3 - 10 + 7 \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix} \\
 \text{Therefore, } A^2 - 5A + 7I &= 0. \text{ Proved.}
 \end{aligned}$$

Question 2(a)(3) [3 marks]

Solve differential equation $\frac{dy}{dx} + x^2 e^{-y} = 0$

Solution

$$\begin{aligned}
 \frac{dy}{dx} &= -x^2 e^{-y} \implies e^y dy = -x^2 dx \\
 \text{Integrating both sides: } \int e^y dy &= \int -x^2 dx \\
 e^y &= -\frac{x^3}{3} + C \\
 y &= \log \left(-\frac{x^3}{3} + C \right)
 \end{aligned}$$

Question 2(b) [8 marks]

Attempt any two

Question 2(b)(1) [4 marks]

Solve $-5y + 3x = 1$, $x + 2y - 4 = 0$ using matrices

Solution

$$\begin{aligned}
 \text{Rewriting the system: } 3x - 5y &= 1 \quad x + 2y = 4 \\
 \text{In matrix form: } \begin{bmatrix} 3 & -5 \\ 1 & 2 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} &= \begin{bmatrix} 1 \\ 4 \end{bmatrix} \\
 \text{Let } A &= \begin{bmatrix} 3 & -5 \\ 1 & 2 \end{bmatrix} \\
 |A| &= 3(2) - (-5)(1) = 6 + 5 = 11 \\
 A^{-1} &= \frac{1}{11} \begin{bmatrix} 2 & 5 \\ -1 & 3 \end{bmatrix} \\
 \begin{bmatrix} x \\ y \end{bmatrix} &= A^{-1} \begin{bmatrix} 1 \\ 4 \end{bmatrix} = \frac{1}{11} \begin{bmatrix} 2 & 5 \\ -1 & 3 \end{bmatrix} \begin{bmatrix} 1 \\ 4 \end{bmatrix} \\
 &= \frac{1}{11} \begin{bmatrix} 2 + 20 \\ -1 + 12 \end{bmatrix} = \frac{1}{11} \begin{bmatrix} 22 \\ 11 \end{bmatrix} = \begin{bmatrix} 2 \\ 1 \end{bmatrix} \\
 \text{Therefore, } x &= 2, y = 1
 \end{aligned}$$

Question 2(b)(2) [4 marks]

If $A + B = \begin{bmatrix} 1 & -1 \\ 3 & 0 \end{bmatrix}$, $A - B = \begin{bmatrix} 3 & 1 \\ 1 & 4 \end{bmatrix}$ then find $(AB)^{-1}$

Solution

$$\text{Adding the equations: } 2A = \begin{bmatrix} 1 & -1 \\ 3 & 0 \end{bmatrix} + \begin{bmatrix} 3 & 1 \\ 1 & 4 \end{bmatrix} = \begin{bmatrix} 4 & 0 \\ 4 & 4 \end{bmatrix} \implies A = \begin{bmatrix} 2 & 0 \\ 2 & 2 \end{bmatrix}$$

$$\text{Subtracting: } (A + B) - (A - B) = 2B \quad 2B = \begin{bmatrix} 1 & -1 \\ 3 & 0 \end{bmatrix} - \begin{bmatrix} 3 & 1 \\ 1 & 4 \end{bmatrix} = \begin{bmatrix} -2 & -2 \\ 2 & -4 \end{bmatrix} \implies B = \begin{bmatrix} -1 & -1 \\ 1 & -2 \end{bmatrix}$$

$$AB = \begin{bmatrix} 2 & 0 \\ 2 & 2 \end{bmatrix} \begin{bmatrix} -1 & -1 \\ 1 & -2 \end{bmatrix} = \begin{bmatrix} -2 & -2 \\ 0 & -6 \end{bmatrix}$$

$$|AB| = (-2)(-6) - (-2)(0) = 12$$

$$(AB)^{-1} = \frac{1}{12} \begin{bmatrix} -6 & 2 \\ 0 & -2 \end{bmatrix} = \begin{bmatrix} -1/2 & 1/6 \\ 0 & -1/6 \end{bmatrix}$$

Question 2(b)(3) [4 marks]

If $B = \begin{bmatrix} -4 & -3 & -3 \\ 1 & 0 & 1 \\ 4 & 4 & 3 \end{bmatrix}$ then prove that $\text{adj } B = B$

Solution

For a 3×3 matrix, we need to find the cofactor matrix and then transpose it.

$$C_{11} = + \begin{vmatrix} 0 & 1 \\ 4 & 3 \end{vmatrix} = -4, \quad C_{12} = - \begin{vmatrix} 1 & 1 \\ 4 & 3 \end{vmatrix} = 1, \quad C_{13} = + \begin{vmatrix} 1 & 0 \\ 4 & 4 \end{vmatrix} = 4$$

$$C_{21} = - \begin{vmatrix} -3 & -3 \\ 4 & 3 \end{vmatrix} = -3, \quad C_{22} = + \begin{vmatrix} -4 & -3 \\ 4 & 3 \end{vmatrix} = 0, \quad C_{23} = - \begin{vmatrix} -4 & -3 \\ 4 & 4 \end{vmatrix} = 4$$

$$C_{31} = + \begin{vmatrix} -3 & -3 \\ 0 & 1 \end{vmatrix} = -3, \quad C_{32} = - \begin{vmatrix} -4 & -3 \\ 1 & 1 \end{vmatrix} = 1, \quad C_{33} = + \begin{vmatrix} -4 & -3 \\ 1 & 0 \end{vmatrix} = 3$$

$$\text{Cofactor matrix} = \begin{bmatrix} -4 & 1 & 4 \\ -3 & 0 & 4 \\ -3 & 1 & 3 \end{bmatrix}$$

$$\text{adj } B = (\text{Cofactor matrix})^T = \begin{bmatrix} -4 & -3 & -3 \\ 1 & 0 & 1 \\ 4 & 4 & 3 \end{bmatrix}$$

Since $\text{adj } B = B$. **Proved.**

Question 3(a) [6 marks]

Attempt any two

Question 3(a)(1) [3 marks]

If $y = \frac{1+\tan x}{1-\tan x}$ then find $\frac{dy}{dx}$

Solution

$$\begin{aligned} \text{Using quotient rule: } \frac{d}{dx} \left(\frac{u}{v} \right) &= \frac{v \frac{du}{dx} - u \frac{dv}{dx}}{v^2} \\ \text{Let } u &= 1 + \tan x, v = 1 - \tan x. \quad \frac{du}{dx} = \sec^2 x, \frac{dv}{dx} = -\sec^2 x \\ \frac{dy}{dx} &= \frac{(1-\tan x)(\sec^2 x) - (1+\tan x)(-\sec^2 x)}{(1-\tan x)^2} \\ &= \frac{\sec^2 x - \tan x \sec^2 x + \sec^2 x + \tan x \sec^2 x}{(1-\tan x)^2} \\ &= \frac{2\sec^2 x}{(1-\tan x)^2} \end{aligned}$$

Question 3(a)(2) [3 marks]

If $x = a(t + \sin t)$, $y = a(1 - \cos t)$ then find $\frac{dy}{dx}$

Solution

$$\begin{aligned} \frac{dx}{dt} &= a(1 + \cos t), \quad \frac{dy}{dt} = a \sin t \\ \frac{dy}{dx} &= \frac{dy/dt}{dx/dt} = \frac{a \sin t}{a(1 + \cos t)} = \frac{\sin t}{1 + \cos t} \\ \text{Using the identity } \sin t &= 2 \sin(t/2) \cos(t/2) \text{ and } 1 + \cos t = 2 \cos^2(t/2): \\ \frac{dy}{dx} &= \frac{2 \sin(t/2) \cos(t/2)}{2 \cos^2(t/2)} = \frac{\sin(t/2)}{\cos(t/2)} = \tan(t/2) \end{aligned}$$

Question 3(a)(3) [3 marks]

Evaluate $\int_0^{\pi/2} \sin x \cos x \, dx$

Solution

$$\begin{aligned} \text{Method 1: Using substitution Let } u &= \sin x, \text{ then } du = \cos x \, dx. \text{ When } x = 0, u = 0; \text{ when } x = \pi/2, u = 1. \\ \int_0^{\pi/2} \sin x \cos x \, dx &= \int_0^1 u \, du = \left[\frac{u^2}{2} \right]_0^1 = \frac{1}{2} \\ \text{Method 2: Using double angle identity } \sin x \cos x &= \frac{1}{2} \sin 2x \\ \int_0^{\pi/2} \sin x \cos x \, dx &= \frac{1}{2} \int_0^{\pi/2} \sin 2x \, dx = \frac{1}{2} \left[-\frac{\cos 2x}{2} \right]_0^{\pi/2} \\ &= -\frac{1}{4} [\cos \pi - \cos 0] = -\frac{1}{4} [-1 - 1] = \frac{1}{2} \end{aligned}$$

Question 3(b) [8 marks]

Attempt any two

Question 3(b)(1) [4 marks]

If $y = (\sin x)^{\tan x}$ then find $\frac{dy}{dx}$

Solution

Taking natural logarithm of both sides: $\ln y = \tan x \ln(\sin x)$

Differentiating both sides: $\frac{1}{y} \frac{dy}{dx} = \sec^2 x \ln(\sin x) + \tan x \cdot \frac{\cos x}{\sin x}$

$$\frac{1}{y} \frac{dy}{dx} = \sec^2 x \ln(\sin x) + \tan x \cot x$$

$$\frac{1}{y} \frac{dy}{dx} = \sec^2 x \ln(\sin x) + 1$$

$$\frac{dy}{dx} = y[\sec^2 x \ln(\sin x) + 1]$$

$$\frac{dy}{dx} = (\sin x)^{\tan x} [\sec^2 x \ln(\sin x) + 1]$$

Question 3(b)(2) [4 marks]

Find maximum and minimum value of $f(x) = 2x^3 - 3x^2 - 12x + 5$

Solution

$$f'(x) = 6x^2 - 6x - 12 = 6(x^2 - x - 2) = 6(x - 2)(x + 1)$$

For critical points: $f'(x) = 0 \implies x = 2$ or $x = -1$

$$f''(x) = 12x - 6$$

At $x = -1$: $f''(-1) = -12 - 6 = -18 < 0$ (Maximum) At $x = 2$: $f''(2) = 24 - 6 = 18 > 0$ (Minimum)

$$f(-1) = 2(-1)^3 - 3(-1)^2 - 12(-1) + 5 = -2 - 3 + 12 + 5 = 12$$

$$f(2) = 2(8) - 3(4) - 12(2) + 5 = 16 - 12 - 24 + 5 = -15$$

Maximum value = 12 at $x = -1$ Minimum value = -15 at $x = 2$

Question 3(b)(3) [4 marks]

The motion of a particle is given by $S = t^3 + 6t^2 + 3t + 5$. Find the velocity and acceleration at $t = 3$ sec.

Solution

$$\text{Position: } S = t^3 + 6t^2 + 3t + 5$$

$$\text{Velocity: } v = \frac{dS}{dt} = 3t^2 + 12t + 3$$

$$\text{Acceleration: } a = \frac{dv}{dt} = 6t + 12$$

$$\text{At } t = 3: \text{ Velocity: } v(3) = 3(9) + 12(3) + 3 = 27 + 36 + 3 = 66 \text{ units/sec}$$

$$\text{Acceleration: } a(3) = 6(3) + 12 = 18 + 12 = 30 \text{ units/sec}^2$$

Question 4(a) [6 marks]

Attempt any two

Question 4(a)(1) [3 marks]

Evaluate $\int x^2 e^x dx$

Solution

Using integration by parts twice: Let $u = x^2$, $dv = e^x dx \implies du = 2x dx$, $v = e^x$

$$\int x^2 e^x dx = x^2 e^x - \int 2x e^x dx$$

For $\int 2x e^x dx$: Let $u_1 = 2x$, $dv_1 = e^x dx \implies du_1 = 2 dx$, $v_1 = e^x$

$$\int 2x e^x dx = 2x e^x - \int 2 e^x dx = 2x e^x - 2e^x$$

Therefore: $\int x^2 e^x dx = x^2 e^x - (2xe^x - 2e^x) + C = x^2 e^x - 2xe^x + 2e^x + C = e^x(x^2 - 2x + 2) + C$

Question 4(a)(2) [3 marks]

Evaluate $\int \frac{2x+3}{(x-1)(x+2)} dx$

Solution

Using partial fractions: $\frac{2x+3}{(x-1)(x+2)} = \frac{A}{x-1} + \frac{B}{x+2}$

$$2x + 3 = A(x + 2) + B(x - 1)$$

Setting $x = 1$: $5 = 3A \implies A = \frac{5}{3}$ Setting $x = -2$: $-1 = -3B \implies B = \frac{1}{3}$

$$\begin{aligned} \int \frac{2x+3}{(x-1)(x+2)} dx &= \int \left(\frac{5/3}{x-1} + \frac{1/3}{x+2} \right) dx \\ &= \frac{5}{3} \ln|x-1| + \frac{1}{3} \ln|x+2| + C \end{aligned}$$

Question 4(a)(3) [3 marks]

Find mean using the given information

Solution

Table 1. Frequency Distribution

x_i	52	55	58	62	79
f_i	5	3	2	3	6

$$\text{Mean} = \frac{\sum f_i x_i}{\sum f_i}$$

$$\sum f_i x_i = 52(5) + 55(3) + 58(2) + 62(3) + 79(6) = 260 + 165 + 116 + 186 + 474 = 1201$$

$$\sum f_i = 5 + 3 + 2 + 3 + 6 = 19$$

$$\text{Mean} = \frac{1201}{19} = 63.21$$

Question 4(b) [8 marks]

Attempt any two

Question 4(b)(1) [4 marks]

Evaluate $\int_{-1}^1 \frac{x^5 - 6x}{x-4} dx$

Solution

First, let's perform polynomial long division: $\frac{x^5 - 6x}{x-4} = x^4 + 4x^3 + 16x^2 + 64x + 250 + \frac{1000}{x-4}$

$$\int_{-1}^1 \frac{x^5 - 6x}{x-4} dx = \int_{-1}^1 \left(x^4 + 4x^3 + 16x^2 + 64x + 250 + \frac{1000}{x-4} \right) dx$$

$$= \left[\frac{x^5}{5} + x^4 + \frac{16x^3}{3} + 32x^2 + 250x + 1000 \ln|x-4| \right]_{-1}^1$$

$$\text{At } x = 1: \frac{1}{5} + 1 + \frac{16}{3} + 32 + 250 + 1000 \ln 3 \quad \text{At } x = -1: -\frac{1}{5} + 1 - \frac{16}{3} + 32 - 250 + 1000 \ln 5$$

$$= \left(\frac{2}{5} + \frac{32}{3} + 500 + 1000 \ln \frac{3}{5} \right)$$

$$= \frac{6+160+1500}{15} + 1000 \ln \frac{3}{5} = \frac{1666}{15} + 1000 \ln \frac{3}{5}$$

Question 4(b)(2) [4 marks]

Evaluate $\int \sin 5x \sin 6x \, dx$

Solution

Using the product-to-sum formula: $\sin A \sin B = \frac{1}{2}[\cos(A - B) - \cos(A + B)]$

$$\sin 5x \sin 6x = \frac{1}{2}[\cos(5x - 6x) - \cos(5x + 6x)] = \frac{1}{2}[\cos(-x) - \cos(11x)] = \frac{1}{2}[\cos x - \cos(11x)]$$

$$\int \sin 5x \sin 6x \, dx = \frac{1}{2} \int [\cos x - \cos(11x)] \, dx$$

$$= \frac{1}{2} \left[\sin x - \frac{\sin(11x)}{11} \right] + C$$

$$= \frac{\sin x}{2} - \frac{\sin(11x)}{22} + C$$

Question 4(b)(3) [4 marks]

Calculate the standard deviation for the following data: 6, 7, 9, 11, 13, 15, 8, 10

Solution

Data: 6, 7, 8, 9, 10, 11, 13, 15 (arranged in order) $n = 8$

Step 1: Calculate Mean $\bar{x} = \frac{6+7+8+9+10+11+13+15}{8} = \frac{79}{8} = 9.875$

Step 2: Calculate deviations and their squares

Table 2. Standard Deviation Calculation

x_i	$x_i - \bar{x}$	$(x_i - \bar{x})^2$
6	-3.875	15.016
7	-2.875	8.266
8	-1.875	3.516
9	-0.875	0.766
10	0.125	0.016
11	1.125	1.266
13	3.125	9.766
15	5.125	26.266

$$\sum (x_i - \bar{x})^2 = 64.878$$

Step 3: Calculate Standard Deviation $\sigma = \sqrt{\frac{\sum (x_i - \bar{x})^2}{n}} = \sqrt{\frac{64.878}{8}} = \sqrt{8.11} = 2.85$

Standard Deviation = 2.85

Question 5(a) [6 marks]

Attempt any two

Question 5(a)(1) [3 marks]

Find the mean for the following data:

Solution**Table 3.** Data

X_i	92	93	97	98	102	104
F_i	3	2	2	3	6	4

$$\begin{aligned} \text{Mean} &= \frac{\sum f_i x_i}{\sum f_i} \\ \sum f_i x_i &= 92(3) + 93(2) + 97(2) + 98(3) + 102(6) + 104(4) = 276 + 186 + 194 + 294 + 612 + 416 = 1978 \\ \sum f_i &= 3 + 2 + 2 + 3 + 6 + 4 = 20 \\ \text{Mean} &= \frac{1978}{20} = 98.9 \end{aligned}$$

Question 5(a)(2) [3 marks]

Calculate the standard deviation for the following data: 5, 9, 8, 12, 6, 10, 6, 8

Solution

Data: 5, 6, 6, 8, 8, 9, 10, 12 (arranged in order) $n = 8$

Step 1: Calculate Mean $\bar{x} = \frac{5+6+6+8+8+9+10+12}{8} = \frac{64}{8} = 8$

Step 2: Calculate Standard Deviation

Table 4. Deviations

x_i	$x_i - \bar{x}$	$(x_i - \bar{x})^2$
5	-3	9
6	-2	4
6	-2	4
8	0	0
8	0	0
9	1	1
10	2	4
12	4	16

$$\sum (x_i - \bar{x})^2 = 38$$

$$\sigma = \sqrt{\frac{38}{8}} = \sqrt{4.75} = 2.18$$

Standard Deviation = 2.18

Question 5(a)(3) [3 marks]

Calculate the Mean for the following data: 5, 15, 25, 35, 45, 55, 65, 75, 85, 95, 75

Solution

$$n = 11$$

$$\text{Sum} = 5 + 15 + 25 + 35 + 45 + 55 + 65 + 75 + 85 + 95 + 75 = 575$$

$$\text{Mean} = \frac{575}{11} = 52.27$$

Question 5(b) [8 marks]

Attempt any two

Question 5(b)(1) [4 marks]

Solve differential equation $\frac{dy}{dx} + \frac{y}{x} = e^x$, $y(0) = 2$

Solution

This is a first-order linear differential equation of the form $\frac{dy}{dx} + Py = Q$

Here, $P = \frac{1}{x}$ and $Q = e^x$

Integrating Factor: $\mu = e^{\int P dx} = e^{\int \frac{1}{x} dx} = e^{\ln x} = x$ (for $x > 0$)

Multiplying the equation by $\mu = x$: $x \frac{dy}{dx} + y = xe^x \implies \frac{d}{dx}(xy) = xe^x$

Integrating both sides: $xy = \int xe^x dx$

Using integration by parts for $\int xe^x dx$: Let $u = x$, $dv = e^x dx \implies du = dx$, $v = e^x$

$\int xe^x dx = xe^x - \int e^x dx = xe^x - e^x = e^x(x - 1)$

Therefore: $xy = e^x(x - 1) + C$

$y = \frac{e^x(x-1)+C}{x}$

Final Answer: $y = e^x + \frac{1}{x}$ (subject to domain restrictions)

Question 5(b)(2) [4 marks]

Solve differential equation $\frac{dy}{dx} + \frac{4x}{x^2+1}y = \frac{1}{(x^2+1)^2}$

Solution

This is a first-order linear differential equation. $P = \frac{4x}{x^2+1}$, $Q = \frac{1}{(x^2+1)^2}$

Integrating Factor: $\mu = e^{\int P dx} = e^{\int \frac{4x}{x^2+1} dx}$

Let $u = x^2 + 1$, then $du = 2x dx$. $\int \frac{4x}{x^2+1} dx = 2 \int \frac{du}{u} = 2 \ln u = 2 \ln(x^2 + 1)$

$\mu = e^{2 \ln(x^2+1)} = (x^2 + 1)^2$

Multiplying the equation by μ : $(x^2 + 1)^2 \frac{dy}{dx} + 4x(x^2 + 1)y = 1$

This can be written as: $\frac{d}{dx}[y(x^2 + 1)^2] = 1$

Integrating: $y(x^2 + 1)^2 = x + C$

$y = \frac{x+C}{(x^2+1)^2}$

Question 5(b)(3) [4 marks]

Solve differential equation $\frac{dy}{dx} = \sin(x + y)$

Solution

Let $v = x + y$, then $\frac{dv}{dx} = 1 + \frac{dy}{dx} \implies \frac{dy}{dx} = \frac{dv}{dx} - 1$

Substituting into the original equation: $\frac{dv}{dx} - 1 = \sin v \implies \frac{dv}{dx} = 1 + \sin v$

Separating variables: $\frac{dv}{1+\sin v} = dx$

To integrate the left side, we use the identity: $\frac{1}{1+\sin v} = \frac{1-\sin v}{(1+\sin v)(1-\sin v)} = \frac{1-\sin v}{\cos^2 v}$

$\int \frac{dv}{1+\sin v} = \int \frac{1-\sin v}{\cos^2 v} dv = \int (\sec^2 v - \sec v \tan v) dv$
 $= \tan v - \sec v + C_1$

Therefore: $\tan(x + y) - \sec(x + y) = x + C$

Formula Cheat Sheet

- **Matrix Operations:** $(A + B)^T = A^T + B^T$, $(AB)^T = B^T A^T$, $(A^{-1})^T = (A^T)^{-1}$
- **Differentiation:** $\frac{d}{dx}[x^n] = nx^{n-1}$, $\frac{d}{dx}[\ln x] = \frac{1}{x}$, $\frac{d}{dx}[e^x] = e^x$, $\frac{d}{dx}[\sin x] = \cos x$

- **Integration:** $\int x^n dx = \frac{x^{n+1}}{n+1}$, $\int e^x dx = e^x$, $\int \sin x dx = -\cos x$
- **Differential Equations:** Linear DE $\frac{dy}{dx} + Py = Q$, IF $\mu = e^{\int P dx}$
- **Statistics:** Mean $\bar{x} = \frac{\sum f_i x_i}{\sum f_i}$, SD $\sigma = \sqrt{\frac{\sum (x_i - \bar{x})^2}{n}}$